

# Higher AVCO<sub>2</sub> and Lactate Gradient Combined with SOFA Score as a Mortality Predictor during the 6-Hours of Resuscitation of Septic Shock

Morteza Talebi Doluee<sup>1</sup>, Sahar Azam Ghasemzadeh<sup>1</sup>, Ahmad Bagheri Moghadam<sup>2</sup>, Zahra Abbasishaye<sup>1</sup>, Davood Soroosh<sup>3</sup>, Elnaz Vafadar Moradi<sup>1\*</sup>

<sup>1</sup>Department of Emergency Medicine, Faculty of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran.

<sup>2</sup>Department of Anesthesiology and Critical Care, Faculty of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran.

<sup>3</sup>Department of Forensic Medicine, Mashhad University of Medical Sciences, Mashhad, Iran.

## ARTICLE INFO

### Article history:

Received 24 June 2024

Revised 17 July 2024

Accepted 02 August 2024

### Keywords:

Sepsis;

Lactate;

Prognosis;

Intensive care unite

## ABSTRACT

**Background:** Sepsis is a severe and life-threatening condition leading to widespread inflammation and organ dysfunction. It is a medical emergency that requires immediate attention and treatment. One of the key indicators used to assess the severity and prognosis of sepsis is lactate level. Another key indicator of sepsis severity is a significant difference in the level of carbon dioxide (CO<sub>2</sub>) between veins and arteries.

**Methods:** In this study, we aimed to evaluate the differences in the venous and arterial PCO<sub>2</sub> and lactate levels during the first 6 hours of treatment of septic shock. In this prospective observational-analytical study patients with septic shock admitted to the ICU were evaluated. Sepsis is defined as patients who had 1 or more of the SIRS criteria with a possible or proven source of infection and hypotension despite appropriate fluid therapy who needed to receive vasopressors.

**Results:** Among 85 patients the mean age was 64±17 years and 48 (56%) were men. Of these patients, 15 (17%) died, of them 8 (53.33%) were male, 14 (93%) were diabetic, 11 (73.33%) were hypertensive, 11 (73.33%) had ischemic Heart disease and 9 (60%) patients had Chronic Obstructive Pulmonary Disease (COPD). The mean HR, SBP, and DBP were significantly higher in lived patients; the SOFA scores were significantly lower in these patients.

**Conclusion:** Overall, the gradient of AV PCO<sub>2</sub> and lactate clearance combined with SOFA score can be a valuable tool for clinicians in predicting mortality risk in critically ill patients and guiding treatment decisions.

## Introduction

Sepsis is a severe and life-threatening condition that occurs when the body's response to an infection becomes dysregulated, leading to widespread inflammation and organ dysfunction. It is a medical emergency that requires immediate attention and treatment [1].

One of the key indicators used to assess the severity and prognosis of sepsis is lactate level. Lactate is a byproduct of anaerobic metabolism, which occurs when there is insufficient oxygen supply to meet the energy demand of cells [2]. In sepsis, lactate levels rise due to tissue hypoperfusion, inadequate oxygen delivery, and impaired cellular metabolism [3-4]. Another key indicator of sepsis severity is a significant difference in the level of carbon dioxide (CO<sub>2</sub>) between veins and arteries [5].

The authors declare no conflicts of interest.

\*Corresponding author.

E-mail address: vafadarme@mums.ac.ir

Copyright © 2025 Tehran University of Medical Sciences. Published by Tehran University of Medical Sciences.



This work is licensed under a Creative Commons Attribution-NonCommercial 4.0 International license (<https://creativecommons.org/licenses/by-nc/4.0/>). Noncommercial uses of the work are permitted, provided the original work is properly cited.

Understanding these differences is crucial for diagnosis and managing septic patients effectively [6]. Veins and arteries have distinct roles in the circulatory system. Arteries carry oxygenated blood back to the heart [7]. In sepsis, a severe infection that can lead to organ dysfunction, both veins and arteries are affected. One of the key differences between vein and arterial CO<sub>2</sub> in sepsis is the partial pressure of carbon dioxide (PCO<sub>2</sub>) [8]. In normal conditions, arterial PCO<sub>2</sub> is slightly higher than venous PCO<sub>2</sub> due to CO<sub>2</sub> production by tissue during metabolism [9]. However, in sepsis, this relationship can change significantly and often leads to tissue hypo perfusion or inadequate blood flow to organs and tissues [10]. When tissue perfusion is compromised, cells switch from aerobic metabolism to anaerobic metabolism [11]. As a result, arterial PCO<sub>2</sub> decreases due to reduced CO<sub>2</sub> production [12]. On the other hand, venous PCO<sub>2</sub> increases in sepsis due to impaired clearance of CO<sub>2</sub> by tissues [1,4]. This accumulation of CO<sub>2</sub> leads to an increase in venous PCO<sub>2</sub> levels [13-14]. The difference between arterial and venous PCO<sub>2</sub> is known as the arteriovenous carbon dioxide gradient (AVCO<sub>2</sub> gradient) [12]. In sepsis, this gradient becomes wider than normal due to decreased arterial PCO<sub>2</sub> and increased venous PCO<sub>2</sub> [7]. Monitoring and analyzing the AVCO<sub>2</sub> gradient can provide information about tissue perfusion and the severity of sepsis [15]. Clinicians often use these differences in CO<sub>2</sub> levels to guide resuscitation efforts in septic shock [11]. In this study, we aimed to evaluate the differences in the venous and arterial PCO<sub>2</sub> and lactate levels during the first 6 hours of treatment of septic shock

## Methods

### Study design

This study is a prospective observational-analytical study on patients with septic shock admitted to the ICU of Imam Reza Hospital from 2018 to 2019. Sepsis is defined as patients who had 1 or more of the SIRS criteria with a possible or proven source of infection and hypotension despite appropriate fluid therapy who needed to receive vasopressors.

### Participants

Patients with septic shock were enrolled in this study; inclusion criteria were confirmation of septic shock based on the criteria explained and patients under 18 years of age, pregnant, known heart disease, liver disease, severe pulmonary airway obstruction, and whom with the

inability to insert a central venous catheter are excluded from the study.

Once the diagnosis was confirmed, 1 cc of arterial blood was taken as an ABG and checked lactate level, arterial carbon dioxide pressure, and a venous blood gas (VBG) sample was also taken from the central vein to for carbon dioxide pressure and oxygen pressure. Finally, the patient's outcome, daily SOFA score, length of stay in ICU, and 11-day mortality were determined.

It should be noted that blood cultures and other necessary cultures were taken. Broad-spectrum antibiotics were started in all patients in the first hour. Crystalloid was administered at the rate of 30 kg/cc in the first three hours up to mean arterial blood pressure (MAP) of more than 65.

Three hours later, fluid therapy was performed as a fluid challenge based on hemodynamic status.

### Statistical Analysis and Sample Size

Descriptive statistical methods such as mean, standard deviation, frequency, and frequency percentage were used to describe the data. The quantitative variables were analyzed by T-test due to normality of data distribution, and qualitative variables with Fisher's exact test.

The sample size was calculated at 71-patients considering the correlation coefficient between the difference in venous and arterial carbon dioxide pressure with changes in central venous oxygen pressure ( $r^2=0.55$ ), and alpha 05 0.0 and beta 0.2; to increase the power of the study we considered 85-patients [7].

## Results

Among 85 patients enrolled in this study with a mean age of  $64 \pm 17$  years 48 (56%) were men. Patient's vital signs, length of ICU- admission, lactate level, and the differentiation between the vein and arterial PCO<sub>2</sub> are summarized in Table 1.

Of these patients, 15 (17%) died, of them 8 (53.33%) were male, 14 (93%) were diabetic, 11(73.33%) were hypertensive, 11 (73.33%) had ischemic Heart disease and 9 (60%) patients had Chronic Obstructive Pulmonary Disease (COPD).

Patient's vital signs and also lactate level and AVCO<sub>2</sub> in both dead and lived groups were summarized in Table 2. The t-test showed that mean HR, SBP, and DBP were significantly higher in lived; on the other hand, mean RR was significantly lower in these patients.

In this study, we evaluated the patient's SOFA score for 13 days, in this way, we found that patients' SOFA scores were significantly lower in lived patients (Table-3).

**Table 1- Study variables.**

	Min	Max	Mean	Standard deviation
1. Patients Vital Sign				
Age(year)	37	95	64±17	17.00
Heart Rate (per minute)	36	140	112±25	25.00
SBP (mmhg)	61	133	68 ± 25	25.00
DBP (mmhg)	40	93	59 ± 16.08	16.00
RR (per minutes)	15	40	24 ± 9	9.02
O 2 saturation (%)	60	100	77 ± 12	12.00
2. Length of ICU Stay				
Length of stay at ICU (Day)	0	13	6	3
3. Laboratory findings				
Lactate at admission	1.0	3.1	1.00	0.00
Lactate after 6-hours	0.0	2.0	0.00	0.00
Lactate gradient (admission and after 6-hour)	0.02	4.10	1.00	0.00
AV PCO2 at admission	4.0	11.0	6.00	1.00
AV PCO2 after 6-hours	5.0	7.0	6.00	0.00
AV PCO2 gradient (admission and after 6-hour)	- 3.02	12.00	1.00	4.00

HR: Heart Rate; SBP: Systolic Blood Pressure; DBP: Diastolic Blood Pressure; RR: Respiratory Rate; ICU: Intensive Care units; AV PCO2: Arteria-venous PCO2

**Table 2- Patients prognosis based on study variables**

	Lived (N: 70)	Died (N:15)	P value
1. Vital Signs / Mean (Standard deviation)			
Heart Rate (per minute)	112 (27.00)	112 (14.00)	0.001
SBP (mmhg)	87 (25.00)	83 (23.00)	0.001
DBP (mmhg)	60 (16.00)	58 (13.00)	0.001
RR (per minutes)	24 (8.00)	26 (10.00)	0.001
O 2 saturation (%)	78 (12.00)	71 (11.00)	0.04
2. Length of ICU Stay/ Mean (Standard deviation)			
Length of stay at ICU (Day)	6 (3.00)	5 (3.00)	0.001
3. Laboratory findings/ Mean (Standard deviation)			
Lactate at admission	1 (0.00)	3.02 (0.04)	0.001
Lactate after 6-hours	0 (0.00)	-	-
Lactate gradient (admission and after 6-hour)	0 (0.00)	4.02 (0.04)	0.001
AV PCO2 at admission	6.01 (1.00)	9.00 (1.00)	0.001
AV PCO2 after 6-hours	6 (0.00)	-	-
AV PCO2 gradient (admission and after 6-hour)	0.00 (1.00)	10 (1.0)	0.001

**Table 3- SOFA score from the first day of admission till 13 day**

	N	Mean	Standard deviation	P value
Admission	Lived	70	3.00	0.00
	Dead	15	10.00	2.00
Day-1	Lived	69	3.00	0.00
	Dead	13	10.07	2.00
Day-2	Lived	64	3.00	1.03
	Dead	13	9.00	2.00
Day-3	Lived	58	3.00	1.00
	Dead	12	9.00	1.00
Day-4	Lived	51	3.07	1.00
	Dead	10	9.00	1.00
Day-5	Lived	48	3.08	1.00
	Dead	8	9.00	1.00
Day-6	Lived	42	2.00	1.00
	Dead	8	9.00	1.00
Day-7	Lived	35	2.00	1.00
	Dead	7	8.00	1.00
Day-8	Lived	27	2.00	1.00

	Dead	6	8.00	2.04	
Day-9	Lived	25	2.00	1.00	0.001
	Dead	5	8.00	2.00	
Day-10	Lived	19	2.00	1.00	0.001
	Dead	3	10.00	2.00	
Day-11	Lived	12	2.00	1.00	0.001
	Dead	2	10.00	2.00	
Day-12	Lived	8	2.00	2.00	0.001
	Dead	-			
Day-13	Lived	4	3.00	1.00	0.001
	Dead	-			

## Discussion

Sepsis is a systemic inflammatory response to infection after the invasion of microbial pathogens into the bloodstream, which leads to suppression of the immune system, multiple organ failure, and susceptibility to hospital infections. It is very important to predict the risk of mortality, especially in high-cost departments such as the emergency department and ICU.

Estimating the patient's mortality risk, in addition to informing the patient's relatives, can be a useful tool to evaluate the quality of emergency department services, as well as evaluate the effectiveness of the treatments; therefore, a different scoring system has been designed. The present study is designed based on predicting hospital mortality of septic patients and includes changes in central venous and arterial carbon dioxide pressure, changes in lactate clearance, as well as daily SOFA score.

We showed that the mean lactate level and its changes in the dead patients were significantly higher than survived. These findings showed the importance of the lactate level by considering it as a prognostic factor due to the emphasis of recent studies on this factor [1].

Lactate level has been used in predicting the adverse outcomes of sepsis and trauma in patients [1-3]. Studies have shown that lactate secretion of more than 32% within 6 hours after resuscitation can reduce the survival rate of patients suffering from septic shock [4]. Researchers showed that a lactate level greater than 4 L/mmol was a criterion for trauma with a worse prognosis [5]. Others have proven that patients with higher lactate levels compared to normal levels had a higher risk of death [12].

Gustavo and colleagues studied 60 septic shock patients and found that persistent high-pressure differences in arterial-venous carbon dioxide (Pv-aCO<sub>2</sub>) are associated with a worse prognosis in these patients [11]. We found similar findings to them, patients with higher AVCO<sub>2</sub> gradient were more likely to die, which was consistent with the results of the study by Gustavo et al.

In confirmation of our study, we can refer to the Huai-wu study in 2016 in China, which investigated the relationship between changes in the difference between Central venous and arterial carbon dioxide pressure (P (v-

a) CO<sub>2</sub>/C (a-v) O<sub>2</sub>) and lactate clearance in septic shock patient's prognosis. In this study, 84 septic shock patients were monitored within 8 hours of resuscitation. They showed that higher P (v-a) CO<sub>2</sub>/C (a-v) O<sub>2</sub>) is associated with higher lactate clearance [13], like us.

Mallat et al. investigated the relationship between lactate and Pv-aCO<sub>2</sub> in septic patients responding to crystalloid therapy. This study which was conducted on 80 patients showed that central venous and arterial carbon dioxide pressure combined with changes in lactate can be used as a marker to check the effectiveness of resuscitation [14]. By combining these markers, clinicians can get a more comprehensive picture of a patient's response to resuscitation interventions. Improvements in central venous and arterial CO<sub>2</sub> pressures, along with lactate clearance, can indicate successful resuscitation and tissue perfusion. Conversely, persistent abnormalities in these markers, as we showed in our study, may suggest ongoing tissue hypo perfusion need for further interventions and poor prognosis.

Zhu et al evaluated the relationship between Pv-aCO<sub>2</sub>/Ca-vO<sub>2</sub> and lactate levels and prognosis of patients with septic shock. Their study was conducted on 144 patients and showed that the ratio of changes in Pv-aCO<sub>2</sub>/Ca-vO<sub>2</sub> and lactate level is more accurate for estimating the prognosis and 11-day mortality of septic patients than each of these factors alone [15].

The results of all these studies are consistent with the results of our study and indicate the poor prognosis of patients with higher lactate gradients and differences in arterial and venous pCO<sub>2</sub>.

Higher daily SOFA scores also were accompanied by poor outcomes. Balchi et al evaluated the value of SOFA in predicting the mortality of septic and non-septic patients, which was conducted on 120 ICU-admitted patients. They showed, that when SOFA scores increase, the patient's mortality increases. Therefore, examining the patients' SOFA Score increases the power of predicting the results of septic patients [16]. While the SOFA score can be a useful tool for predicting mortality in critically ill patients, it is important to note that it is just one of many factors that should be considered when making treatment decisions. Therefore, combining the SOFA

score with the gradient of lactate clearance and Pv-aCO<sub>2</sub> may be helpful to better decisions.

Also, Sawika et al. conducted a study on 99 malignant hematologic ICU-admitted patients to investigate the effectiveness of SOFA in estimating their prognosis and analyzed the risk factors related to mortality using univariate logistic regression analysis. They found that SOFA was one of the independent risk factors of death [2]. Thakur et al. conducted a study on surgical patients with sepsis and used SOFA score and Apache- II to predict mortality. They studied 72 surgical patients with sepsis and showed that Apache- II, and SOFA scores were equally effective in the mortality assessment of surgical patients with sepsis [17].

Therefore, the predictive power of SOFA Score, like other common scoring systems such as Apache- II, has been mentioned as acceptable and excellent in most previous studies, such as ours. In a study aimed at investigating SOFA and Apache- II scores in predicting mortality in sepsis patients, Desal et al. examined SOFA scores on days 1, 3, 7, and Apache on the day of admission. Finally, they had 41% mortality, and 84% of their patients had multi-organ dysfunction syndrome (MODS), and the most affected organ was the lung. On the third day, 9.1% of patients with a SOFA score of less than 9 and 78% with a SOFA score of more than 9 had mortality; however, the Apache- II score had a slight difference in these patients. They showed that the Apache-II score was inefficient in determining the prognosis of patients and it was in line with our results [18].

## Conclusion

Based on our study, it seems that the mean lactate, the gradient of AV PCO<sub>2</sub> and lactate, and daily SOFA scores are important predictors for in-hospital death in patients with sepsis. We found that combining these three variables can provide a more accurate prediction of mortality in critically ill patients instead of using any single variable alone. This combined approach takes into account both tissue perfusion, providing a more comprehensive assessment of the septic shock patient's condition and prognosis. Overall, the gradient of AV PCO<sub>2</sub> and lactate clearance combined with SOFA score can be a valuable tool for clinicians in predicting mortality risk in critically ill patients and guiding treatment decisions.

## Strengths and limitations

This study had several strengths, including the simultaneous use of models to predict septic in-hospital mortality, as well as a new idea in the emergency department (ED) context.

Among the limitations of the current research, it can be mentioned that the research was conducted in a single

center, which reduces the generalizability of the results. Since there were few studies in this field, the possibility of comparison with other studies was limited. It should be noted that few studies in this field have been conducted in the intensive care unit, and considering that the present study was conducted in the emergency department, this issue can be one of the limitations of our study and to some extent affect the results.

## Acknowledgments

This study was supported by the Vice chancellors of Research of Mashhad University of Medical Science (grant number 960521).

## References

- [1] Batul R, Gulzar U, Qayoom O. The Outcome of Sepsis Patients Admitted to the Surgical Intensive Care Unit. *Arch Anesthesiol Crit Care*. 2023; 9(2):126-31.
- [2] Sawicka W, Owczuk R, Wujtewicz M, Wujtewicz M. The effectiveness of the APACHE II, SAPS II and SOFA prognostic scoring systems in patients with haematological malignancies in the intensive care unit. *Anestezjol Intens Ter*. 2019; 46(3):177-82.
- [3] Bahtouee M, Eghbali SS, Maleki N, Rastgou V, Motamed N. Use of the APACHE II score for the assessment of outcome and mortality prediction in an Iranian medical-surgical intensive care unit. *Arch Anesthesiol Crit Care*. 2018; 4(4):521-6.
- [4] Kumar R, Kumar N. Validation of lactate clearance at 6 h for mortality prediction in critically ill children. *Indian journal of critical care medicine: peer-reviewed, Indian J Crit Care Med*. 2016; 20(10):570.
- [5] Purohit A, Yadav A, Yadav R, Kumar S, Prasad S, Gupta D. Lactate Albumin Ratio as Predictor of 28 Days Mortality in Septic Shock Patients and its Correlation to SOFA Score for Assessment of Organ Dysfunction: A Prospective Observational Study. *Arch Anesthesiol Crit Care*. 2024; 10(2).
- [6] Guzasiah FY, Nurdin H, Muchtar F, Rum HM, Palinrunji AS. Low procalcitonin clearance correlates with mortality treated with culture-matched antibiotics in intensive care unit: A retrospective, observational study. *Bali J Anesthesiol*. 2023; 7(3):150-3.
- [7] Mohammadi H, Haghghi M. Survey relationship of mortality rate of hospitalized patients in ICU with different degrees of APACHE II. *J Guilan Univ Med Sci*. 2006; 15(59):85-90.
- [8] Seymour CW, Liu VX, Iwashyna TJ, Brunkhorst FM, Rea TD, Scherag A, Rubenfeld G, Kahn JM, Shankar-Hari M, Singer M, Deutschman CS. Assessment of clinical criteria for sepsis: for the third international consensus definitions for sepsis and septic shock (Sepsis-3). *Jama*. 2016;

- 315(8):762-74.
- [9] Rhodes LA, Erwin WC, Borasino S, Cleveland DC, Alten JA. Central venous to arterial CO<sub>2</sub> difference after cardiac surgery in infants and neonates. *J.Pediatr Crit Care*. 2017; 18(3):228-33.
- [10] Yuan S, He H, Long Y. Interpretation of venous-to-arterial carbon dioxide difference in the resuscitation of septic shock patients. *J Thorac Dis*. 2019;11(Suppl 11): S1538.
- [11] Ospina-Tascón GA, Bautista-Rincón DF, Umaña M, Tafur JD, Gutiérrez A, García AF, Bermúdez W, Granados M, Arango-Dávila C, Hernández G. Persistently high venous-to-arterial carbon dioxide differences during early resuscitation are associated with poor outcomes in septic shock. *J Crit Care*. 2013;1-10.
- [12] Pan J, Peng M, Liao C, Hu X, Wang A, Li X. Relative efficacy and safety of early lactate clearance-guided therapy resuscitation in patients with sepsis: A meta-analysis. *J Med*. 2019; 98(8): e14453.
- [13] He HW, Liu DW, Long Y, Wang XT. High central venous-to-arterial CO<sub>2</sub> difference/arterial-central venous O<sub>2</sub> difference ratio is associated with poor lactate clearance in septic patients after resuscitation. *J Crit Care*. 2016; 31(1):76-81.
- [14] Mallat J, Pepy F, Lemyze M, Gasan G, Vangrunderbeeck N, Tronchon L, Vallet B, Thevenin D. Central venous-to-arterial carbon dioxide partial pressure difference in early resuscitation from septic shock: a prospective observational study. *Eur J Anaesthesiol*. 2014;31(7):371-80.
- [15] Zhou J, Song J, Gong S, Li L, Zhang H, Wang M. Persistent hyperlactatemia-high central venous-arterial carbon dioxide to arterial-venous oxygen content ratio is associated with poor outcomes in early resuscitation of septic shock. *Am J Emerg Med*. 2017; 35(8):1136-41.
- [16] Balcı C, Sungurtekin H, Gürses EL, Sungurtekin U. Septik ve nonseptik hastalarda APACHE II, APACHE III, SOFA skorlama sistemleri, trombosit düzeyleri ve mortalite. *Ulus Travma Acil Cerrahi Derg*. 2005;11(1):29-34.
- [17] Thakur R, Rohith VN, Arora JK. Mean SOFA score in comparison with APACHE II score in predicting mortality in surgical patients with sepsis. *Cureus*. 2023;15(3).
- [18] Desai S, Lakhani JD. Utility of SOFA and APACHE II score in sepsis in rural set up MICU. *J Assoc Physicians India*. 2013; 61(9):608-11.