

# The Effect of Paracetamol (Acetaminophen) for Postoperative Pain Control Breast Surgery: Systematic Review and Meta-Analysis Study

Mahfuz Ghaderi, Mohammad Esmail Akbari\*, Atieh Akbari, Sadegh Khoddam, Bashirjamal Wahidi

Cancer Research Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran.

## ARTICLE INFO

### Article history:

Received 19 July 2024  
Revised 10 August 2024  
Accepted 24 August 2024

### Keywords:

Breast surgery;  
Paracetamol;  
Acetaminophen;  
Systematic review;  
Meta-analysis study

## ABSTRACT

**Background:** Breast surgery (BS) is one of the types of surgery that will leave complications for patients. This systematic review study was conducted with the aim of determining the effect of Paracetamol (acetaminophen) on pain after BCS using a systematic review method.

**Methods:** This study aimed to determine the effect of Paracetamol (acetaminophen) on pain after BCS by systematic review method and using PARISMA checklist. The search was conducted in all articles published in Scopus, ISI, PubMed, EBSCO, Embase and Google scholar search engine in the period of 1.1.2000 to 6.1.2024. The checklist included the type of surgery performed, the year the article was published, the type of pain assessment tool, the drug intervention performed, and the results of the drug's effect on the pain of BS patients. Data analysis was done descriptively with the help of Endnote software.

**Results:** According to the findings in the initial search, a number of 545 articles were extracted, and according to the performed screenings, finally, 9 articles were included in the systematic review stage. In relation to pain measurement tool, VAS tool was used in 7 articles. But in the article of De Oliveira et al and colleagues, the QOR-40 scale was used, and in the article of Ohnesorge et al and colleagues, the NRS tool was used. The articles were published between 2005 and 2024. In all the articles, at least two groups were used. The findings of all studies have shown the effect of Paracetamol (acetaminophen) on pain reduction in patients after BS.

**Conclusions:** Considering the effect of Paracetamol (acetaminophen) on reducing the pain of patients with BS, it is suggested that this drug be prescribed to reduce the pain of these patients.

## Introduction

Pain is a mental and complex phenomenon that is defined differently by the patient or the treatment staff [1]. Pain is one of the important complications of diseases that affects many people [2-3]. Considering that pain imposes many physical and psychological complications on patients, it is necessary to identify the factors that are effective in causing pain, and it is

necessary to take necessary measures in this field [4-5]. Also, if the experience of pain relief is insufficient, it affects the patient's social performance and well-being and causes anxiety, depression, and a decrease in the patient's quality of life [6-7].

Reviewing and evaluating sufficient documentation in the field of pain management and regular evaluation is necessary to check the pain status of patients after surgery [8]. Many barriers, including barriers related to employees (nurses, doctors, other personnel), barriers

The authors declare no conflicts of interest.

\*Corresponding author.

E-mail address: profmeakbari@gmail.com

Copyright © 2025 Tehran University of Medical Sciences. Published by Tehran University of Medical Sciences.



This work is licensed under a Creative Commons Attribution-NonCommercial 4.0 International license (<https://creativecommons.org/licenses/by-nc/4.0/>). Noncommercial uses of the work are permitted, provided the original work is properly cited.

related to the system, and barriers related to the environment are effective in pain management [9-10]. Considering the high severity of pain after surgery, for this reason, pain management of patients after surgery should be prioritized [11]. So that it has been estimated that 80% of patients experienced pain after surgery and about 70% of these patients were in severe pain in terms of perceived pain intensity [12-13].

Pain management after surgery is considered one of the most important priorities of the treatment staff. Pain experience varies depending on the type of disease. In fact, pain is influenced by various factors such as the type of surgery, the duration of surgery, the method of surgery, etc. [14-16]. If the pain is not relieved after surgery, complications such as personal suffering, re-admission of the patient to the hospital, cardio-respiratory complications and care pressure will occur in the patient [1].

Breast surgery (BS) is one of the types of surgery that will leave complications for patients [17-19]. One of the reasons for performing BS is the presence of BC in patients [20]. Breast cancer is one of the most common types of cancer among women, and in 2020, with the rate of 2.26 million new cases, it has the highest incidence of cancer among patients. Also, in 2020, more than two-thirds of breast cancer deaths occurred in less developed countries, which is considered a serious health risk in these countries [21-23]. The reported rate of chronic persistent postsurgical pain (CPSP) related to BS has been reported to be high, which is why it is a priority to identify drugs effective in reducing the pain of these patients [24-26].

### **Aim**

This systematic review study was conducted with the aim of determining the effect of Paracetamol (acetaminophen) on pain after BCS using a systematic review method.

### **Methods**

This study aimed to determine the effect of Paracetamol (acetaminophen) on pain after BCS by systematic review method and using PARISMA checklist [1].

### **Selection of studies**

The PICO criterion was used to select the articles.

**P:** Patients with BCS;

**I:** Pain score of patients with BCS;

**C:** Comparison of pain in all of group (Experimental/Control or Placebo);

**O:** The effect of Paracetamol (acetaminophen) drug on patients' pain scores;

### **Paper selection and data extraction**

The search was conducted in all articles published in Scopus, ISI, PubMed, EBSCO, Embase and Google scholar search engine in the period of 1.1.2000 to 6.1.2024.

### **Inclusion and exclusion criteria**

All the original articles in English language that investigated the effect of Paracetamol (acetaminophen) on the pain status of patients with BS were included in the study. If the full file of the article is not accessible for any reason, it is excluded from the study.

### **Data extraction**

The checklist included the type of surgery performed, the year the article was published, the type of pain assessment tool, the drug intervention performed, and the results of the drug's effect on the pain of BS patients.

### **Statistical analysis**

Data analysis was done descriptively with the help of Endnote software.

### **Results**

(Figure 1) shows the findings of the articles included in the systematic review. According to the findings in the initial search, a number of 545 articles were extracted, and according to the performed screenings, finally, 9 articles were included in the systematic review stage.

In relation to pain measurement tool, VAS tool was used in 7 articles. But in the article of De Oliveira et al [27] and colleagues, the QOR-40 scale was used, and in the article of Ohnesorge et al [28] and colleagues, the NRS tool was used. The articles were published between 2005 and 2024. In all the articles, at least two groups were used. The findings of all studies have shown the effect of Paracetamol (acetaminophen) on pain reduction in patients after BS (Table 1).




Figure 1- Flowcharts for systematic review.

Table 1- Characteristics of articles with the prescription of Paracetamol (acetaminophen) to the systematic review stage to reduce pain

-	Author Name	Year	Type of surgery	Scale	Intervention	Result	Conclusion	
1	Mitchell A et al[29]	2008	Outpatient BS	VAS (0–100 mm)	AcIBU: 400 mg of ibuprofen+650 mg of acetaminophen+ T3: acetaminophen + caffeine codeine	AcIBU Group Days 0 = 26.9 (18.1) Days 2 = 19.6(18.2) Days 5 = 14.7(16.4) Days 7 = 10.7(14.6)	T3 Group Days 0 = 30.1(20.6) Days 2 = 20.4(17.4) Days 5 = 14.8(16.5) Days 7 = 12.3(14.9)	According to the findings, the patients' pain was reduced by taking medicine in both groups, but there was no difference between the pain intensity in the AcIBU group and the T3 group. Also, 92% of AcIBU group patients and 89% of T3 group patients were satisfied with pain control.

2	Wagle et al [30]	2018	MRM	VAS	<p><b>Group-I:</b> paracetamol 1g (prior to induction)</p> <p><b>Group-II:</b> paracetamol 1g (end of the surgery)</p>	<p><b>Group-I</b> 0 Mins: 1.9(0.9) 30 Mins: 2(1.4) 1 hrs: 2.5(1.2) 2 hrs: 2.5(0.6) 6 hrs: 1.8(1.5) 12 hrs: 3.6(1.6) 24 hrs: 1.4(0.9)</p>	<p><b>Group-I</b> 0 Mins: 2.9(1.1) 30 Mins: 3.1(1.9) 1 hrs: 3.6(1.9) 2 hrs: 3(0.9) 6 hrs: 2.7(1.2) 12 hrs: 3.9(1.8) 24 hrs: 1.7(1.1)</p>	<p>A significant difference was observed between the two groups' pain from the beginning of the intervention to 6 hours after the intervention (p&lt;0.05), but this difference was not significant at 12 and 24 hours after the intervention (p&gt;0.05). The use of Acetaminophen caused a significant reduction in the pain of patients (P&lt;0.006).</p>
3	De Oliveira et al [27]	2018	Mastectomy	QOR-40 scale	<p><b>Group-I:</b> paracetamol 1g (end of the surgery)</p> <p><b>Group-II:</b> Placebo</p>	<p>Acetaminophen: 31 (29 to 33)</p>	<p>Placebo: 27 (22 to 30)</p>	<p>The pain reduction rate of patients in the group receiving Paracetamol was higher than the patients in the metamizole group. The amount of pain in group 1 and group 4 patients was higher than group 2 and group 3 patients. In all investigated groups, the pain level was reported at the lowest level 6 hours after surgery. The amount of preventive analgesic effect in the group of patients receiving diclofenac was higher than that of patients in the acetaminophen group. Also, if these two drugs were used in combination (group 3), the analgesic effect of this drug was reported as the highest amount of pain reduction.</p>
4	Ohnesorge et al [28]	2005	BCS	NRS (0-10)	<p><b>Group-I:</b> Paracetamol</p> <p><b>Group-II:</b> Metamizole</p> <p><b>Group-III:</b> Placebo</p>	-	-	<p>The pain reduction rate of patients in the group receiving Paracetamol was higher than the patients in the metamizole group. The amount of pain in group 1 and group 4 patients was higher than group 2 and group 3 patients. In all investigated groups, the pain level was reported at the lowest level 6 hours after surgery. The amount of preventive analgesic effect in the group of patients receiving diclofenac was higher than that of patients in the acetaminophen group. Also, if these two drugs were used in combination (group 3), the analgesic effect of this drug was reported as the highest amount of pain reduction.</p>
5	Wadhwa [31]	2015		VAS	<p><b>Group-I:</b> acetaminophen 10mg/kg</p> <p><b>Group-II:</b> diclofenac sodium 1mg/kg</p> <p><b>Group-III:</b> acetaminophen + diclofenac sodium</p> <p><b>Group-III:</b> placebo.</p>	<p><b>Mean VAS</b> <b>G-I:</b> 4.2(0.12) <b>Group-II:</b> 3.47(0.12) <b>Group-III:</b> 2.84(0.12) <b>Group-III:</b> 4.46(0.12)</p>	-	<p>The pain reduction rate of patients in the group receiving Paracetamol was higher than the patients in the metamizole group. The amount of pain in group 1 and group 4 patients was higher than group 2 and group 3 patients. In all investigated groups, the pain level was reported at the lowest level 6 hours after surgery. The amount of preventive analgesic effect in the group of patients receiving diclofenac was higher than that of patients in the acetaminophen group. Also, if these two drugs were used in combination (group 3), the analgesic effect of this drug was reported as the highest amount of pain reduction.</p>

6	Emre et al [32]	2009	Mastectomy	VAS	<b>Group-I:</b> IV paracetamol <b>Group-II:</b> saline	-	-	Paracetamol had reduced the patients' pain, but this reduction was not statistically significant compared to the control group.
7	Kampe et al [33]	2006	BCS	VAS (0–100 mm)	<b>Group-I:</b> IV paracetamol <b>Group-II:</b> saline solution of IV dipyrone	Paracetamol 4 h after: 17.3(18.7) 8 h after: 12(11.6) 20 h after: 11.7(15.4) 30 h after: 7.5(8.7)	Dipyrone 4 h after: 20.9(21.4) 8 h after: 15.3(19.9) 20 h after: 9(15.2) 30 h after: 5.7(9.8)	The use of Paracetamol had reduced the pain of the patients.
8	Nonaka et al [34]	2016	Mastectomy	VAS	<b>Group-I:</b> 1 G acetaminophen <b>Group-II:</b> 50 mg flurbiprofen	-	-	The analgesic effect of flurbiprofen and acetaminophen was equal in patients.
9	Alsaadi et al[35]	2024	BCS	-	<b>Group-I:</b> IV paracetamol <b>Group-II:</b> Placebo	-	-	The use of Paracetamol had reduced the pain of the patients.

## Discussion

This study was conducted with the aim of determining the effect of Paracetamol (acetaminophen) on the pain of patients undergoing BS surgery. According to the findings, the drug Paracetamol (acetaminophen) had reduced the pain of the patients. Paracetamol (acetaminophen) drug is effective in reducing the pain of patients and various studies conducted in this field have confirmed this issue. So, in the study of Coulthard et al., Paracetamol reduced the pain of patients undergoing oral surgery [36] and it is consistent with the results of this study. A person suffering from a disease leads to many complications, including physical and mental complications in the person. Also, a large economic burden is left for the patient and the patient's quality of life undergoes changes [40-43].

In systematic review studies, Paracetamol has also reduced the pain of patients. So that in the study of Mahdavi et al., which had 5 original articles on the effect of Paracetamol on the pain status of Cardiac Surgery patients, this drug had reduced the number of patients [37]. In Ghaffarpasand et al.'s study, which included 5 original articles on the effect of Paracetamol on pain after craniotomy patients, the injection of this drug (before surgery) reduced the pain of patients after surgery [38]. The results of the study by Ghaffarpasand et al. and the study by Mahdavi et al. are consistent with the results of this study.

On the other hand, in the study of Machado et al., the effect of Paracetamol on patients' back or neck pain was investigated in 12 original articles. The findings of a

review study by Machado et al showed that paracetamol did not have a positive effect on the pain, disability and quality of life of patients [39]. From the difference between the results of the current review study in the group of BS patients and the results of the study by Machado et al., it can be pointed out the difference in the type of patients investigated, which may have different effects on the response of the drug to the patients' pain.

## Conclusion

Considering the effect of Paracetamol (acetaminophen) on reducing the pain of patients with BS, it is suggested that this drug be prescribed to reduce the pain of these patients.

## Acknowledgements

Deputy of research and technology in Shahid Beheshti University of medical sciences.

## References

- [1] Mahama F, Ninnoni JP. Assessment and management of postoperative pain among nurses at a resource-constraint teaching Hospital in Ghana. *Nurs Res Pract.* 2019; 2019(1):9091467.
- [2] Vasigh A, Tarjoman A, Borji M. The effect of spiritual-religious interventions on patients' pain status: Systematic review. *Anaesth Pain Intensive Care.* 2018; 22(4).

- [3] Komlakh K, Hatefi M, Soltany B. Comparison of Pain Score in Patients with Brain Disorders Using Care Pain Observation Tool (CPOT) and Nonverbal Pain Scale (NVPS). *Arch Neurosci*. 2022; 9(4).
- [4] Hatefi M, Parvizi R, Borji M, Tarjoman A. Effect of self-management program on pain and disability index in elderly men with osteoarthritis. *Anesth Pain Med*. 2019; 9(4).
- [5] Mohammadi HR, Rahmatian A, Hatefi M, Sadeghi S. Comparison of Pain Intensity, Fear of Movement, and Disability Before and After Lumbar Spine Surgery. *Arch Neurosci*. 2023; 10(2).
- [6] Snijders RA, Brom L, Theunissen M, van den Beuken-van Everdingen MH. Update on prevalence of pain in patients with cancer 2022: a systematic literature review and meta-analysis. *Cancers*. 2023; 15(3):591.
- [7] Hatefi M, Nouri L. Evaluating the Relationship Between Myofascial Pain Syndrome (MPS) and Physical and Mental Health Status in Patients with Stroke. *Arch Neurosci*. 2022; 9(1).
- [8] Temesgen MH, Brihanu A, bekele Teshome Z. Post-operative pain assessment, management compliance with WHO guidelines and its barriers in hospitals of West Shoa zone, central of Ethiopia, 2021. *Ann Med Surg*. 2022; 84.
- [9] Carr E. Barriers to effective pain management. *J Perioper Pract*. 2007;17(5):200-8.
- [10] Mędrzycka-Dąbrowska W, Dąbrowski S, Basiński A. Problems and Barriers in Ensuring Effective Acute and Post-Operative Pain Management--an International Perspective. *Advances in clinical and experimental medicine: official organ Wroclaw Medical University*. 2015; 24(5):905-10.
- [11] Saffari TM, Saffari S, Brower KI, Janis JE. Management of acute surgical pain in *Plast Reconstr Surg*. 2024; 153(4):838e-49e.
- [12] Park R, Mohiuddin M, Arellano R, Pogatzki-Zahn E, Klar G, Gilron I. Prevalence of postoperative pain after hospital discharge: systematic review and meta-analysis. *Pain Reports*. 2023; 8(3):e1075.
- [13] Apfelbaum JL, Chen C, Mehta SS, Gan TJ. Postoperative pain experience: results from a national survey suggest postoperative pain continues to be undermanaged. *Anesth Analg*. 2003; 97(2):534-40.
- [14] Akire SC, Bayraktar N. Outcomes of Pain Management Among Postoperative Patients: A Cross-sectional Study. *J Perianesth Nurs*. 2024; 39(2):240-5.
- [15] Dağcan Şahin N, Gürol Arslan G. Perspectives of patients, families and nurses on pain after cardiac surgery: A qualitative study. *Nurs Crit Care*. 2024; 29(3):501-11.
- [16] de Ladoucette A. Management of perioperative pain after TKA. *Orthopaedics & Traumatology: Surg Res*. 2023; 109(1):103443.
- [17] Alghamdi M, Alsayed B, Albenmoussa F, Salleh SB, Alqaysi L, Alfayez J, et al. Local Anesthetic Infiltration for Pain Control in Aesthetic Breast Reduction Surgery: A Systematic Review and Meta-analysis of Randomized Controlled Trials. *Aesthet Plast Surg*. 2024; 1-8.
- [18] Feili F, Rozegar M. Knowledge of Iranian dentists Regarding of oral cancer: a systematic review and meta-analysis, *International Journal of Cancer Management*, 2024;17(1): <https://brieflands.com/journals/international-journal-of-cancer-management>
- [19] Villa G, Mandarano R, Scirè-Calabrisotto C, Rizzelli V, Del Duca M, Montin DP, et al. Chronic pain after breast surgery: incidence, associated factors, and impact on quality of life, an observational prospective study. *Perioper Med*. 2021; 10:1-1.
- [20] Min J, Kim JY, Ryu J, Park S, Courneya KS, Ligibel J, et al. Early Implementation of Exercise to Facilitate Recovery After Breast Cancer Surgery: A Randomized Clinical Trial. *JAMA surgery*. 2024.
- [21] Xu Y, Gong M, Wang Y, Yang Y, Liu S, Zeng Q. Global trends and forecasts of breast cancer incidence and deaths. *Sci Data*. 2023; 10(1):334.
- [22] Sung H, Ferlay J, Siegel RL, Laversanne M, Soerjomataram I, Jemal A, et al. Global cancer statistics 2020: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *A Cancer J Clin*. 2021; 71(3):209-49.
- [23] Parsae S, Jahani N, Salehiniya H, Raieson M. Effectiveness of Mindfulness Training in Emotion Regulation and Distress Tolerance among Breast Cancer Patients. *Iran J Breast Dis*. 2024; 17(1)
- [24] Yu L, Shen XJ, Liu H, Zhou YT, Zhang Q, Zhang ZD, et al. Effect of ultrasound-guided continuous erector spinae plane block on postoperative pain and inflammatory response in patients undergoing modified radical mastectomy for breast cancer: study protocol for a randomised controlled trial. *Trials*. 2024; 25(1):51.
- [25] Garcia V, Walle J, Leroux-Bromberg N, Delbrouck D, Hannebicque K, Oune FB, et al. Incidence and characteristics of chronic postsurgical pain at 6 months after total mastectomy under pectoroserratus and interpectoral plane block combined with general anesthesia: a prospective cohort study. *Reg Anesth Pain Med*. 2024; 49(1):36-40.
- [26] Tricco AC, Lillie E, Zarin W, O'Brien KK, Colquhoun H, Levac D, et al. PRISMA extension for scoping reviews (PRISMA-ScR): checklist and explanation. *Ann Intern Med*. 2018; 169(7):467-73.
- [27] De Oliveira Jr GS, Rodes ME, Bialek J, Kendall MC, McCarthy RJ. Single dose systemic acetaminophen to improve patient reported quality of recovery after ambulatory segmental mastectomy: a prospective, randomized, double-blinded, placebo controlled, clinical trial. *Breast J*. 2018; 24(3):240-4.
- [28] Ohnesorge H, Bein B, Hanss R, Francksen H, Mayer L, Scholz J, et al. Paracetamol versus metamizol in the treatment of postoperative pain after breast surgery: a randomized, controlled trial. *Eur J Anaesthesiol*. 2009; 26(8):648-53.

- [29] Mitchell A, van Zanten SV, Inglis K, Porter G. A randomized controlled trial comparing acetaminophen plus ibuprofen versus acetaminophen plus codeine plus caffeine after outpatient general surgery. *J Am Coll Surg.* 2008; 206(3):472-9.
- [30] Wagle B, Regmi Y, Shrestha S, Thapa P, Vishwakarma S, Mandal MR, et al. Pre-Emptive Analgesic Effect of Intravenous Paracetamol in Modified radical mastectomy. *Nepal J Cancer.* 2018; 2(1):22-6.
- [31] Wadhwa R. A Randomised Double Blind Study Comparing Preemptive Analgesic Efficacy Of Oral Acetaminophen, Diclofenac And Combination Of Acetaminophen And Diclofenac In Modified Radical Mastectomy Surgery. 2015.
- [32] Emre C, Tuncel G, Kaya M, Canoler Ö, Yıldırım MÖ, Kadioğulları N. The Effect of Intravenous Paracetamol on Morphine Consumption Used for Post-Operative Pain After Modified Radical Mastectomy. *Acta Oncologica Turcica.* 2009; 42(3):100-4.
- [33] Kampe S, Warm M, Landwehr S, Dagtekin O, Haussmann S, Paul M, et al. Clinical equivalence of IV paracetamol compared to IV dipyron for postoperative analgesia after surgery for breast cancer. *Curr Med Res Opin.* 2006; 22(10):1949-54.
- [34] Nonaka T, Hara M, Miyamoto C, Sugita M, Yamamoto T. Comparison of the analgesic effect of intravenous acetaminophen with that of flurbiprofen axetil on post-breast surgery pain: a randomized controlled trial. *J Anesth.* 2016; 30:405-9.
- [35] Alsaadi D, Low L, Ting J, Craughwell M, McDonnell J, Lowery A, et al. Pre-emptive paracetamol reduces intra-operative opioid use in patients undergoing day-case oncologic breast surgery. *EXCLI J.* 2024; 23:356.
- [36] Coulthard P, Bailey E, Patel N. Paracetamol (acetaminophen) for pain after oral surgery. *Oral Surg.* 2014; 7(2):81-6.
- [37] Mahdavi E, Soltani G, Amini S. An Investigation of Safety and Efficacy of Intravenous Paracetamol in Pain Management Following Cardiac Surgery. *J Cardio-Thorac Med.* 2015; 3(3):324-8.
- [38] Ghaffarpassand F, Dadgostar E, Ilami G, Shoaee F, Niakan A, Aghabaklou S, et al. Intravenous acetaminophen (paracetamol) for postcraniotomy pain: systematic review and meta-analysis of randomized controlled trials. *World Neurosurg.* 2020; 134:569-76.
- [39] Machado GC, Maher CG, Ferreira PH, Pinheiro MB, Lin CW, Day RO, et al. Efficacy and safety of paracetamol for spinal pain and osteoarthritis: systematic review and meta-analysis of randomised placebo-controlled trials. *BMJ.* 2015; 350.
- [40] Roozegar MA, Nourmohammadi H, Havasian M, Seidkhani H, Jamdar S. Evaluation of Oral and Dental complications caused by Cyclophosphamide in patients with breast cancer in the Shahid-Mostafa Hospital, Ilam, 2019. *Research Journal of Pharmacy and Technology.* 2022;15(12):5517-20.
- [41] Esmailikia M, Gholami-Parizad E, Ghazanfari Z, Abedzadeh MS, Roozegar MA. Investigation of Oral Health Status (DMFT-index) among 3-6 Years Old Children in Ilam (Western Iran), 2015. *Research J. Pharm. and Tech.* 2020; 13(4): 1876-1880.
- [42] Roozegar MA, Abdi Z, Matin S, Khorshidi A, Azizian M. The effect of 1.2 mg simvastatin mouthwash on the level of interleukin-6 in patients with chronic periodontitis. *Indian Journal of Forensic Medicine & Toxicology.* 2019; 13(3):463-67.
- [43] Roozegar MA, Pakzad I, Mohammadi TM, Hoshmand B. Analyzing the osteogenic stimulatory effect of the combination dexamethasone and low levelled laser irradiation (LLLI) on periodontal ligament stem cell (PDLSc). *Der Pharma Chemica,* 2015, 7(11):226-230.