

Review Article



A Mini Review on the Multifaceted Role of TGF- β in Metastasis Progression: Molecular Mechanisms and Novel Therapeutic Strategies

Yaser Mohammadi ^{1,2}, Mitra Nourbakhsh ^{2*}¹ Student Research Committee, Iran University of Medical Sciences, Tehran, Iran² Department of Biochemistry, School of Medicine, Iran University of Medical Sciences, Tehran, Iran

Article info:

Received: 20 June 2024

Revised: 25 July 2024

Accepted: 29 July 2024

* Corresponding Author:

Mitra Nourbakhsh, Department of Biochemistry, School of Medicine, Iran University of Medical Sciences, Tehran, Iran
Email: Nourbakhsh.m@iums.ac.ir

ABSTRACT

Transforming growth factor-beta (TGF- β) plays a dual role in tumor regulation and metastasis. In the early stages of cancer, TGF- β functions as a tumor suppressor, maintaining cellular homeostasis. However, as cancer progresses, this cytokine can promote tumor invasion and metastasis by modulating the tumor microenvironment and immune responses. TGF- β regulates multiple cellular functions, including growth control, differentiation, apoptosis, and signaling, through SMAD and non-SMAD pathways. In cancer, TGF- β signaling activates the epithelial-mesenchymal transition (EMT), leading to decreased cell adhesion, increased motility, and enhanced tumor invasiveness. These alterations are also linked to chemotherapy resistance. Additionally, TGF- β stimulates angiogenesis by inducing pro-angiogenic factors such as vascular endothelial growth factor (VEGF) and platelet-derived growth factor (PDGF), both of which are essential for metastatic tumor expansion. Furthermore, TGF- β contributes to tumor immune evasion by suppressing T-cell and natural killer (NK) cell activity while promoting regulatory T cells (Tregs). Targeted therapies against TGF- β —including ligand receptor inhibitors, monoclonal antibodies, kinase inhibitors, and antisense oligonucleotides (ASO)—have shown promise in preclinical and clinical studies. However, challenges such as tumor resistance and adverse effects resulting from broad TGF- β inhibition remain. Therefore, ongoing research aims to refine TGF- β -based therapies and integrate them with other treatment modalities, including immunotherapy.

Keywords: Transforming Growth Factor beta (TGF- β); Metastasis; Cancer; Antineoplastic Agents

Use your device to scan and read the article online



Citation: Mohammadi Y. A Mini Review on the Multifaceted Role of TGF- β in Metastasis Progression: Molecular Mechanisms and Novel Therapeutic Strategies. Acta Biochimica Iranica. 2024;2(3):125-131.

doi <https://doi.org/10.18502/abi.v2i3.19096>

Introduction

Cancerous transformations are part of a series of complex changes in cells and tissues that occur in response to internal and external stimuli (1). These modifications enhance the tumor's capacity to penetrate and spread, in addition to increasing

cell growth and proliferation. Numerous cytokines, including transforming growth factor-beta (TGF- β), play a major role in tumor regulation and metastasis from several angles (2).

TGF- β is one of the key cytokines involved in regulating cellular and tissue activities. This cytokine



participates in various processes, including cell growth regulation, differentiation, apoptosis, and cell-to-cell interactions (3). The TGF- β family includes inhibitors, activins, bone morphogenetic proteins (BMPs), and growth differentiation factors (GDFs) (4). Under normal circumstances, TGF- β functions as a tumor suppressor, promoting cellular equilibrium and homeostasis (5).

Nonetheless, TGF- β has a dual function in diseases such as cancer (6). In the early stages, TGF- β suppresses tumors, but as cancer progresses, it can promote invasion and metastasis by influencing the immune system and the tumor microenvironment (7). Furthermore, TGF- β -related signaling pathways may lead to reduced expression of epithelial markers such as E-cadherin and increased expression of mesenchymal markers such as vimentin and N-cadherin (8). Although epithelial-mesenchymal transition (EMT) is essential for normal embryonic development, in cancer progression, it is hijacked to facilitate tumor invasion and metastasis (9).

There are three isoforms of TGF- β : TGF- β 1, TGF- β 2, and TGF- β 3. Cells produce TGF- β from precursor molecules (10). TGF- β homodimers interact with latency-associated peptides (LAP) to form small latent complexes (SLCs). Upon binding to the extracellular matrix, proteolysis activates these latent complexes, enabling TGF- β to bind to its receptors (TGFbRI, TGFbRII, and TGFbRIII) and initiate intracellular signaling (11). This signaling, primarily mediated by suppressor of mothers against decapentaplegic (SMADs), results in changes such as enhanced metastatic spread, alterations in cellular markers, and differentiation, all contributing to cancer formation and progression (12).

Given the intricate role of TGF- β in metastasis, the primary goal of this review is to provide a concise and informative review of the molecular mechanisms of TGF- β in metastasis and its implications for cancer therapy.

Molecular mechanisms of TGF- β in metastasis

SMAD and non-SMAD signaling pathways

SMADs are intracellular signaling proteins and transcription factors within the TGF- β signaling family (13). Activated TGF- β complexes bind to type 2 serine/threonine kinase receptors. TGFbRII then recruits and phosphorylates TGFbRI (14). Zhang et al. demonstrated that in the absence of TGF- β , type 1 and 2 receptors exist as monomers on the cell surface. SMAD2/3 (R-SMADs) are phosphorylated by TGFbRI and subsequently form a heteromeric complex with SMAD4 (coSMAD). This complex functions as a transcription factor for numerous genes after translocating to the nucleus (15, 16), including the activation of the mitogen-activated protein kinase 8 (MAPK8) pathway, which triggers apoptosis (14). SMAD7—an inhibitory SMAD (i-SMAD) that can interact with TGFbRI and prevent SMAD2/3 activation—

E3 ligases (enzymes that ubiquitinate and degrade SMAD2/3 and SMAD4), and SMAD phosphatases (which dephosphorylate SMAD2/3 and SMAD4, leading to their inactivation) are some of the mechanisms that regulate this signaling pathway (11, 17).

Beyond the SMAD-mediated TGF- β signaling route, TGF- β receptors can also activate other intracellular signaling pathways. A significant portion of the complexity and diversity of cellular responses to TGF- β is attributed to these non-SMAD pathways (18). The PI3K-AKT-mTOR pathway is essential for controlling cell division, survival, and metabolism. By interacting with PI3K's p85 subunit, TGF- β can activate the protein, which in turn triggers AKT and subsequently mTOR. mTOR then regulates protein translation (19).

The RhoA/ROCK pathway is crucial for migration, cell motility, and cytoskeletal remodeling. TGF- β can activate ROCK via the RhoA GTPase, leading to cytoskeletal modifications and enhanced cell motility (20). Additionally, three key MAPK pathways—p38, JNK, and Erk—can be activated through phosphorylation and activation of MAPK kinases (MKKs), which are regulated by activated TAK1. The MAPK pathways play a critical role in TGF- β -induced cellular processes: Erk activation enhances gene expression and cell proliferation, while JNK and p38 MAPK activation can lead to cell death (21).

Furthermore, TGF- β can activate other signaling pathways, including NF- κ B, Wnt, and Notch, which regulate gene expression, cell division, and cell death in various ways (18).

Epithelial-mesenchymal transition (EMT)

In healthy cells, EMT is essential for tissue repair and development. On the other hand, TGF- β stimulates EMT in cancer cells at advanced stages of the disease, aiding the growth, metastasis, and advancement of tumor cells (22). Vimentin, E-cadherin, and N-cadherin are among the markers that alter throughout the EMT process, causing a decrease in cell adhesion as well as a loss of cell polarity (23). Furthermore, as mesenchymal traits are acquired, there is an increase in motility, invasiveness, and, eventually, metastasis (22, 23).

Research has indicated that chemotherapy resistance is linked to mesenchymal cancer cells (24). Current research casts doubt on EMT's ability to trigger metastasis. EMT is not necessary for the spread of lung cancer or pancreatic adenocarcinoma, according to studies conducted by Zheng et al. and Fischer et al. These findings suggest that EMT may only be a factor in chemoresistance (25, 26).

TGF- β -induced EMT is mediated by a complex set of signaling pathways and regulatory molecules. TGF- β stimulates the production of SNAIL, ZEB, and TWIST—proteins known to suppress epithelial markers and enhance the expression of mesenchymal genes—via




Figure 1. TGF- β functions through two main signaling pathways: **SMAD pathway:** In this pathway, the TGF- β molecule first binds to its cell surface receptors, namely T β R II, and activates this receptor. Then, T β R II recruits T β R I, and together they phosphorylate receptor-associated SMAD proteins (R-SMAD). This phosphorylation leads to the formation of an R-SMAD complex with co-SMAD proteins. Subsequently, this complex is transported into the cell nucleus, where it regulates the transcription of specific genes, such as the CAGA gene. This pathway directly plays a role in gene regulation and cellular changes. **Non-SMAD pathway:** This pathway is activated when TGF- β binds to other downstream factors. These factors include proteins such as SHC/GRB2/SOS, TRAF4/6, PAR6, and PI3K. Unlike the SMAD pathway, which directly influences gene regulation, the non-SMAD pathway operates through more complex mechanisms involving a variety of proteins and can regulate processes such as cell survival, cell migration, and immune responses.

the SMAD signaling pathway (27). Additionally, non-SMAD signaling pathways, including PI3K/Akt and MAPK pathways (such as ERK, JNK, and p38), are activated by TGF- β . By regulating the expression and activity of transcription factors linked to EMT, as well as downstream effectors, these pathways enhance the EMT process and converge with SMAD pathways (18).

TGF- β also regulates mesenchymal and epithelial markers. Mesenchymal markers, including vimentin, fibronectin, and N-cadherin, are increased, whereas epithelial markers, such as E-cadherin and cytokeratins, are downregulated (28). Notably, TGF- β -induced EMT can be influenced by extracellular matrix components, cytokines, and growth factors (29). The crosstalk between cancer and stromal cells in the tumor microenvironment can further promote the EMT process and contribute to tumor progression and metastasis (28, 29).

The Role of TGF- β in Promoting Metastasis

The intricate and multi-step process of metastasis includes the following steps: a) invasion; b) intravasation, survival, and extravasation from the bloodstream; and c) establishment of new colonies in distant organs (30). Changes in the EMT process of cancer cells and alterations in the tumor microenvironment are two key

variables that enable cancer stem cells (CSCs) to leave the original tumor. As previously discussed, EMT is a complex process. TNF- α , IL-6, and chemokines 4/12 are examples of factors that can further promote EMT within the tumor microenvironment (31). Moreover, TGF- β and TNF- α enhance EMT (30, 31).

Additionally, growth hormones such as FGF and IGF are secreted by cancer cells, driving the tumor microenvironment toward high interstitial fluid pressure (IFP), hypoxia, and acidity (32). These conditions facilitate the remodeling of the extracellular matrix (ECM) surrounding the tumor, allowing cancer-associated fibroblasts (CAFs) to promote metastasis more efficiently (30, 32). TGF- β plays a crucial role in metastasis to various organs, including the lungs, prostate, liver, and bones (33).

The delicate balance of bone remodeling is disrupted by cancer cells that have metastasized to the bone, fostering the formation of bone tumors (34). TGF- β secreted from bone by osteoclasts enhances PTHrP signaling in osteoblasts, leading to reduced OPG expression and increased RANKL expression (34, 35). This imbalance favors RANKL, which increases osteoclastic activity, thereby triggering further production of TGF- β . This creates a positive feedback loop that drives bone metastasis (36).

The Multifaceted Role of TGF- β in Angiogenesis

Angiogenesis is the process by which pre-existing blood vessels divide to generate new ones. It plays a crucial role in both physiological and pathological conditions, such as wound healing and tumor growth (37). TGF- β stimulates angiogenesis through multiple mechanisms. It induces the expression of numerous pro-angiogenic factors, including platelet-derived growth factor (PDGF), fibroblast growth factor (FGF), and vascular endothelial growth factor (VEGF) (38). Additionally, it directly influences endothelial cells, promoting migration and proliferation—two essential phases of angiogenesis (39).

TGF- β regulates the production and accumulation of extracellular matrix (ECM) components, such as glycoproteins, collagen, and fibronectin (40). This complex ECM network provides structural support to newly formed blood vessels (41). Furthermore, TGF- β facilitates the recruitment of smooth muscle cells (SMCs) and pericytes to developing arteries. These supporting cells are essential for maintaining vascular stability and enhancing the maturity and functionality of blood vessels (42).

Additionally, TGF- β modulates the expression of cell adhesion molecules and junctional proteins in endothelial cells, regulating their interactions with neighboring cells. This regulation of cell-cell interactions is critical for coordinating endothelial cell behavior during angiogenesis (43). Overall, TGF- β plays a multifaceted role in angiogenesis by orchestrating various cellular processes involved in blood vessel formation and remodeling.

TGF- β Promotes Immune Evasion

Under normal circumstances, T lymphocytes and natural killer (NK) cells can identify and destroy cancer cells. However, TGF- β helps tumor cells evade the body's immune system. Although the exact mechanism of TGF- β remains unknown, several potential pathways have been proposed, which we will briefly discuss. Gorelik et al. demonstrated that the primary role of TGF- β is to negatively regulate T lymphocytes. TGF- β can impede T and B cell growth, reduce immune factor synthesis by B lymphocytes, affect NK cell function, and hinder Th1 cell development (44).

Regulatory T cells (Tregs) play a crucial role in tumor immune evasion. TGF- β stimulates FoxP3 expression and induces Tregs that suppress immune responses against tumors (45). Dendritic cells (DCs) prime and activate T cells during the immune response. TGF- β promotes apoptosis and prevents DC migration, facilitating immune evasion (46). Tumor-associated macrophages (TAMs) are polarized by TGF- β from an M1 anti-tumor phenotype to an M2 pro-tumor phenotype (47). Additionally, TGF- β can inhibit neutrophil

degranulation and promote the synthesis of interleukin 10 (IL-10).

Overall, TGF- β plays a significant role in tumor immune evasion by regulating multiple immune cells and promoting immunosuppressive factors (48).

Targeted TGF- β Therapy Strategies, Challenges, and Future Prospects

Therapeutic Strategies

TGF- β is a molecule with dual functions in cancer development. It may act as a tumor growth inhibitor or, in some cases, facilitate disease progression. Thus, developing effective cancer treatments requires a thorough understanding of the TGF- β signaling system. Fortunately, scientific research in this field has advanced significantly. Several investigations have demonstrated that pharmacological therapies targeting different components of the TGF- β signaling system have produced encouraging outcomes in preclinical and clinical trials. TGF- β -based treatment strategies can generally be categorized into five groups: 1) Ligand traps; 2) Monoclonal antibodies; 3) Receptor kinase inhibitors; 4) Antisense oligonucleotides; and 5) Aptamers (peptide- and nucleotide-based) (49).

A comprehensive assessment of several factors must be conducted to optimize therapeutic applications. Key considerations include evaluating TGF- β levels and isoforms, the expression of genes responsive to TGF- β , the types of cells influenced by TGF- β , genetic analysis of tumors for expression and mutations in T β RII, T β RI, and SMADs, and the extent of host cellular responses (49, 50).

The monoclonal antibody 1D11, which binds to all three isoforms of TGF- β and inhibits its activity to prevent bone tumor development, was developed for use in mice (51). For potential patient applications, Genzyme Co. has developed human versions of these neutralizing antibodies, including Lerdelimumab, Fresolimumab, and Metelimumab (52). Numerous T β RI kinase inhibitors have been designed and are currently in clinical trials for cancer treatment. The T β RI kinase inhibitor Galunisertib (LY2157299) has demonstrated satisfactory safety and tolerability in early-phase trials involving Japanese patients with advanced solid malignancies (53). Furthermore, it is currently being investigated in combination with immunotherapy and anti-PD-1 antibodies such as nivolumab and durvalumab (Clinical Trials: NCT02423343) (54). Studies have shown that T β RI inhibitors, such as SM16, and TGF- β neutralizing antibodies, such as 1D11, synergistically prevent metastasis when combined with OX40 agonist antibodies (55). In melanoma and glioma models, the T β RI inhibitor SD208 has demonstrated efficacy in halting bone metastases and slowing the progression of osteolytic lesions by blocking the TGF- β /SMAD pathway (56).

Antisense oligonucleotide (ASO) therapies represent another cutting-edge treatment strategy that can inhibit TGF- β -induced metastasis by reducing its production. ASOs bind to corresponding RNA sequences, leading to mRNA degradation and decreased TGF- β protein synthesis (57). For instance, Trabedersen (AP 12009) is an ASO specific to human TGF- β 2 coding RNA that inhibits the production of this isoform, thereby preventing TGF- β -driven metastasis (58). Another ASO, Belagenpumatucel-L (Lucanix), has demonstrated anti-tumor efficacy in non-small cell lung cancer (NSCLC) by inhibiting TGF- β 2 to reduce its immunosuppressive effects (59).

Due to their specificity in target binding, small molecule peptides offer another therapeutic approach that can suppress the TGF- β signaling pathway while minimizing adverse effects compared to other treatment modalities (60). Han Kang K et al. demonstrated that the TAT-SNX9 peptide exhibits anti-fibrotic properties against TGF- β in mouse and human lung fibroblast cells (61). SNX9 inhibits the TGF- β signaling pathway by binding to SMAD3 and preventing its nuclear translocation (57). Additionally, Trx-SARA inhibits the formation of the SMAD2-3/SMAD4 complex. Studies have also shown that Trx-SARA can suppress the TGF- β -induced epithelial-mesenchymal transition (EMT) process (62).

Advantages and Disadvantages

Each of these treatment approaches has unique benefits and drawbacks. While receptor and ligand inhibitors broadly block TGF- β signaling throughout the body, they also pose a risk of adverse effects. Monoclonal antibodies, on the other hand, offer a more precise means of targeting TGF- β in specific cells or tissues, potentially reducing side effects. However, the high cost and complexity of monoclonal antibody production present additional challenges.

Challenges

The development of effective TGF- β therapies faces several hurdles. Key challenges include:

- **Improving the specificity of TGF- β inhibitors** – Current inhibitors block all three isoforms of TGF- β , potentially leading to side effects. Developing isoform-specific inhibitors is essential.
- **Overcoming resistance to TGF- β therapy** – Some tumors exhibit resistance to TGF- β -targeted treatments. Understanding these resistance mechanisms and formulating strategies to counteract them is crucial.
- **Developing combination therapies** – TGF- β therapy may be more effective when paired with other treatments such as immunotherapy or chemotherapy. Identifying optimal therapeutic combinations is a priority.

Future Prospects

Researchers continue to explore novel approaches to improve TGF- β therapies. Some of the most promising areas of investigation include:

- **Nano-based therapies** – Nanotechnology can enhance the targeted delivery of TGF- β inhibitors to tumors or specific cells, thereby minimizing side effects and improving efficacy.
- **Gene therapy** – Gene therapy offers a potential long-term solution by introducing genes into cells that express TGF- β inhibitors.
- **Personalized medicine** – Scientists are working to identify patients who are most likely to benefit from TGF- β therapy, ensuring that treatments are tailored to individual needs.

TGF- β remains a promising target for cancer treatment. However, further research is necessary to develop more effective therapies. Some of these treatments are currently undergoing clinical trials, with the hope that they will receive approval for broader use in the coming years.

Acknowledgements

The authors would like to express their gratitude to Mrs. Asma Rafiei for her valuable assistance in the grammatical editing of this manuscript.

Conflict of Interest

The authors declare that they have no conflicts of interest.

References

1. Lang BJ, Guerrero-Giménez ME, Prince TL, Ackerman A, Bonorino C, Calderwood SK. Heat Shock Proteins Are Essential Components in Transformation and Tumor Progression: Cancer Cell Intrinsic Pathways and Beyond. *Int J Mol Sci*. 2019 Sep 11;20(18):4507. <https://doi.org/10.3390/ijms20184507>
2. Gholinejad Z, Kheiripour N, Nourbakhsh M, Ilbeigi D, Behroozfar K, Hesari Z, et al. Extracellular NAMPT/Visfatin induces proliferation through ERK1/2 and AKT and inhibits apoptosis in breast cancer cells. *Peptides*. 2017 Jun;92:9-15. <https://doi.org/10.1016/j.peptides.2017.04.007>
3. Wang A, Huang Y, Yang X. TGF- β promotes proliferation and inhibits apoptosis of liver cancer Huh-7 cells by regulating MiR-182/CADM1. *Trop J Pharm Res*. 2023;22(9):1785-95. <https://doi.org/10.4314/tjpr.v22i9.4>
4. Golán-Cancela I, Caja L. The TGF- β Family in Glioblastoma. *Int J Mol Sci*. 2024 Jan 15;25(2):1067. <https://doi.org/10.3390/ijms25021067>
5. David CJ, Huang YH, Chen M, Su J, Zou Y, Bardeesy N, et al. TGF- β Tumor Suppression through a Lethal EMT.

- Cell. 2016 Feb 25;164(5):1015-30. <https://doi.org/10.1016/j.cell.2016.01.009>
6. Zhang M, Zhang YY, Chen Y, Wang J, Wang Q, Lu H. TGF- β Signaling and Resistance to Cancer Therapy. *Front Cell Dev Biol.* 2021 Nov 30;9:786728. <https://doi.org/10.3389/fcell.2021.786728>
 7. Li L, Yu R, Cai T, Chen Z, Lan M, Zou T, et al. Effects of immune cells and cytokines on inflammation and immunosuppression in the tumor microenvironment. *Int Immunopharmacol.* 2020 Nov;88:106939. <https://doi.org/10.1016/j.intimp.2020.106939>
 8. Ji Q, Liu X, Han Z, Zhou L, Sui H, Yan L, et al. Resveratrol suppresses epithelial-to-mesenchymal transition in colorectal cancer through TGF- β /Smads signaling pathway mediated Snail/E-cadherin expression. *BMC Cancer.* 2015 Mar 5;15:97. <https://doi.org/10.1186/s12885-015-1119-y>
 9. Zhu Y, Wu J, Ma W, Zhang H, Wang D. Expression of TGF- β 1, Snail, E-cadherin and N-cadherin in gastric cancer and its significance. *Chin J Clin Oncol* 2007;4:384-9. <https://doi.org/10.1007/s11805-007-0384-1>
 10. Minto AW, Wilson HM, Rees AJ, Quigg RJ, Brown PA. Selective expression of TGF- β 2 and TGF- β 3 isoforms in early mesangioproliferative glomerulonephritis. *Nephron Exp Nephrol.* 2004;96(4):e111-8. <https://doi.org/10.1159/000077377>
 11. Li Y, Fan W, Link F, Wang S, Dooley S. Transforming growth factor β latency: A mechanism of cytokine storage and signalling regulation in liver homeostasis and disease. *JHEP Rep.* 2021 Nov 18;4(2):100397. <https://doi.org/10.1016/j.jhepr.2021.100397>
 12. Wu M-Z, Yuan Y-c, Huang B-Y, Chen J-X, Li B-K, Fang J-H, et al. Identification of a TGF- β /SMAD/Inc-UTGF positive feedback loop and its role in hepatoma metastasis. *Signal Transduct Target Ther.* 2021 Nov 17;6(1):395. <https://doi.org/10.1038/s41392-021-00781-3>
 13. Afshari H, Noori S, Nourbakhsh M, Daraei A, Azami Movahed M, Zarghi A. A novel imidazo[1,2-a]pyridine derivative and its co-administration with curcumin exert anti-inflammatory effects by modulating the STAT3/NF- κ B/iNOS/COX-2 signaling pathway in breast and ovarian cancer cell lines. *Bioimpacts.* 2024;14(2):27618. <https://doi.org/10.34172/bi.2023.27618>
 14. Wrana JL. Signaling by the TGF β superfamily. *Cold Spring Harb Perspect Biol.* 2013 Oct 1;5(10):a011197. <https://doi.org/10.1101/cshperspect.a011197>
 15. Budi EH, Duan D, Derynck R. Transforming Growth Factor- β Receptors and Smads: Regulatory Complexity and Functional Versatility. *Trends Cell Biol.* 2017 Sep;27(9):658-72. <https://doi.org/10.1016/j.tcb.2017.04.005>
 16. Nakao A, Imamura T, Souchelnytskyi S, Kawabata M, Ishisaki A, Oeda E, et al. TGF-beta receptor-mediated signalling through Smad2, Smad3 and Smad4. *EMBO J.* 1997 Sep 1;16(17):5353-62. <https://doi.org/10.1093/emboj/16.17.5353>
 17. Li J, Zou L, Zhou Y, Li L, Zhu Y, Yang Y, et al. A low-frequency variant in SMAD7 modulates TGF- β signaling and confers risk for colorectal cancer in Chinese population. *Mol Carcinog.* 2017 Jul;56(7):1798-807. <https://doi.org/10.1002/mc.22637>
 18. Aashaq S, Batool A, Mir SA, Beigh MA, Andrabi KI, Shah ZA. TGF- β signaling: A recap of SMAD-independent and SMAD-dependent pathways. *J Cell Physiol.* 2022 Jan;237(1):59-85. <https://doi.org/10.1002/jcp.30529>
 19. Baek SH, Ko JH, Lee JH, Kim C, Lee H, Nam D, et al. Ginkgolic Acid Inhibits Invasion and Migration and TGF- β -Induced EMT of Lung Cancer Cells Through PI3K/Akt/mTOR Inactivation. *J Cell Physiol.* 2017 Feb;232(2):346-54. <https://doi.org/10.1002/jcp.25426>
 20. Manickam N, Patel M, Griendling KK, Gorin Y, Barnes JL. RhoA/Rho kinase mediates TGF- β 1-induced kidney myofibroblast activation through Poldip2/Nox4-derived reactive oxygen species. *Am J Physiol Renal Physiol.* 2014 Jul 15;307(2):F159-71. <https://doi.org/10.1152/ajprenal.00546.2013>
 21. Li J, Zhao Z, Liu J, Huang N, Long D, Wang J, et al. MEK/ERK and p38 MAPK regulate chondrogenesis of rat bone marrow mesenchymal stem cells through delicate interaction with TGF-beta1/Smads pathway. *Cell Prolif.* 2010 Aug;43(4):333-43. <https://doi.org/10.1111/j.1365-2184.2010.00682.x>
 22. Papageorgis P. TGF β Signaling in Tumor Initiation, Epithelial-to-Mesenchymal Transition, and Metastasis. *J Oncol.* 2015;2015:587193. <https://doi.org/10.1155/2015/587193>
 23. Loh C-Y, Chai JY, Tang TF, Wong WF, Sethi G, Shanmugam MK, et al. The E-Cadherin and N-Cadherin Switch in Epithelial-to-Mesenchymal Transition: Signaling, Therapeutic Implications, and Challenges. *Cells.* 2019 Sep 20;8(10):1118. <https://doi.org/10.3390/cells8101118>
 24. Roodhart JM, Daenen LG, Stigter EC, Prins HJ, Gerrits J, Houthuijzen JM, et al. Mesenchymal stem cells induce resistance to chemotherapy through the release of platinum-induced fatty acids. *Cancer Cell.* 2011 Sep 13;20(3):370-83. <https://doi.org/10.1016/j.ccr.2011.08.010>
 25. Fischer KR, Durrans A, Lee S, Sheng J, Li F, Wong ST, et al. Epithelial-to-mesenchymal transition is not required for lung metastasis but contributes to chemoresistance. *Nature.* 2015;527(7579):472-6. <https://doi.org/10.1038/nature15748>
 26. Zheng X, Carstens JL, Kim J, Scheible M, Kaye J, Sugimoto H, et al. Epithelial-to-mesenchymal transition is dispensable for metastasis but induces chemoresistance in pancreatic cancer. *Nature.* 2015 Nov 26;527(7579):525-30. <https://doi.org/10.1038/nature16064>
 27. Lamouille S, Xu J, Derynck R. Molecular mechanisms of epithelial-mesenchymal transition. *Nat Rev Mol Cell Biol.* 2014 Mar;15(3):178-96. <https://doi.org/10.1038/nrm3758>
 28. Shirakihara T, Saitoh M, Miyazono K. Differential regulation of epithelial and mesenchymal markers by deltaEF1 proteins in epithelial mesenchymal transition induced by TGF-beta. *Mol Biol Cell.* 2007 Sep;18(9):3533-44. <https://doi.org/10.1091/mbc.e07-03-0249>
 29. Boye A. A cytokine in turmoil: Transforming growth factor beta in cancer. *Biomed Pharmacother.* 2021 Jul;139:111657. <https://doi.org/10.1016/j.biopha.2021.111657>
 30. Majidpoor J, Mortezaee K. Steps in metastasis: an updated review. *Med Oncol.* 2021 Jan 4;38(1):3. <https://doi.org/10.1007/s12032-020-01447-w>
 31. Chen Q, Yang D, Zong H, Zhu L, Wang L, Wang X, et al. Growth-induced stress enhances epithelial-mesenchymal transition induced by IL-6 in clear cell renal cell carcinoma via the Akt/GSK-3 β / β -catenin signaling pathway. *Oncogenesis.* 2017 Aug 28;6(8):e375. <https://doi.org/10.1038/oncsis.2017.74>
 32. Jung HY, Fattet L, Yang J. Molecular pathways: linking tumor microenvironment to epithelial-mesenchymal transition in metastasis. *Clin Cancer Res.* 2015 Mar 1;21(5):962-8. <https://doi.org/10.1158/1078-0432.ccr-13-3173>
 33. Nguyen DX, Bos PD, Massagué J. Metastasis: from dissemination to organ-specific colonization. *Nat Rev Cancer.* 2009 Apr;9(4):274-84. <https://doi.org/10.1038/nrc2622>
 34. Yin JJ, Selander K, Chirgwin JM, Dallas M, Grubbs BG, Wieser R, et al. TGF-beta signaling blockade inhibits PTHrP secretion by breast cancer cells and bone metastases development. *J Clin Invest.* 1999 Jan;103(2):197-206. <https://doi.org/10.1172/jci3523>
 35. Kang Y, Siegel PM, Shu W, Drobnjak M, Kakonen SM, Cordon-Cardo C, et al. A multigenic program mediating breast cancer metastasis to bone. *Cancer Cell.* 2003 Jun;3(6):537-49. [https://doi.org/10.1016/s1535-6108\(03\)00132-6](https://doi.org/10.1016/s1535-6108(03)00132-6)

36. Li B, Wang P, Jiao J, Wei H, Xu W, Zhou P. Roles of the RANKL-RANK Axis in Immunity-Implications for Pathogenesis and Treatment of Bone Metastasis. *Front Immunol*. 2022 Mar 21;13:824117. <https://doi.org/10.3389/fimmu.2022.824117>
37. Guo J, Hu Z, Yan F, Lei S, Li T, Li X, et al. Angelica dahurica promoted angiogenesis and accelerated wound healing in db/db mice via the HIF-1 α /PDGF- β signaling pathway. *Free Radic Biol Med*. 2020 Nov 20;160:447-57. <https://doi.org/10.1016/j.freeradbiomed.2020.08.015>
38. Jiang X, Wang J, Deng X, Xiong F, Zhang S, Gong Z, et al. The role of microenvironment in tumor angiogenesis. *J Exp Clin Cancer Res*. 2020 Sep 30;39(1):204. <https://doi.org/10.1186/s13046-020-01709-5>
39. Zong J, Jiang J, Shi P, Liu J, Wang W, Li B, et al. Fatty acid extracts facilitate cutaneous wound healing through activating AKT, ERK, and TGF- β /Smad3 signaling and promoting angiogenesis. *Am J Transl Res*. 2020 Feb 15;12(2):478-92.
40. Tsuda T. Extracellular Interactions between Fibulins and Transforming Growth Factor (TGF)- β in Physiological and Pathological Conditions. *Int J Mol Sci*. 2018 Sep 17;19(9):2787. <https://doi.org/10.3390/ijms19092787>
41. Davis GE, Senger DR. Endothelial extracellular matrix: biosynthesis, remodeling, and functions during vascular morphogenesis and neovessel stabilization. *Circ Res*. 2005 Nov 25;97(11):1093-107. <https://doi.org/10.1161/01.res.0000191547.64391.e3>
42. Ricard N, Tu L, Le Hirsch M, Huertas A, Phan C, Thuillet R, et al. Increased pericyte coverage mediated by endothelial-derived fibroblast growth factor-2 and interleukin-6 is a source of smooth muscle-like cells in pulmonary hypertension. *Circulation*. 2014 Apr 15;129(15):1586-97. <https://doi.org/10.1161/circulationaha.113.007469>
43. R Rossi E, Bernabeu C, Smadja DM. Endoglin as an Adhesion Molecule in Mature and Progenitor Endothelial Cells: A Function Beyond TGF- β . *Front Med (Lausanne)*. 2019 Jan 30;6:10. <https://doi.org/10.3389/fmed.2019.00010>
44. Gorelik L, Flavell RA. Abrogation of TGF β signaling in T cells leads to spontaneous T cell differentiation and autoimmune disease. *Immunity*. 2000 Feb;12(2):171-81. [https://doi.org/10.1016/s1074-7613\(00\)80170-3](https://doi.org/10.1016/s1074-7613(00)80170-3)
45. Ma X, Gao Y, Chen Y, Liu J, Yang C, Bao C, et al. M2-Type Macrophages Induce Tregs Generation by Activating the TGF- β /Smad Signalling Pathway to Promote Colorectal Cancer Development. *Onco Targets Ther*. 2021 Dec 7;14:5391-402. <https://doi.org/10.2147/ott.s336548>
46. Fucikova J, Palova-Jelinkova L, Bartunkova J, Spisek R. Induction of Tolerance and Immunity by Dendritic Cells: Mechanisms and Clinical Applications. *Front Immunol*. 2019 Oct 29;10:2393. <https://doi.org/10.3389/fimmu.2019.02393>
47. Ge Z, Ding S. The Crosstalk Between Tumor-Associated Macrophages (TAMs) and Tumor Cells and the Corresponding Targeted Therapy. *Front Oncol*. 2020 Nov 3;10:590941. <https://doi.org/10.3389/fonc.2020.590941>
48. Li MO, Sanjabi S, Flavell RA. Transforming growth factor-beta controls development, homeostasis, and tolerance of T cells by regulatory T cell-dependent and -independent mechanisms. *Immunity*. 2006 Sep;25(3):455-71. <https://doi.org/10.1016/j.immuni.2006.07.011>
49. Danielpour D. Advances and Challenges in Targeting TGF- β Isoforms for Therapeutic Intervention of Cancer: A Mechanism-Based Perspective. *Pharmaceuticals (Basel)*. 2024 Apr 20;17(4):533. <https://doi.org/10.3390/ph17040533>
50. Teixeira AF, Ten Dijke P, Zhu HJ. On-Target Anti-TGF- β Therapies Are Not Succeeding in Clinical Cancer Treatments: What Are Remaining Challenges? *Front Cell Dev Biol*. 2020 Jul 8;8:605. <https://doi.org/10.3389/fcell.2020.00605>
51. Edwards JR, Nyman JS, Lwin ST, Moore MM, Esparza J, O'Quinn EC, et al. Inhibition of TGF- β signaling by 1D11 antibody treatment increases bone mass and quality in vivo. *J Bone Miner Res*. 2010 Nov;25(11):2419-26. <https://doi.org/10.1002/jbmr.139>
52. Cordeiro MF, Gay JA, Khaw PT. Human anti-transforming growth factor-beta2 antibody: a new glaucoma anti-scarring agent. *Invest Ophthalmol Vis Sci*. 1999 Sep;40(10):2225-34.
53. Fujiwara Y, Nokihara H, Yamada Y, Yamamoto N, Sunami K, Utsumi H, et al. Phase 1 study of galunisertib, a TGF-beta receptor I kinase inhibitor, in Japanese patients with advanced solid tumors. *Cancer Chemother Pharmacol*. 2015 Dec;76(6):1143-52. <https://doi.org/10.1007/s00280-015-2895-4>
54. Nadal E, Saleh M, Aix SP, Ochoa-de-Olza M, Patel SP, Antonia S, et al. A phase Ib/II study of galunisertib in combination with nivolumab in solid tumors and non-small cell lung cancer. *BMC Cancer*. 2023 Jul 28;23(1):708. <https://doi.org/10.1186/s12885-023-11153-1>
55. Garrison K, Hahn T, Lee WC, Ling LE, Weinberg AD, Akporiaye ET. The small molecule TGF- β signaling inhibitor SM16 synergizes with agonistic OX40 antibody to suppress established mammary tumors and reduce spontaneous metastasis. *Cancer Immunol Immunother*. 2012 Apr;61(4):511-21. <https://doi.org/10.1007/s00262-011-1119-y>
56. Mohammad KS, Javelaud D, Fournier PG, Niewolna M, McKenna CR, Peng XH, et al. TGF-beta-RI kinase inhibitor SD-208 reduces the development and progression of melanoma bone metastases. *Cancer Res*. 2011 Jan 1;71(1):175-84. <https://doi.org/10.1158/0008-5472.can-10-2651>
57. Xie F, Ling L, van Dam H, Zhou F, Zhang L. TGF- β signaling in cancer metastasis. *Acta Biochim Biophys Sin (Shanghai)*. 2018 Jan 1;50(1):121-32. <https://doi.org/10.1093/abbs/gmx123>
58. Schlingensiepen KH, Jaschinski F, Lang SA, Moser C, Geissler EK, Schlitt HJ, et al. Transforming growth factor-beta 2 gene silencing with trabedersen (AP 12009) in pancreatic cancer. *Cancer science*. 2011;102(6):1193-200. <https://doi.org/10.1111/j.1349-7006.2011.01917.x>
59. Zappa C, Mousa SA. Non-small cell lung cancer: current treatment and future advances. *Transl Lung Cancer Res*. 2016 Jun;5(3):288-300. <https://doi.org/10.21037/tlcr.2016.06.07>
60. Ramsey JD, Flynn NH. Cell-penetrating peptides transport therapeutics into cells. *Pharmacol Ther*. 2015 Oct;154:78-86. <https://doi.org/10.1016/j.pharmthera.2015.07.003>
61. Kang JH, Jung MY, Yin X, Andrianifahanana M, Hernandez DM, Leof EB. Cell-penetrating peptides selectively targeting SMAD3 inhibit profibrotic TGF- β signaling. *J Clin Invest*. 2017 Jun 30;127(7):2541-54. <https://doi.org/10.1172/jci88696>
62. Zhao BM, Hoffmann FM. Inhibition of transforming growth factor-beta1-induced signaling and epithelial-to-mesenchymal transition by the Smad-binding peptide aptamer Trx-SARA. *Mol Biol Cell*. 2006 Sep;17(9):3819-31. <https://doi.org/10.1091/mbc.e05-10-0990>