



Activity of *Citrus aurantium* and *Lavandula angustifolia* in Alzheimer's Disease Symptoms in Male Wistar Rats

Amir Arasteh^{1*}, Morteza Karimpour¹, Faezeh Fallah², Sara Kiani² and Maedeh Kakavan³

1. Department of Biology, Rasht branch, Islamic Azad University, Rasht, Iran

2. Department of Biology, Islamic Azad University, Central Tehran branch, Tehran, Iran

3. Department of Medical Microbiology, Faculty of Medicine, Mazandaran University of Medical Sciences, Sari, Iran

Abstract

Background: Alzheimer's Disease (AD) is one of the most prevalent chronic neurodegenerative disorders. The present study aims to better understand the mechanism by which *Citrus aurantium* (*C. aurantium*) and *Lavandula angustifolia* (*L. angustifolia*) hydro-alcoholic extracts were used to treat AD and anti-oxidant issues in a laboratory model.

Methods: 15 male Wistar rats, weighing 250 ± 20 gr, aged 6–8 weeks, were used. Amyloids in the brain were found and identified using the shuttle box and Congo red test. ELISA testing for norepinephrine and serotonin, Superoxide Dismutase (SOD), Malondialdehyde (MDA), and Real-time PCR for expression of the *APP* and *GLT1* genes were done.

Results: The shuttle box test demonstrated that AD produces behavioral harm, since it significantly reduces passive avoidance learning. The Congo red test revealed that the AD models had much more amyloid beta in their brain tissue than the control. Norepinephrine levels were also decreased by using both extracts in test group. Treatment with both extracts led to a substantial rise in SOD activity and fall in MDA concentration.

Conclusion: The gene expression data indicated that the relative expression of the *APP* and *GLT1* genes was shown to be lower in the groups treated with both extracts. *C. aurantium* and *L. angustifolia* may therefore offer a multi-target treatment strategy for AD, which calls for more research in this area.

Keywords: Amyloid beta-peptides, Antioxidants, Brain, Citrus, Lavandula, Neurodegenerative diseases, Norepinephrine, Rats, Serotonin

To cite this article: Arasteh A, Karimpour M, Fallah F, Kiani S, Kakavan M. Activity of *Citrus aurantium* and *Lavandula angustifolia* in Alzheimer's Disease Symptoms in Male Wistar Rats. Avicenna J Med Biotech 2023;15(4):223-231.

* Corresponding author:
Amir Arasteh, Ph.D.,
Department of Biology, Rasht
Branch, Islamic Azad University,
Rasht, Iran
Tel: +98 13 33423308
E-mail:
arasteh@iaurasht.ac.ir
Received: 12 Mar 2023
Accepted: 22 Jul 2023

Introduction

One of the most prevalent chronic neurodegenerative disorders, Alzheimer's Disease (AD), is characterized by neural plaques and neurofibrillary tangles as pathological signs. Both the hyper-phosphorylation of tau protein linked with microtubules in neurons and the accumulation of amyloid beta peptide ($A\beta$) in brain tissue are responsible for these symptoms. The amyloidogenic processing is influenced by a variety of genetic and environmental variables¹. $A\beta$ (abeta) accumulation in the brain starts a pathogenic cascade that eventually results in synaptic dysfunction and loss, neuronal death, and finally cognitive failure. Inflammation and oxidative stress are also a part of this process, which contribute to neurological dysfunction²⁻⁴. The

only available treatments for AD are one partial N-Methyl-D-aspartate (NMDA) receptor antagonist (memantine) and four cholinesterase inhibitors (tacrine, rivastigmine, donepezil, and galantamine).

Herbal and medicinal extracts are also being studied in addition to therapeutic substances. These studies, which include the incidence and prevalence of AD in many cultures and nations, are based in part on nutritional and epidemiological investigations. For instance, a considerable reduced incidence of AD (4.4-fold) was seen among older Indians compared to an American reference sample in a major epidemiologic investigation⁵.

In the primary healthcare system, medicinal plants

are seen as being quite significant. Numerous herbal therapies have been examined in clinical studies and other AD-related models, and they have all demonstrated positive results ⁶.

Citrus aurantium var. *amara* L. (*C. aurantium*) (www.theplantlist.org, record 28100679), the plant known by its scientific name, is native to Asia's tropical regions; however, it may also be found in other tropical and subtropical areas. In Asia, *C. aurantium* is used as a brew or tea and *C. aurantium* sweat. The sweat of *C. aurantium* is used in salad dressing, as an air freshener and in the bedroom for better sleep. In Iran, to reduce the complications of AD in the elderly, plant syrup containing one third of a glass of *C. aurantium* sweat, one third of water and one third of rose water is used. Blossom of the orange tree, is widely used in perfumery and making essential oils, all kinds of seasonal drinks, and preparing jam. This herb is widely utilized in the treatment of several ailments and is freely accessible ⁷. Flavonoids, including hesperidin, naringin, and alkaloids, primarily synephrine, are the primary physiologically active components of *C. aurantium* and have positive medicinal benefits on human health. These substances have the reputation of a novel drug due to a variety of therapeutic benefits including disinfectant, anticancer, anti-inflammatory, and anti-oxidant, as well as its involvement in the treatment of cardiovascular, mental, and neurological problems ⁸.

Another plant whose anti-Alzheimer effects were investigated in this research is *Lavandula angustifolia* (*L. angustifolia*) Mill (www.theplantlist.org, record 108971). This plant (English name: Lavender), is widely used in Iran and Mediterranean regions to reduce Alzheimer's complications. In Iran, *L. angustifolia* brew is used to prevent and reduce the side effects of AD. It has been demonstrated in multiple researches ⁹ to have a variety of therapeutic benefits, including sedative, anticonvulsant, analgesic, and local anesthetic properties. Additionally, the nervous system is soothed and relaxed by this herb ¹⁰. In Alzheimer's patients, the amount of norepinephrine and Malondialdehyde (MDA) (an indicator of oxidative stress) increases, but the concentration of serotonin and the activity of the Superoxide Dismutase (SOD) enzyme decrease. Also, the expression of the Glutamate transporter-1 (*GLT1*) gene increases, and promising results were obtained in all cases with the use of medicinal plants. The present research aims to better understand the mechanism of

action of *C. aurantium* and *L. angustifolia* hydro-alcoholic extracts in treating anti-oxidant issues typical of AD in an induced laboratory rat model. This understanding may protect the biodiversity rights of indigenous people and open the door to potential therapeutic applications.

Materials and Methods

Preparation of herbal extracts

Identified and documented genus and species of *C. aurantium* and *L. angustifolia* plants were purchased from Medicinal Plants Research Center, Tehran University of Medical Sciences. The extraction was performed by the maceration method (ethanol 96%) for both plants. For example, 10 g of the dried plant powder was poured into 200 ml of 96% ethanol solution. The solution was stirred for 24 hr and filtered through a filter paper to obtain the extract.

GC-MS analysis

GC/MS analyses were performed by 7890B series gas chromatography device (Agilent Company of America), Column: 30 mm×0.25 mm and 0.25 mm film thickness. For *L. angustifolia*, the oven temperature program started from 60–220°C for 20 min at a rate of 6°C/min. Transfer line temperature was 280°C; injector temperature was 250°C. Components were detected by comparing the resulting spectrum to the mass pattern found in the analyzer library ¹¹. For *C. aurantium*, an aliquot of 10 µl was placed in GC-MS syringe and oven temperature of 60°C for 10 min. The temperature was increased to a final temperature of 250°C in rate of 5°C/min. Injector and detector temperatures were 230°C and 300°C, respectively ¹².

Animal studies, model development and treatment

15 male Wistar rats, weighing 250±20 g, were acquired from the Pasteur Institute of Iran for this investigation. They were housed at a temperature of 23±3°C and a humidity of 50±10 throughout the experiment. Up to the completion of the treatment, the rats had unrestricted access to animal food, including water and special rat chow. The rats' sleeping and waking times were fixed to be 12 hr in a dark environment and 12 hr in light, and were categorized as shown in table 1. The research was conducted in accordance with accepted principles for laboratory animal use and care.

Rats were put to sleep using ketamine (100 mg/kg) and xylazine (10 mg/kg) before being moved to stereotaxic equipment to produce the model (RWD, 0755-

Table 1. The studied groupings for investigation of *C. aurantium* and *L. angustifolia*

Group name	Number	Gender	Title
G1	3	Male	Healthy
G2	3	Male	Alzheimer
G3	3	Male	Alzheimer + <i>Citrus aurantium</i>
G4	3	Male	Alzheimer + <i>Lavandula angustifolia</i>
G5	3	Male	Alzheimer + <i>Citrus aurantium</i> + <i>Lavandula angustifolia</i>

86111281). The Alzheimer's model was injected and generated using β -Amyloid Peptide (1-42) (Sigma, CAS number: 107761) by injecting in the CA1 region of the hippocampus with the coordinates utilized from the Paxinos atlas (ML=-2.6, AP=-3.8, DV=2.2) and a concentration of 5 $\mu\text{g}/\mu\text{l}$ and 2 μl on each side of the hippocampus using a 5 μl syringe (Hamilton, USA) ¹³. Following the model's introduction, 200 μl of *C. aurantium* and *L. angustifolia* extracts with concentration of 100 mg/kg were administered by gavage over 4 weeks, five days per week ¹⁴⁻¹⁷. The behavioral shuttle box test was conducted 30 days following the last treatment. The rats were subsequently put to death with carbon dioxide gas, and the brain tissue was removed and fixed in 10% formalin ¹³.

Shuttle box test

A shuttle box device (type ST-500) with two chambers-one light and one dark-and bottoms wrapped in steel wire with a diameter of 1 to 2 mm was used for this test. Also, in the dark chamber, a light shock of 75 mA and alternating current for three s was delivered to the mouse's leg just once using an electric current generator. The mice were all first placed in this room for ten min to become used to the open guillotine door, after which they were all moved to the light chamber, and as soon as they entered the dark chamber, the guillotine door was shut and the mice's legs were shocked with electricity. Mice were tested for passive avoidance memory 24 hr later, and the delay time for entering the dark compartment was assessed in seconds ¹⁸.

Congo red test

Once the tissue had been verified, ethanol was employed to dehydrate the subject (starting from low degrees to absolute alcohol). The alcohol was then eliminated by xylol. The samples were then mounted on a silanized slide after being microtome-cut to a thickness of 5 μm . The dehydration procedure and deparaffinization of the slides were done first. They were then immersed in Congo Red 1 solution (Sigma-C-6277) for 30 min, rinsed with distilled water, and then immersed 5-10 times in an alkaline alcohol solution (1% sodium hydroxide and 50% alcohol, Sigma-S0899). After washing the slides in water for 5 min, hematoxylin (Sigma-H9627) for 30 s, and ammonium hydroxide (Sigma-1336-21-6) for 30 s were used and they were once again put in water for washing. Dewatering and clarifying were completed in the end, and an optical microscope (LABOMED) was used to capture images ¹⁹.

ELISA test for norepinephrine and serotonin

One ml of Ripa buffer (Cat. NB: DB9719) was used to homogenize about 100 mg of tissue before centrifuging it for protein extraction. The BCA test was used to identify the supernatant solution that contained protein. After creating the standard graph, the Biotek-refelx800 (US) ELISA equipment was used to evaluate the amounts of norepinephrine and serotonin protein in

the samples according to the ELISA kit's methodology ²⁰.

Superoxide dismutase (SOD) and malondialdehyde (MDA) assay

100 mg of tissue was combined with 1 ml of PBS (Cat. NB: A-0018) before being homogenized and centrifuged. Using the SOD kit (CAT NO: ZB-SOD), the SOD test was conducted on the supernatant of the samples following the kit's instructions. Using a tetrazolium salt that, when reduced with superoxide anion, yields a water-soluble formazan dye, this kit assessed SOD activity. The quantity of produced color was quantified at a wavelength of 440-460 nm with a Biotek-refelx800 (US) ELISA equipment when it was discovered that the presence of SOD in the media inhibits the rate of formazan production. The material was extracted using Ripa buffer, and the MDA kit (ZB-MDA) was used under the kit's instructions to test MDA. Then, using a Biotek-refelx800 (US) ELISA equipment, the quantity of color generated was measured at a wavelength of 535 nm ²¹.

Expression of Amyloid precursor protein (APP) and Glutamate transporter-1 (GLT1) genes

Each sample's total RNA was manually extracted using Triazol (Qiazol, Kiazist) following the manufacturer's instructions from 50 mg of tissue. Simply, 500 ng of RNA from each sample was added to the cDNA synthesis procedure using the Easy cDNA Synthesis Kit (Parstous). Afterwards the extracted RNA was read at a wavelength of 260 nm and a ratio of 260/280 and 230/260 was checked to ensure its quality and purity. Primer3 software was used to create primers for genes (Table 2). Then, using a real-time PCR thermocycler (ABI Stepone, US) and SYBR Master with high Rox (2X) (Addbio), relative changes in the expression of the APP and GLT1 genes in various groups as compared to the control group and GAPDH internal control gene were evaluated ²².

Software called Graph Pad Prism version 9 was utilized to evaluate the tests' statistical significance. ANOVA and Tukay's multiple comparison test were employed in one way. A p-value of 0.05 and a 95% confidence interval were taken into account. Each group included three tested samples. Three repetitions for each sample were run through each test. The results were expressed as mean \pm standard error (mean \pm SE).

Table 2. Primer sequence designed by primer 3 software

Symbol gene	Sequence 5'-3'
rAPP-F	AGATGTCAGAAGTGAAGATGGA
rAPP-R	GCAATGATGGATGGATGTGT
rGLT1-F	AGAGAGGCTGCCGTTAAATAC
rGLT1-R	AAACACAGTCAGTGAGAGCAGG
rGap-R	TGTAGACCATGTAGTTGAGGTCA
rGap-F	AGGTCGGTGTGAACGGATTTG

Results

Phytochemical evaluation

The results of GC–MS for *C. Aurantium* and of *L. angustifolia* are shown in tables 3 and 4. As shown in the table 3, D-Limonen was reported as the active constituent of the *C. Aurantium* hydro-alcoholic extract. Also, the eucalyptol was reported as the active constituent of the *L. angustifolia*.

Shuttle box behavioral test

The findings demonstrated that model induces behavioral impairment so that passive avoidance learning is significantly less than in the control group. Figure 1A, shows that on the first day of measurement, the model group's initial delay time to enter the dark room after the induction of an electric shock was shorter than that of the control group, which is statistically significant at $p < 0.05$. The decrease in the treated groups when compared to the control is not statistically significant, but when compared to the model group, particularly with the treatment using both extracts, it resulted in an increase in learning, which is statistically significant. The experiment was measured again on the second day, and Figure 1B, depicted the same procedure for all control and treatment groups with additional seconds.

To evaluate memory recovery, the Time of Dark Chamber (TDC) was also investigated. Figure 2 shows a statistically significant ($p < 0.05$) increase in TDC in the model group when compared to the control groups. In comparison to the Alzheimer's group, treatment with *C. aurantium* and *L. angustifolia* and both medications simultaneously decreased the amount of time spent in




Figure 1. Initial delay time to enter the dark room after the induction of an electric shock. A) the amount of step through latency (STLa) in seconds on the first day, B) the amount of step through latency (STLr) in seconds on the second day.

the dark room. This difference is statistically significant at $p < 0.05$ for the *L. angustifolia* group and both extracts.

Congo red test

Amyloids in brain tissue were found and identified

Table 3. Chemical composition of Hydro-alcoholic extract of *C. Aurantium*

Number	Compound Name	Retention time (min)	Probability (%)	Percentage (%)
1	D-Limonen	10.43	98	5.54
2	Pyrrolidinone,	10.921	98	1.17
3	Linalool	12.11	91	2.06
4	D-Glucuronic acid	18.840	82	9.53
5	alpha-D-Mannofuranoside	18.954	94	1.72
6	alpha-Guaiene	20.24	92	0.59
7	alpha-D-Glucopyranoside	22.04	47	1.26
8	Daphnetin	22.49	43	3.73
9	Caffeine	25.65	96	4.42

Table 4. Chemical composition of Hydro-alcoholic extract of *L. angustifolia*

Number	Compound name	Retention time (min)	Probability (%)	Percentage (%)
1	Eucalyptol	10.66	98	1.1
2	Camphor	12.78	98	1.1
3	Borneol	13.13	90	2.4
4	Benzene	20.46	50	0.8
5	Benzo[h]quinoline	32.34	38	22.9
6	Silicic acid	32.37	45	5.4




Figure 2. Time of Dark Chamber (TDC) in seconds.

using the Congo red staining method. The outcomes of brain tissue staining in various groups have been shown in figure 3. Figure 4 compares the groups based on the three times each sample's amyloid plaque count was counted. The graphic shows that the Alzheimer's model mice had more amyloid in their brain tissue than the control and other treatment groups. The maximum significance is $p < 0.001$ compared to each extract with the model group in the groups treated with *C. aurantium* and *L. angustifolia* extracts in terms of lowering the amount of amyloid compared to the model group. There are no significant variations between the two extracts.

ELISA test for norepinephrine and serotonin

As shown in figure 5A, after illness induction, there is a significant increase in norepinephrine protein levels in the model group compared to the control group ($p < 0.001$). In the groups treated with *C. aurantium*



Figure 3. The Congo red staining images of a cross-section of brain tissue in mice of different groups, as shown in the figure, with the use of *C. aurantium* and *L. angustifolia* extracts. the number of amyloid plaques (denoted by white arrows) has decreased significantly compared to the sample with Alzheimer's disease. This reduction has been more with the simultaneous use of two extracts. Scale bars for the first column from the left are $200 \mu\text{m}$ and for the second and third columns are $100 \mu\text{m}$.




Figure 4. The number of amyloid plaques in different groups, similar signs are not significantly different. Non-similar signs have a statistically significant difference of $p < 0.05$.




Figure 5. Norepinephrine (A) and Serotonin (B) protein expression levels in different groups, similar signs are not significantly different. Non-similar signs have statistically significant differences ($p \leq 0.001$).

tium and the two extracts, there was a reduction in this protein when compared to the model group; this reduction in protein is significant when compared to the model group. The differences between the therapy groups are negligible. Additionally, figure 5B, shows that following illness induction, the model group's level of serotonin protein decreased significantly ($p < 0.001$) compared to the control group. There is hardly any distinction across the treatment groups. Furthermore, figure 5B demonstrates that, in comparison to the control group, the model group's level of serotonin protein reduced dramatically ($p < 0.001$) after illness induction.

Superoxide dismutase (SOD) and malondialdehyde (MDA) tests

As shown in figure 6A, when compared to the control group, the model group's SOD activity was

Activity of *Citrus aurantium* and *Lavandula angustifolia*




Figure 6. SOD activity (a), and MDA concentration (b), in the studied groups. Similar signs are not significantly different. Non-similar signs have statistically significant differences ($p \leq 0.05$).

significantly lower ($p < 0.001$). When compared to the model group, treatment with *L. angustifolia* extract and both extracts combined significantly increased SOD activity. Among the groups treated with one extract, the increase in SOD activity in the group treated with both extracts relative to the model group showed maximum significance. As shown in figure 6B, in comparison to the control group, the MDA concentration in the model group has significantly increased ($p < 0.001$). MDA concentration was significantly lower after treatment with each extract separately and in combination than it was in the model group. Among the groups treated with one extract, the reduction in MDA levels in the group treated with both extracts relative to the model group showed maximum significance.

Expression measurement of APP and GLT1 genes by Real-time PCR method

As shown in figure 7A, in comparison to the control group, there is a significant increase ($p < 0.01$) in the relative expression of the APP gene in the model group. In comparison to the model group, the groups treated with individual and combined extracts showed a drop in the relative expression of this gene. In the *C. aurantium* group and both extracts together, this decrease is significant. Figure 7B displays the level of GLT1 gene expression in different groups. In comparison to the control group, the relative expression of this gene in the model group has dramatically risen ($p < 0.01$). The relative expression of this gene relative to the model group has significantly decreased in the *L. angustifolia* group and both extracts combined after the effects of the extracts both alone and together. The




Figure 7. Changes in the relative expression of (A) APP and (B) GLT1 genes in different groups. Similar signs are not significantly different. Non-similar signs have statistically significant differences ($p \leq 0.005$).

relative level of expression of both genes has been impacted by both extracts.

Discussion

Numerous studies have demonstrated that food and nutrition can positively influence various pathophysiological consequences in AD so far. Additionally, some studies have suggested that dietary components such as polyphenols may be useful in avoiding and postponing age-related diseases^{23,24}. *C. aurantium* is a traditional herbal remedy that is frequently used as both an antioxidant and an anticancer agent²⁵. Additionally, *L. angustifolia* is recommended in conventional medicine as a therapy for AD. The therapeutic medications employed are mostly anti-oxidants, anti-inflammatory drugs, cholinergic agonists, estrogen, neurotrophic factor, etc. due to the numerous pathogenic factors in AD²⁶. Although organic synthesis can yield two or more pharmacophores related to multifunctional AD inhibitors, the implementation of this strategy is greatly hampered by unpredictable side effects. As a result, researchers' attention has gradually shifted away from organic synthesis and toward multifunctional compounds in natural products²⁷.

As the findings of the current study demonstrated, the behavioral test revealed that although the decrease in the treated groups when compared to the control is not statistically significant, the treatment with both extracts specifically resulted in an increase in learning that is statistically significant when compared to the model group. As the behavioral studies by Zhang *et al*

indicated, *Rhodiola crenulata* Extract (RCE) can enhance cognitive abilities in A β_{1-42} induced AD mice models, including learning and memory²⁸. According to Rabies's study the shuttle box results showed that the initial delay in the Alzheimer's mouse model that was treated with *Cyperus rotundus* ethanolic extract was significantly decreased²⁹. The A β plaque system plays a role in neuropsychiatric processes including memory and learning. Disorders of learning and memory are usually caused by an increase in A β in certain brain areas³⁰. The findings of this research indicated that a significant increase exists in the level of A β in the mouse models' brains, which might be the reason for the cognitive deterioration found in the current study. These defects were corrected by treatment with *C. aurantium* and *L. angustifolia* extract, indicating that lavender and orange spring may improve the A β plaque system's functioning. As a result, *C. aurantium* and *L. angustifolia* may enhance learning and memory in behavioral tests through a variety of processes. The findings of Congo red staining of the brain tissue indicated that the groups treated with *C. aurantium* and *L. angustifolia* extracts had the maximum significance at $p < 0.001$ compared to each extract alone with the model group in terms of lowering the quantity of amyloid plaques. Similarly, Dhanasekaran *et al* demonstrated using Congo red staining that a dosage of 5.0 mg/kg of *Centella asiatica* and a long-term treatment plan can diminish fibrillar amyloid plaques¹⁹. Additionally, research by Soheili's team showed that *L. angustifolia* might significantly decrease the number of amyloid plaques in mice used as an Alzheimer's model³¹.

Numerous practical research has shown that deficiencies in the anti-oxidant defense system and oxidative stress associated with A β are important factors in the pathophysiology and etiology of AD⁹. Compared to the model group, the findings of the current study indicated that treatment with *L. angustifolia* extract and both extracts combined significantly increased SOD activity. Among the groups treated with one extract, the increase in SOD activity in the group treated with both extracts relative to the model group showed maximum significance. Additionally, compared to the model group, treatment with both individual and combined extracts resulted in a significant drop in MDA levels. Zhang *et al* demonstrated that RCE therapy might stop MDA accumulation and the reduction in SOD activity in AD mice models. The present findings are also in line with earlier research, which found that in a rat model of streptozotocin-induced hippocampus cell injury, MDA levels considerably rose along with significant deficits in learning and spatial memory, and the pretreatment with *Rhodiola rosea* extract greatly reduced these aberrations³².

The findings of the ELISA test revealed that a significant increase exists in the protein level of norepinephrine in the model group compared to the

control after the illness was induced. This protein was lower after treatment with each extract alone and after treatment with both extracts compared to the model group. Additionally, after illness induction, the model group's level of serotonin protein decreased significantly ($p < 0.001$) compared to the control group. This protein was significantly elevated following treatment with each extract alone as well as treatment with combined extracts in comparison to the model group. In the study conducted by Rapaka *et al*, pathological and behavioral changes caused by AL-induced Alzheimer's mice were associated with a decrease in dopamine and serotonin levels in the cerebral cortex and hippocampus, and treatment with *Benincasa hispida* hindered this in the AD model³³. Elsayi's study also demonstrated that *Lagerstroemia indica* extract might change the levels of the hormones norepinephrine and serotonin in the Alzheimer's model group in comparison to the control group, respectively³⁴.

In comparison to the model group, the findings reveal that the relative expression of the *APP* gene in the groups treated with the extracts individually and together demonstrate lower expression levels. In the *C. aurantium* group and when the two extracts are combined, this decrease is significant. Furthermore, the *L. angustifolia* group and both extracts taken combined showed a significantly lower expression of *GLT1* following the effects of the extracts than the model group.

Similarly, Yaghmaei group research showed that silymarin extract treatment also lowers the number of amyloid plaques in the brain. Moreover, silymarin was able to reduce *APP* expression as seen by the comparison of treated and untreated groups' *APP* gene expression³⁵. Ji *et al* also discovered that Akt phosphorylation was necessary for insulin stimulation to enhance total and surface *GLT1* expression in cultured astrocytes, which was linked with *KBBP* expression and *GLT1* mRNA transcription³⁶. According to abundant evidence, oxidative stress-induced abnormal and excessive Ca²⁺ release from the Endoplasmic Reticulum (ER) via the Ryanodine Receptor (RyR) which then plays a significant role in memory loss and cognitive dysfunction in AD patients through activation of the enzyme calcineurin phosphatase and increased activities of the enzymes β -calcineurin and γ -secretases in the generation and deposition of A β_{42} senile plaques and the *APP* gene. Inhibiting excessive Ca²⁺ release from the ER, as well as RyR over-activation and calcineurin activation in hippocampus neurons, may therefore be viable therapeutic targets for the treatment of AD³⁷.

Conclusion

The aim of the present study was to better understand the mechanism of action of two plants, *C. aurantium* and *L. angustifolia*, as widely used ethnic herbal medicines, which are widely used in Iran and Mediter-

ranean regions³⁸ to reduce the complications of AD. The results of this research showed that the *C. aurantium* and *L. angustifolia* are able to modulate the expression of *APP* and *GLT1* proteins, decrease oxidative stress, protect neurons from degeneration or apoptosis, and greatly enhance the cognitive performance of AD rat in the behavioral test. Therefore, *C. aurantium* and *L. angustifolia* provide a multi-target therapeutic strategy for AD and can be introduced as nootropic agents to reduce the complications of cognitive diseases such as AD.

Acknowledgement

The authors of this article would like to express their deepest gratitude to Mr. Talebi, who helped to improve research quality, as well as Histogenotec Research Laboratory. The costs are borne by the authors.

Conflict of Interest

The authors declare that they have no conflict of interest.

References

- Mohebbali N, Shahzadeh Fazeli SA, Ghafoori H, Farahmand Z, MohammadKhani E, Vakhshiteh F, et al. Effect of flavonoids rich extract of *Capparis spinosa* on inflammatory involved genes in amyloid-beta peptide injected rat model of Alzheimer's disease. *Nutr Neurosci* 2018;21(2):143-50.
- Ahmad A, Ali T, Park HY, Badshah H, Rehman SU, Kim MO. Neuroprotective effect of fisetin against amyloid-beta-induced cognitive/synaptic dysfunction, neuroinflammation, and neurodegeneration in adult mice. *Mol Neurobiol* 2017;54(3):2269-85.
- Busche MA, Hyman BT. Synergy between amyloid- β and tau in Alzheimer's disease. *Nat Neurosci* 2020;23(10):1183-1193.
- de Oliveira J, Kucharska E, Garcez ML, Rodrigues MS, Quevedo J, Moreno-Gonzalez I, et al. Inflammatory cascade in Alzheimer's disease pathogenesis: A review of experimental findings. *Cells* 2021;10(10):2581.
- Ray B, Chauhan N, K Lahiri D. The "Aged Garlic Extract"(AGE) and one of its active ingredients S-Allyl-LCysteine (SAC) as potential preventive and therapeutic agents for Alzheimer's disease (AD). *Curr Medicinal Chem* 2011;18(22): 3306-13.
- Alzobaidi N, Quasimi H, Emad NA, Alhalmi A, Naqvi M. Bioactive compounds and traditional herbal medicine: Promising approaches for the treatment of dementia. *Degener Neurol Neuromuscul Dis* 2021;11:1.
- Ben Hsouna A, Gargouri M, Dhifi W, Ben Saad R, Sayahi N, Mnif W, et al. Potential anti-inflammatory and antioxidant effects of *Citrus aurantium* essential oil against carbon tetrachloride-mediated hepatotoxicity: A biochemical, molecular and histopathological changes in adult rats. *Environ Toxicol* 2019;34(4): 388-400.
- Maksoud S, Abdel-Massih RM, Rajha HN, Louka N, Chemat F, Barba FJ, Debs E. *Citrus aurantium* L. active constituents, biological effects and extraction methods. an updated review. *Molecules* 2021;26(19):5832.
- Soheili M, Salami M. *Lavandula angustifolia* biological characteristics: An in vitro study. *J Cell Physiol* 2019; 234(9):16424-30.
- Badr MM, Badawy ME, Taktak NE. Characterization, antimicrobial activity, and antioxidant activity of the nanoemulsions of *Lavandula spica* essential oil and its main monoterpenes. *Journal of Drug Delivery Science and Technology* 2021;65:102732.
- Torabbeigi M, Aberoomand Azar P. Analysis of essential oil compositions of *Lavandula angustifolia* by HS-SPME and MAHS-SPME followed by GC and GC-MS. *Acta Chromatographica* 2013;25(3):571-9.
- Azhdarzadeh F, Hojjati M. Chemical composition and antimicrobial activity of leaf, ripe and unripe peel of bitter orange (*Citrus aurantium*) essential oils. *Nutrition and Food Sciences Research* 2016;3(1):43-50.
- Rahman SO, Panda BP, Parvez S, Kaundal M, Hussain S, Akhtar M, et al. Neuroprotective role of astaxanthin in hippocampal insulin resistance induced by A β peptides in animal model of Alzheimer's disease. *Biomed Pharmacother* 2019; 110:47-58.
- Jasim SA, Ali SA-J, Fadhil OQ, Rakhmatova MK, Kzar HH, Margiana R, et al. Investigating the effects of hydro-alcoholic *Urtica dioica* extract and retinoic acid on follicular development: An animal study. *Medi J Islam Repub Iran* 2023;37.
- Adamu SS, Umaru HA, Albert HO, Muhammad AL. The effect of green synthesized zinc oxide nanoparticles using *Allium cepa* extracts on Triton X-100 induced hyperlipidemia in rats. *International Journal of Nutrition Sciences* 2023;8(1):36-46.
- Ogechukwu OC, Salt AP. In vivo antimalarial activity and phytochemical screening of tree bark extract of *Ficus elastica*. *Journal of Science and Technology Research* 2023;5(2).
- Fathollahy I, Barzegar Asl A. Aqueous extract of *Senjed* (*Elaeagnus angustifolia* L.) peel: characteristics and effect on physico-chemical properties of cold-pressed sesame oil. *Journal of Food Measurement and Characterization* 2023:1-10.
- Zavvari F, Karimzadeh F. A review on the behavioral tests for learning and memory assessments in rat. *The Neuroscience Journal of Shefaye Khatam* 2017;5(4):110-124.
- Dhanasekaran M, Holcomb LA, Hitt AR, Tharakan B, Porter JW, Young KA, et al. *Centella asiatica* extract selectively decreases amyloid β levels in hippocampus of Alzheimer's disease animal model. *Phytother Res* 2009; 23(1):14-9.
- Heneka MT, Nadrigny F, Regen T, Martinez-Hernandez A, Dumitrescu-Ozimek L, Terwel D, et al. *Locus ceruleus* controls Alzheimer's disease pathology by modulating microglial functions through norepinephrine. *Proc Natl Acad Sci USA* 2010;107(13):6058-6063.
- Shunan D, Yu M, Guan H, Zhou Y. Neuroprotective effect of betalain against A β 1-42-induced Alzheimer's disease in sprague dawley rats via putative modulation of

- oxidative stress and nuclear factor kappa B (NF- κ B) signaling pathway. *Biomed Pharmacother* 2021;137:111369.
22. Giacomeli R, Izoton JC, Dos Santos RB, Boeira SP, Jesse CR, Haas SE. Neuroprotective effects of curcumin lipid-core nanocapsules in a model Alzheimer's disease induced by β -amyloid 1-42 peptide in aged female mice. *Brain Res* 2019;1721:146325.
 23. Sarubbo F, Moranta D, Asensio VJ, Miralles A, Esteban S. Effects of resveratrol and other polyphenols on the most common brain age-related diseases. *Curr Med Chem* 2017;24(38):4245-66.
 24. Kumar MR, Azizi NF, Yeap SK, Abdullah JO, Khalid M, Omar AR, et al. Clinical and preclinical studies of fermented foods and their effects on Alzheimer's disease. *Antioxidants (Basel)* 2022;11 (5):883.
 25. Sutar I, Khan H, Patel S, Celano R, Rastrelli L. An overview on *Citrus aurantium* L.: Its functions as food ingredient and therapeutic agent. *Oxid Med Cell Longev* 2018 May 2;2018:7864269.
 26. Dey A, Bhattacharya R, Mukherjee A, Pandey DK. Natural products against Alzheimer's disease: Pharmacotherapeutics and biotechnological interventions. *Biotechnol Adv* 2017;35(2):178-216.
 27. Zeng G-f, Zhang Z-y, Lu L, Xiao D-q, Zong S-h, He J-m. Protective effects of ginger root extract on Alzheimer disease-induced behavioral dysfunction in rats. *Rejuvenation Res* 2013;16(2):124-33.
 28. Zhang X, Wang X, Hu X, Chu X, Li X, Han F. Neuroprotective effects of a *Rhodiola crenulata* extract on amyloid- β peptides (A β 1-42)-induced cognitive deficits in rat models of Alzheimer's disease. *Phyto-medicine* 2019;57:331-8.
 29. Rabiei Z, Hojjati M, Rafieian-Kopaeia M, Alibabaei Z. Effect of *Cyperus rotundus* tubers ethanolic extract on learning and memory in animal model of Alzheimer. *Biomedicine & Aging Pathology* 2013;3(4):185-91.
 30. Kent SA, Spires-Jones TL, Durrant CS. The physiological roles of tau and A β : implications for Alzheimer's disease pathology and therapeutics. *Acta neuropathologica* 2020;140(4):417-47.
 31. Soheili M, Tavirani MR, Salami M. *Lavandula angustifolia* extract improves deteriorated synaptic plasticity in an animal model of Alzheimer's disease. *Iran J Basic Med Sci* 2015;18(11):1147.
 32. Yu H, Zhang P, Zhang C, Zhang X, Li Z, Li W, Fu A. [Effects of *rhodiola rosea* on oxidative stress and negative emotional states in patients with obstructive sleep apnea]. *Lin chuang er bi yan hou tou jing wai ke za zhi= Journal of clinical otorhinolaryngology, head, and neck surgery* 2019;33(10):954-957. Chinese.
 33. Rapaka D, Bitra VR, Ummidi R, Akula A. *Benincasa hispida* alleviates amyloid pathology by inhibition of Keap1/Nrf2-axis: Emphasis on oxidative and inflammatory stress involved in Alzheimer's disease model. *Neuropeptides* 2021; 88:102151.
 34. Elsawi SA, Aly HF, Elbatany MM, Maamoun AA, Mowawad DM. Phytochemical evaluation of *Lagerstroemia indica* (L.) Pers leaves as anti-Alzheimer's. *J Mater Environ Sci* 2018;9(9):2575-86.
 35. Yaghmaei P, Azarfar K, Dezfulian M, Ebrahim-Habibi A. Silymarin effect on amyloid- β plaque accumulation and gene expression of APP in an Alzheimer's disease rat model. *Daru* 2014;22 (1):24.
 36. Ji Y-F, Xu S-M, Zhu J, Wang X-X, Shen Y. Insulin increases glutamate transporter GLT1 in cultured astrocytes. *Biochem Biophys Res Commun* 2011;405(4):691-696.
 37. Dutta BJ, Singh S, Seksaria S, Gupta GD, Bodakhe SH, Singh A. Potential role of IP3/Ca2+ signaling and phosphodiesterases: Relevance to neurodegeneration in Alzheimer's disease and possible therapeutic strategies. *Biochem Pharmacol* 2022;201:115071.
 38. Shahrajabian MH, Sun W. Survey on medicinal plants and herbs in traditional Iranian medicine with anti-oxidant, anti-viral, anti-microbial, and anti-inflammation properties. *Letters in Drug Design & Discovery* 2023;20 (11):1707-43.