



## The Novel Drug Discovery to Combat COVID-19 by Repressing Important Virus Proteins Involved in Pathogenesis Using Medicinal Herbal Compounds

Samira Mahmoudi <sup>1</sup>, Negar Balmeh <sup>2</sup>, Niloofar Mohammadi <sup>3</sup>, and Tahereh Sadeghian-Rizi <sup>4\*</sup>

1. Department of Microbial Biotechnology, Faculty of Biological Sciences, Tehran North Branch, Islamic Azad University, Tehran, Iran

2. Department of Cell and Molecular Biology, Faculty of Biology, Nour Danesh Institution of Higher Education, Meymeh, Iran

3. Department of Sciences and Agricultural Engineering, School of Agricultural Sciences, Pir Bakran Branch, Payame Noor University, Isfahan, Iran

4. Department of Biotechnology, Faculty of Biological Sciences and Technology, Shahid Ashrafi Esfahani University, Isfahan, Iran

### Abstract

**Background:** The cause of COVID-19 global pandemic is SARS-CoV-2. Given the outbreak of this disease, it is so important to find a treatment. One strategy to cope with COVID-19 is to use the active ingredients of medicinal plants. In this study, the effect of active substances was surveyed in inhibiting four important druggable targets, including S protein, 3CLpro, RdRp, and N protein. RdRp controls the replication of SARS-CoV-2 and is crucial for its life cycle. 3CLpro is the main protease of the virus and could be another therapeutic target. Moreover, N protein and S protein are responsible for SARS-CoV-2 assembly and attaching, respectively.

**Methods:** The 3D structures of herbal active ingredients were prepared and docked with the mentioned SARS-CoV-2 proteins to obtain their affinity. Then, available antiviral drugs introduced in other investigations were docked using similar tools and compared with the results of this study. Finally, other properties of natural compounds were uncovered for drug designing.

**Results:** The outcomes of the study revealed that Linarin, Amentoflavone, (-)-Catechin Gallate and Hypericin from *Chrysanthemum morifolium*, *Hypericum perforatum*, *Humulus lupulus*, and *Hibiscus sabdariffa* had the highest affinity for these basic proteins and in some cases, their affinity was much higher than antiviral medicines.

**Conclusion:** In addition to having high affinity, these herb active ingredients have antioxidant, vasoprotective, anticarcinogenic, and antiviral properties. Therefore, they can be used as extremely safe therapeutic compounds in drug design studies to control COVID-19.

*Avicenna J Med Biotech 2021; 13(3): 107-115*

**Keywords:** COVID-19, Drug design, Medicinal plants, SARS-CoV-2

### Introduction

Coronavirus disease 2019 (COVID-19) emerged in late 2019 by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) and posed a global public health emergency. SARS-CoV-2 is a new coronavirus and the World Health Organization (WHO) firstly named it 2019-nCoV <sup>1,2</sup>. The reason for naming coronavirus SARS-CoV-2 is the similarities of its genomic sequence to severe acute respiratory syndrome coronavirus (SARS-CoV) and the disease was subsequently called COVID-19 <sup>3</sup>. Coronaviruses, a genus of the

Coronaviridae family, are classified as single-stranded, positive-sense RNA viruses, and their genomes range from 25-32 kb <sup>4</sup>. According to the results of genome-based phylogenetic analysis, SARS-CoV-2 is a new branch of the betacoronavirus genus. SARS-CoV and the Middle East Respiratory Syndrome (MERS) are also from the betacoronavirus genus. It is noteworthy that SARS and MERS also appeared in 2002 and 2012, respectively <sup>5,6</sup>. SARS-CoV caused an epidemic in China and MERS began in Saudi Arabia <sup>7</sup>. The genome

\* **Corresponding author:**  
Tahereh Sadeghian-Rizi, Ph.D.,  
Department of Biotechnology,  
Faculty of Biological Sciences and  
Technology, Shahid Ashrafi  
Esfahani University, Esfahan, Iran  
**Tel:** +98 9132366325  
**E-mail:**  
t.sadeghian@ashrafi.ac.ir  
**Received:** 7 Oct 2020  
**Accepted:** 4 Jan 2021

of SARS-CoV-2 codes for Nonstructural Proteins (NSPs) such as 3-chymotrypsin-like protease (3CLpro), RNA-dependent RNA polymerase (RdRp), accessory proteins, and Structural Proteins (SPs) <sup>8</sup>. SPs include Nucleocapsid (N) protein, Membrane (M) protein, Envelope (E) protein, and Spike (S) protein <sup>9</sup>.

The S protein is responsible for virus-cell interactions during viral entry <sup>10</sup>. S1 and S2 are the two subunits of the S protein. S1 is at the N-terminus and responsible for the virus-receptor binding while S2 is at the C-terminus and responsible for virus-cell membrane fusion <sup>11,12</sup>. S1 itself is divided into two domains called the Receptor-Binding Domain (RBD) and the N-Terminal Domain (NTD) <sup>7</sup>. In fact, SARS-CoV-2 attaches to cells by its S protein and human Angiotensin-Converting Enzyme 2 (ACE2) <sup>13-15</sup>. This binding triggers a cascade of events and leads to a fusion between viral and cellular membranes. Then, RNA viral genome is released into the cytoplasm of host cells <sup>16</sup>. The next step is synthesizing polyproteins for encoding the viral replicase-transcriptase complex <sup>17</sup>. So, targeting S protein could be an initial solution to prevent the infection.

The N protein is in the helical nucleocapsid and has an important role in replication, RNA packaging, and pathogenesis <sup>4</sup>. N protein is more conserved compared to other SPs and more expressed throughout COVID-19 infection; thus, it is one of the immunodominant antigens in SARS-CoV-2 <sup>18,19</sup>. Therefore, N protein inhibition seems to be one of the fastest and most effective actions to prevent the infection progression in patients infected with COVID-19.

During transcription of the SARS-CoV-2 genome, the virus produces a ~800 *kDa* polypeptide and cleaves this polypeptide by 3CLpro to generate NSPs <sup>20</sup>. 3CLpro converts this polypeptide to eleven fragments. The 3CLpro is located at the 3' end of the genome that has excessive variability <sup>21</sup>. In fact, 3CLpro has a critical role in the virus particles' replication so this protein is a potential target for inhibiting SARS-CoV-2 <sup>22</sup>.

RdRp is another important protein for SARS-CoV-2 with a range of 240-450 *kDa* and a catalytic core with differentiated domains <sup>23</sup>. RdRp catalyzes the RNA replication from its RNA templates, *i.e.*, the viral RNA is synthesized by RdRp. The sequences of RdRp in SARS-CoV, SARS-CoV-2, and MERS-CoV are almost similarly encoded <sup>24</sup>. RdRp active site is a deep and huge groove for polymerization of RNA <sup>25</sup>. This active site is the most accessible and conserved region, so targeting this region may be an effective therapeutic target to inhibit viral replication <sup>26</sup>.

SPs and NSPs are both promising targets to inhibit the virus; while SPs are important in the infection processes, NSPs are involved in virus replication and transcription <sup>27</sup>.

Many of the available antiviral drugs undergoing clinical trials are actually being tested for repurposing, such as chloroquine and Hydroxychloroquine, which

are used to cure arthritis and malaria, respectively <sup>28</sup>, and are currently being used in patients with COVID-19 <sup>29</sup>. Multiple bioinformatics studies have even been performed to find the inhibitory effects of various antiviral chemicals on SARS-CoV-2. For instance, in an *in silico* study, a drug-repurposing investigation was performed on quinoline-based inhibitors against three proteins, S protein, 3CLpro, and RdRp. The results exhibited that Saquinavir and quinoline,1,2,3,4-tetrahydro-1-[(2-phenylcyclopropyl)sulfonyl]-trans-(8CI) inhibited 3CLpro, oxolinic acid and elvitegravir inhibited RdRp, and rilapladib inhibited S protein <sup>30</sup>. But no therapeutic drug has yet been approved by the FDA that is applied exclusively for these patients <sup>29</sup>. As a result, it is necessary to perform structural based analyses to identify inhibitors for SARS-CoV-2 proteins.

In recent years, medicinal plants have attracted significant attention to treat infectious diseases. Complex molecular structures and a wide variety of natural compounds make medicinal plants an excellent biological resource for drug discovery. So far, a limited number of plant species have been used for medicinal purposes, so there is much hope for their increasing use in the future <sup>31</sup>. Flavonoids are natural compounds that have a wide range of functions, including promoting general immunity, blocking transcription and translation of pathogenic microorganisms, and being non-toxic to human cells. These factors have led to the notion of flavonoids as antiviral agents in various articles and studies <sup>32</sup>. A laboratory study examined the inhibitory effect of flavonoid compounds and found that pectolarin, herbacetin, and rhoifolin strongly blocked the 3CLpro of SARS-CoV <sup>33</sup>. Thus, in this article, the inhibitory effect of active ingredients of different medicinal plants was investigated on four important virus proteins including S protein, N protein, 3CLpro, and RdRp. These plant-based substances have characteristics such as fewer side effects, cost-effectiveness, and greater availability. The results of this *in silico* approach showed that some of these studied active ingredients have a high affinity for each of the four important viral proteins compared to the inhibitors previously reported for each of these proteins. They could possibly have an inhibitory effect on the SARS-CoV-2 virus and COVID-19.

## Materials and Methods

### Modeling and structural analysis

After obtaining all information about SARS-CoV-2 structure and genome through the ViralZone database ([www.viralzone.expasy.org](http://www.viralzone.expasy.org)) <sup>34</sup>, the sequences of S protein, 3CLpro, RdRp, and N protein were acquired from the NCBI database (<https://www.ncbi.nlm.nih.gov/>) <sup>35</sup> and modeled by the SWISS-MODEL database (<https://swissmodel.expasy.org/>) <sup>36</sup>. Also, the RCSB PDB database (<https://www.rcsb.org/>) was used to download the SARS-CoV-2 structure <sup>37</sup>.

### Collecting the active substances of medicinal herbs

In this study, the active ingredients of medicinal plants were obtained from articles that had previously introduced the active ingredients in each plant using the GC-MS technique. Plants were selected based on their potential or approved antimicrobial properties as well as the information about their active ingredients. Criteria for selecting effective ingredients included having an equal affinity or greater than the available inhibitors to the mentioned proteins (Table 1).

Then, 3D structures of medicinal herbal substances were obtained from the PubChem database (<https://pubchem.ncbi.nlm.nih.gov/>)<sup>38</sup>.

### Molecular analysis

These obtained 3D structures and that of viral proteins were prepared by docking methods through removing optimal H-bond and all water and adding Gasteiger charges. Then, molecular docking analysis was conducted between each herbal compound and viral proteins through AutoDock vina. Figures were created by Chimera software, version 1.14<sup>39</sup>. Also, to ensure the suitability of the proposed herbal compounds in this study, the blockers and antiviral drugs introduced in other studies were docked with selected viral proteins.

### Investigation of characteristics of high scoring compounds

Finally, Way2Drug ([www.pharmaexpert.ru/pass-online/predict.php](http://www.pharmaexpert.ru/pass-online/predict.php))<sup>40</sup>, lazar ([lazar.in-silico.ch/predict](http://lazar.in-silico.ch/predict))<sup>41</sup>, and AVCpred (<http://crdd.osdd.net/servers/avcpred>)<sup>34</sup> were used to obtain the activity, daily dose carcinogenicity, blood-brain barrier penetration, and antiviral features, respectively.

## Results

The best herbal compounds for S protein, 3CLpro, RdRp, and N protein were recognized based on the highest binding affinity. The affinity of reported viral proteins' blockers was obtained and compared with the top scoring natural compounds examined in the present study. The results presented in table 2 have shown that top scoring medicinal herbal compounds for 3CLpro and RdRp had more affinity compared with previously reported blockers. Also, information on all plant chemicals docked with N, 3CL, S, and RdRp proteins is given in tables 1-4, respectively.

The most common medicinal herbal compounds for all mentioned viral proteins were investigated in this study. Among them, Amentoflavone had the highest binding energy with the scores of -8, -8.9, and -9.4 Kcal/mol for S protein, 3CLpro, and RdRp, respectively, and Linarin was the best top scoring compound for interaction with SARS-CoV-2 N protein (Score of -8.1 Kcal/mol) (Table 3).

The important amino acids involved in the binding of mentioned viral proteins that were achieved from previous articles and the interaction of H-bond related to top scoring compound for blocking each viral protein are shown in figure 1.

Based on the lazar database, characteristics of common compounds including respiratory analeptic, anti-tussive, antiviral, RNA synthesis inhibitor, and hepatoprotection were investigated. Other specifications such as carcinogenicity, blood-brain barrier penetration, herbal compounds antiviral ability, and maximum

Table 1. Sixty-four traditional medicinal herbs used in the survey

Name of herbs		
<i>Alpinia officinarum</i>	<i>Cuminum cyminum</i>	<i>Olea europaea</i>
<i>Artemisia dracunculus</i>	<i>Clausena lansium</i>	<i>Plantago lanceolata</i>
<i>Artemisia absinthium</i>	<i>Chrysanthemum</i>	<i>Passiflora caerulea</i>
<i>Apium graveolens</i>	<i>Datura stramonium L</i>	<i>Pimpinella anisum</i>
<i>Astragalus hamosus</i>	<i>Dianthus barbatus</i>	<i>Punica granatum</i>
<i>Acorus calamus</i>	<i>Dracocephalum</i>	<i>Plantago ovata</i>
<i>Anethum graveolens dhi</i>	<i>Elaeagnus angustifolia</i>	<i>Paeonia</i>
<i>Achillea millefolium</i>	<i>Foeniculum vulgare</i>	<i>Rhus</i>
<i>Allium sativum</i>	<i>Ferula assa-foetida</i>	<i>Salvia officinalis</i>
<i>Bacopa</i>	<i>Glycine max</i>	<i>Silybum marianum</i>
<i>Boswellia</i>	<i>Glycyrrhiza glabra</i>	<i>Taraxacum</i>
<i>Beberis vulgaris (Berberidaceae)</i>	<i>Hibiscus sabdariffa</i>	<i>Thymes</i>
<i>Cistanche tubulosa</i>	<i>Hypericum perforatum</i>	<i>Trigonella foenumgraecum</i>
<i>Cassia fistula</i>	<i>Humulus lupulus</i>	<i>Tanacetum parthenium</i>
<i>Centella asiatica</i>	<i>Hymenocrater</i>	<i>Trigonella foenum-graecum</i>
<i>Crocus sativus</i>	<i>Juglans regia</i>	<i>Thymus kotschyanus</i>
<i>Curcuma longa</i>	<i>Jasminum</i>	<i>Taxus baccata</i>
<i>Cuscuta</i>	<i>Lavandula</i>	<i>Valeriana officinalis</i>
<i>Coriandrum sativum</i>	<i>Lavandula angustifolia</i>	<i>Ziziphora</i>
<i>Cinnamomum verum</i>	<i>Matricaria chamomilla</i>	<i>Zingiber officinale</i>
<i>Carum carvi</i>	<i>Mespilus germanica</i>	
<i>Citrus bergamia Risso</i>	<i>Nigella sativa</i>	

## Drug Discovery to Combat COVID-19

Table 2. The binding affinity of top scoring chemical compounds for blocking each viral protein (N protein, 3CL, S protein, and RdRp) compared to available or proposed natural and antiviral drugs

Top scoring herbal compounds for blocking N protein	Herb name	Docking score (kCal/mol)
Linarin	<i>Chrysanthemum morifolium</i>	-8.1
Isorhoifolin	<i>Olea europaea</i>	-8.1
N protein blocker	Blocker	Docking score (kCal/mol)
Conivaptan	Natural compound	-8.3
Amyrin	Natural compound	-7.9
Top scoring herbal compounds for blocking 3CLpro	Herb name	Docking score (kCal/mol)
(-)-Catechin Gallate	<i>Humulus lupulus</i>	-9.2
(-)-Epicatechin Gallate	<i>Humulus lupulus</i>	-9.2
(-)-Epigallocatechin Gallate	<i>Camellia sinensis</i>	-9.2
Hypericin	<i>Hypericum perforatum/Hibiscus sabdariffa</i>	-9.2
Oxycanthine	<i>Berberis integerrima</i>	-9.1
Pseudohypericin	<i>Hypericum perforatum/Hibiscus sabdariffa</i>	-9
Delphinidin 3-O-Beta-D-sambubioside	<i>Hibiscus sabdariffa</i>	-9
3CLpro blocker	Blocker	Docking score (kCal/mol)
Prulifloxacin	Antiviral drug	-8.3
Nelfinavir	Antiviral drug	-8.2
Raltegravir	Antiviral drug	-8.1
Dolutegravir	Antiviral drug	-8.0
Darunavir	Antiviral drug	-8.0
Bictegravir	Antiviral drug	-7.9
Top scoring herbal compounds for blocking S protein	Herb name	Docking score (kCal/mol)
Amentoflavone	<i>Hypericum perforatum</i>	-8
S protein blocker	Blocker	Docking score (kCal/mol)
Loniflavone	Natural compound	-9.0
Top scoring herbal compounds for blocking RdRp	Herb name	Docking score (kCal/mol)
Amentoflavone	<i>Hypericum perforatum</i>	-9.4
Hypericin	<i>Hypericum perforatum/Hibiscus sabdariffa</i>	-9.1
Neeroicitrin	<i>Citrus bergamia Risso</i>	-9
RdRp blocker	Blocker	Docking score (kCal/mol)
Theaflavin	Natural compound	-9.3
Setrobuvir	Antiviral drug	-8.5
Remdesivir	Antiviral drug	-7.8
Cefuroxime	Antiviral drug	-7.4
Sofosbuvir	Antiviral drug	-7.1
Yak	Antiviral drug	-7.0
IDX-184	Antiviral drug	-6.7
Galidesivir	Antiviral drug	-6.4
Tenofovir	Antiviral drug	-6.4
Ribavirin	Antiviral drug	-6.1
Hydroxychloroquine	Antiviral drug	-6.1
Favipiravir	Antiviral drug	-5.1

recommended daily dose were examined and indicated in table 4.

### Discussion

At present, researchers around the world are examining several different technologies, medicines, and vaccines against COVID-19. So far, ten candidate vaccines have been administered to volunteers in safety trials<sup>42</sup>, among which one has reached the second phase of human trials<sup>43</sup>. However, there is still no conclusive evidence that vaccines can make people completely safe; so along with vaccine production, special

attention should be paid to therapeutic drug development<sup>44</sup>.

Blocking the steps of the viral lifecycle provides potential targets to cure COVID-19 infection. SARS-CoV-2 has two types of proteins, including SPs and NSPs. Both proteins are important for virus survival and the infection process, so preventing the action of these proteins could be helpful to create anti-SARS-CoV-2 drugs<sup>45</sup>. Inhibition of S and N proteins can be a quick solution for preventing SARS-COV-2 infection<sup>46</sup> and further, given that 3CLpro and RdRp are involved in virus replication and development, it is nec-

Table 3. The binding affinity of the most common medicinal herbal compounds for each viral protein (N protein, 3CL, S protein, and RdRp)

Common herbal compounds for blocking 3CLpro	Herb name	Docking score (kCal/mol)
Hypericin	<i>Hypericum perforatum/Hibiscus sabdariffa</i>	-9.2
Amentoflavone	<i>Hypericum perforatum</i>	-8.9
Hesperidin	<i>Ziziphora</i>	-8.7
Linarin	<i>Chrysanthemum morifolium</i>	-8.6
Neeriocitrin	<i>Citrus bergamia Risso</i>	-8.4
Hispaglabridin B	<i>Glycyrrhiza glabra</i>	-8.1
Common herbal compounds for blocking RdRp	Herb name	Docking score (kCal/mol)
Amentoflavone	<i>Hypericum perforatum</i>	-9.4
Hypericin	<i>Hypericum perforatum/Hibiscus sabdariffa</i>	-9.1
Neeriocitrin	<i>Citrus bergamia Risso</i>	-9
Linarin	<i>Chrysanthemum morifolium</i>	-8.5
Hesperidin	<i>Ziziphora</i>	-8.2
Hispaglabridin B	<i>Glycyrrhiza glabra</i>	-8
Common herbal compounds for blocking N Protein	Herb name	Docking score (kCal/mol)
Linarin	<i>Chrysanthemum morifolium</i>	-8.1
Hesperidin	<i>Ziziphora</i>	-7.6
Amentoflavone	<i>Hypericum perforatum</i>	-7.4
Hispaglabridin B	<i>Glycyrrhiza glabra</i>	-7.2
Hypericin	<i>Hypericum perforatum/Hibiscus sabdariffa</i>	-7.1
Neeriocitrin	<i>Citrus bergamia Risso</i>	-7
Common herbal compounds for blocking S Protein	Herb name	Docking score (kCal/mol)
Amentoflavone	<i>Hypericum perforatum</i>	-8
Hypericin	<i>Hypericum perforatum</i>	-7.3
Linarin	<i>Chrysanthemum morifolium</i>	-7.2
Hispaglabridin B	<i>Glycyrrhiza glabra</i>	-7.2
Neeriocitrin	<i>Citrus bergamia Risso</i>	-7.2
Hesperidin	<i>Ziziphora</i>	-7.1

essary to identify novel compounds that can repress the 3CLpro and RdRp<sup>47,48</sup>.

There are two strategies for preventing SARS-CoV-2 interaction with the host cell. Blocking virus receptors (ACE2, TMPRSS2, and GRP78) on the surface of host cells and inhibiting virus proteins are the two methods. In our previous study, the effect of 1032 medicinal herbal compounds on inhibiting SARS-CoV-2 receptors was investigated and based on the results, Berbamine, Hesperidin, and Hypericin had the highest affinity to target receptors of S protein<sup>49</sup>. Subsequently, in the present study, previous investigations were extended to screen various medicinal herbal compounds against S protein, 3CLpro, RdRp, and N protein of the virus.

Some studies stated that a few chemical drugs such as Colistin, Nelfinavir, and Prulifloxacin could be used to target SARS-CoV-2 3CLpro<sup>50,51</sup>. One study examined the inhibitory effect of existing antiviral compounds. This research indicated the interaction of about 123 antiviral drugs against 3CLpro and Paritaprevir, Raltegravir, Bictegravir, and Dolutegravir<sup>52</sup>. In an investigation, a medicinal plant database containing nearly 32,297 antiviral phytochemicals was screened, and outcomes approved that 5,7,30,40-tetrahydroxy-2'-

(3,3-dimethylallyl) isoflavone had the highest affinity to 3CLpro<sup>22</sup>. An *in silico* study performed computational drug design methods to determine inhibitors of 3CLpro from available FDA approved antiviral medicines. As a result, three antiviral drugs including Saquinavir, Remdesivir, Darunavir, and two natural compounds including coumarin derivatives and flavone were promising hits and can inhibit SARS-CoV-2 3CLpro function<sup>53</sup>.

Based on a molecular dynamics study, inhibitors were tested against SARS-CoV-2 RdRp and it was revealed that Ribavirin, Sofosbuvir, Remdesivir, Galidesivir, Cefuroxime, Favipiravir, Hydroxychloroquine, and Tenofovir could tightly attach to the RdRp active site. Also, they presented that YAK, Setrobuvir, and IDX-184 can tightly wrap to RdRp and eradicate the virus<sup>48</sup>. In addition to many studies performed to detect the inhibitory effect of chemical compounds on RdRp protein, research has been done to find the effect of traditional Chinese medicinal compounds on this protein. For instance, a research group screened some medicinal compounds and found that Theaflavin had lower binding energy while docking with SARS-CoV-2 RdRp, thus, Theaflavin can be a potential inhibitor for it<sup>25</sup>. Since the advent of SARS-CoV-2, much

Table 4. Different characteristics of common medicinal herbal compounds having a high affinity to N protein, 3CL, S protein, and RdRp

Compound name	Activity	Carcinogenicity in mouse	Penetration from blood-brain barrier in human	Predicted percentage	Maximum daily dose in human
Linarin	Vasoprotector, hemostatic, cardioprotectant, free radical scavenger, anticarcinogenic, chemopreventive, antihypercholesterolemic, antiprotozoal (Leishmania), hepatoprotectant, anaphylatoxin receptor antagonist, effective for proliferative diseases treatment, TP53 expression enhancer, antineoplastic, antioxidant, antihemorrhagic, antifungal, antiviral (Influenza), anti-infective, radioprotector, anti-inflammatory, antimycobacterial, antithrombotic, platelet adhesion inhibitor	Not carcinogenic	-	54.176	-
Hesperidin	Hepatoprotectant, free radical scavenger, chemopreventive, membrane permeability inhibitor, vasoprotector, Caspase 3 stimulant, hemostatic, cardioprotectant, membrane integrity agonist, effective for proliferative diseases treatment, skin whitener, antihypercholesterolemic, antineoplastic, laxative, antiprotozoal (Leishmania), TP53 expression enhancer, antioxidant, antiulcerative, antifungal, vasodilator, expectorant, Caspase 8 stimulant, antihemorrhagic, anti-inflammatory, antitussive, antibacterial, apoptosis agonist, respiratory analeptic, effective in hepatic diseases treatment, RNA synthesis inhibitor, antiviral, anti-infective	Not carcinogenic	No penetration	54.176	13.5 (mg/kg_bw/day)
Amentoflavone	Kinase inhibitor, TP53 expression enhancer, anaphylatoxin receptor antagonist, histidine kinase inhibitor, antimutagenic, peroxidase inhibitor, apoptosis agonist, vasoprotector, antiseborrheic, antineoplastic, histamine release inhibitor, antihemorrhagic, JAK2 expression inhibitor, antineoplastic (Breast cancer), anticarcinogenic, antiseptic, chemopreventive, mucomembranous protector, free radical scavenger, hemostatic, antioxidant, kidney function stimulant, hepatoprotectant, radiosensitizer	Not carcinogenic	-	-	-
Hispaglabridin B	Anti-infective, spasmolytic, chemopreventive, free radical scavenger, apoptosis agonist, skin whitener, antineoplastic, antidyskinetic, TP53 expression enhancer, Myc inhibitor	-	-	-	-
Hypericin	TP53 expression enhancer, membrane integrity agonist, antineoplastic, apoptosis agonist, membrane permeability inhibitor, antiseborrheic, Caspase 3 stimulant, kidney function stimulant, antiseptic, vasoprotector, antimutagenic	-	-	-	0.022 (mmol/kg_bw/day)
Neoeriocitrin	anticarcinogenic, free radical scavenger, lipid peroxidase inhibitor, hepatoprotectant, chemopreventive, proliferative diseases treatment, vasoprotector, antihypercholesterolemic, anaphylatoxin receptor antagonist, antiprotozoal (Leishmania), TP53 expression enhancer, hemostatic, skin whitener, cardioprotectant, antioxidant, antineoplastic, laxative, effective in capillary fragility treatment, anti-infective, antiviral (Influenza), antifungal, respiratory analeptic, expectorant, interleukin 6 antagonist, antitussive, sweetener, apoptosis agonist, analeptic, antithrombotic, anti-inflammatory, antibacterial, effective in hepatic diseases treatment, antiviral (Herpes)	Not carcinogenic	-	-	-

research has been done on the inhibitory effect of Remdesivir on RdRp<sup>54</sup> and ultimately, its effect was approved by FDA<sup>55</sup>. A study detected top natural compounds by virtual screening so that Loniflavone and Simeprevir had the highest affinity to S protein; also, Amyrin and Conivaptan had the highest affinity to the N protein<sup>56</sup>.

According to the results of the present study, Linarin with the score of -8.1 *kCal/mol*, Catechin Gallate with the score of -9.2 *kCal/mol*, Amentoflavone with the score of -8, and Amentoflavone with the score of -9.4 *kCal/mol* had a higher affinity for N protein, 3CLpro, S protein, and RdRp, respectively. Also, the active ingredient Amentoflavone obtained from *Hyper-*

*icum perforatum* plant had a relatively high affinity for all four important proteins of SARS-CoC-2, so this active ingredient may prevent the binding, entry, multiplication, and development of the virus.

Comparison of the binding affinity of previously reported blockers for each four mentioned viral proteins and top scoring herbal compounds investigated in this survey revealed that the compounds obtained in this study have a greater tendency to bind to virus proteins in some cases. The antiviral drug Prulifloxacin had a score of -8.3 *kCal/mol* for binding to the 3CLpro protein, while (-)-Catechin Gallate had by far more tendency to bind to this protein, with a score of -9.2 *kCal/mol*. In the case of RdRp, the antiviral drug Theaflavin




Figure 1. Affinity of top scoring medicinal herbal compounds to inhibit the S protein, 3CLpro, RdRp, and N protein. A) the interaction between Amentoflavone from *Hypericum perforatum* (Purple) and S protein (Orange red); B) the interaction between Linarin from *Chrysanthemum morifolium* (Deep pink) and N protein (Light sea green); C) the interaction between Amentoflavone from *Hypericum perforatum* (Orang) and 3CLpro (Olive drab); D) the interaction between Amentoflavone from *Hypericum perforatum* (Violet red) and RdRp (Sandy brown).

had the  $-9.3$   $kCal/mol$  docking score, while Amentoflavone had the  $-9.4$   $kCal/mol$  docking score. It should be noted that Remdesivir, which has been already approved by the FDA to inhibit RdRp protein, has a much lower score ( $-7.8$   $kCal/mol$ ) than Amentoflavone.

In addition to having fewer side effects, Linarin and Amentoflavone are not carcinogenic. Linarin and Amentoflavone have other properties as they are antioxidants, vasoprotective, anticarcinogenic, and antiviral, which make them an extremely safe therapeutic compound in the respiratory tract.

According to our previous study on the effect of herbal active ingredients in inhibiting receptors of SARS-CoV-2 and their effect on viral proteins in the present study, it was found that Hypericin as an herbal compound has a high affinity for ACE2, TMPRSS2, and GRP78 receptors and four viral proteins, thus, it can be an important blocker/inhibitor for SARS-CoV-2 proteins and target cell receptors in treatment and control of COVID-19. Therefore, laboratory studies of this effective substance are a great help in discovering the proper medicine for the epidemic disease.

### Conclusion

Most studies have examined one or two proteins of SARS-CoV-2, but in this research, the inhibitory effect of medicinal herbal compounds was investigated on S protein, N protein, 3CLpro, and RdRp simultaneously so that basic proteins of SPs and NSPs could be inhibited and definitively the virus would be eradicated. It can be concluded that Linarin, Amentoflavone, (-)-Catechin Gallate, and Hypericin may have an effective role in SARS-CoV-2 infection. In addition to having a great affinity to attach to the viral proteins, these herbal compounds have antioxidant, vasoprotective, anticarcinogenic, and antiviral properties. Thus, they can be applied as extremely safe therapeutic natural com-

pounds and clinical assessments might have notable outcomes for controlling COVID-19.

### Acknowledgement

The authors gratefully acknowledge AJMB Editorial office for English editing of this paper.

### Conflict of Interest

Authors declare no conflict of interest.

### References

- Zhou P, Yang X-L, Wang X-G, Hu B, Zhang L, Zhang W, et al. A pneumonia outbreak associated with a new coronavirus of probable bat origin. *Nature* 2020;579(7798):270-3.
- Wu F, Zhao S, Yu B, Chen YM, Wang W, Song ZG, et al. A new coronavirus associated with human respiratory disease in China. *Nature* 2020;579(7798):265-9.
- Coronaviridae Study Group of the International Committee on Taxonomy of Viruses: Gorbalenya AA, Baker SC, Baric RS, de Groot RJ, Drosten C, Gulyaeva AA, et al. The species severe acute respiratory syndrome-related coronavirus: classifying 2019-nCoV and naming it SARS-CoV-2. *Nat Microbiol* 2020; 2020:536-44.
- Malik YA. Properties of Coronavirus and SARS-CoV-2. *Malays J Pathol* 2020;42(1):3-11.
- Zaki AM, Van Boheemen S, Bestebroer TM, Osterhaus ADME, Fouchier RAM. Isolation of a novel coronavirus from a man with pneumonia in Saudi Arabia. *N Engl J Med* 2012;367(19):1814-20.
- De Wit E, Van Doremalen N, Falzarano D, Munster VJ. SARS and MERS: recent insights into emerging coronaviruses. *Nat Rev Microbiol* 2016;14(8):523-34.
- Liu X, Liu C, Liu G, Luo W, Xia N. COVID-19: Progress in diagnostics, therapy and vaccination. *Theranostics* 2020;10(17):7821-35.
- Li G, De Clercq E. Therapeutic options for the 2019 novel coronavirus (2019-nCoV). *Nature Publishing Group*; 2020.
- Lisi L, Lacal PM, Barbaccia ML, Graziani G. Approaching Coronavirus Disease 2019: mechanisms of action of repurposed drugs with potential activity against SARS-CoV-2. *Biochem Pharmacol* 2020;114:169.
- Walls AC, Park YJ, Tortorici MA, Wall A, McGuire AT, Veerler D. Structure, function, and antigenicity of the SARS-CoV-2 spike glycoprotein. *Cell* 2020;181(2):281-92.
- Du L, He Y, Zhou Y, Liu S, Zheng BJ, Jiang S. The spike protein of SARS-CoV—a target for vaccine and therapeutic development. *Nat Rev Microbiol* 2009;7(3): 226-36.
- Du L, Yang Y, Zhou Y, Lu L, Li F, Jiang S. MERS-CoV spike protein: a key target for antivirals. *Expert Opin Ther Targets* 2017;21(2):131-43.
- Datta PK, Liu F, Fischer T, Rappaport J, Qin X. SARS-CoV-2 pandemic and research gaps: Understanding SARS-CoV-2 interaction with the ACE2 receptor and implications for therapy. *Theranostics* 2020;10(16):7448-

- 64.
14. Hoffmann M, Kleine-Weber H, Schroeder S, Krüger N, Herrler T, Erichsen S, et al. SARS-CoV-2 cell entry depends on ACE2 and TMPRSS2 and is blocked by a clinically proven protease inhibitor. *Cell* 2020;181(2): 271-80.
  15. Ou X, Liu Y, Lei X, Li P, Mi D, Ren L, et al. Characterization of spike glycoprotein of SARS-CoV-2 on virus entry and its immune cross-reactivity with SARS-CoV. *Nat Commun* 2020;11(1):1620.
  16. Chen Y, Liu Q, Guo D. Emerging coronaviruses: genome structure, replication, and pathogenesis. *J Med Virol* 2020;92(4):418-23.
  17. Fung TS, Liu DX. Coronavirus infection, ER stress, apoptosis and innate immunity. *Front Microbiol* 2014;5: 296.
  18. Peng H, Yang L, Wang L, Li J, Huang J, Lu Z, et al. Long-lived memory T lymphocyte responses against SARS coronavirus nucleocapsid protein in SARS-recovered patients. *Virology* 2006;351(2):466-75.
  19. Ahmed SF, Quadeer AA, McKay MR. Preliminary identification of potential vaccine targets for the COVID-19 coronavirus (SARS-CoV-2) based on SARS-CoV immunological studies. *Viruses* 2020;12(3):254.
  20. Anand K, Ziebuhr J, Wadhwani P, Mesters JR, Hilgenfeld R. Coronavirus main proteinase (3CLpro) structure: basis for design of anti-SARS drugs. *Science* 2003;300 (5626):1763-7.
  21. Needle D, Lountos GT, Waugh DS. Structures of the Middle East respiratory syndrome coronavirus 3C-like protease reveal insights into substrate specificity. *Acta Crystallogr Sect D Biol Crystallogr* 2015;71(5):1102-11.
  22. ul Qamar MT, Alqahtani SM, Alamri MA, Chen LL. Structural basis of SARS-CoV-2 3CLpro and anti-COVID-19 drug discovery from medicinal plants. *J Pharm Anal* 2020;10(4):313-9.
  23. Smertina E, Urakova N, Strive T, Frese M. Calicivirus RNA-dependent RNA polymerases: evolution, structure, protein dynamics and function. *Front Microbiol* 2019; 10:1280.
  24. Liu W, Morse JS, Lalonde T, Xu S. Learning from the past: possible urgent prevention and treatment options for severe acute respiratory infections caused by 2019-nCoV. *Chembiochem* 2020; 21(5):730-8.
  25. Lung J, Lin Y, Yang Y, Chou Y, Shu L, Cheng Y, et al. The potential chemical structure of anti-SARS-CoV-2 RNA-dependent RNA polymerase. *J Med Virol* 2020;92 (6):693-7.
  26. Jia H, Gong P. A structure-function diversity survey of the RNA-dependent RNA polymerases from the positive-strand RNA viruses. *Front Microbiol* 2019;10:1945.
  27. Romano M, Ruggiero A, Squeglia F, Maga G, Berisio R. A Structural View of SARS-CoV-2 RNA Replication Machinery: RNA Synthesis, Proofreading and Final Capping. *Cells* 2020;9(5):1267.
  28. Touret F, de Lamballerie X. Of chloroquine and COVID-19. *Antiviral Res* 2020;104762.
  29. Chandel V, Sharma PP, Raj S, Choudhari R, Rathi B, Kumar D. Structure-based drug repurposing for targeting Nsp9 replicase and spike proteins of severe acute respiratory syndrome coronavirus 2. *J Biomol Struct Dyn* 2020;1-14.
  30. Alexpandi R, De Mesquita JF, Pandian SK, Ravi AV. Quinolines-based SARS-CoV-2 3CLpro and RdRp inhibitors and Spike-RBD-ACE2 inhibitor for drug-repurposing against COVID-19: an in silico analysis. *Front Microbiol* 2020;11:1796.
  31. Mosquera-Yuqui F, Lopez-Guerra N, Moncayo-Palacio EA. Targeting the 3CLpro and RdRp of SARS-CoV-2 with phytochemicals from medicinal plants of the Andean Region: molecular docking and molecular dynamics simulations. *J Biomol Struct Dyn* 2020;1-14.
  32. Lalani S, Poh CL. Flavonoids as antiviral agents for Enterovirus A71 (EV-A71). *Viruses* 2020;12(2):184.
  33. Jo S, Kim S, Shin DH, Kim MS. Inhibition of SARS-CoV 3CL protease by flavonoids. *J Enzyme Inhib Med Chem* 2020;35(1):145-51.
  34. Hulo C, De Castro E, Masson P, Bougueleret L, Bairoch A, Xenarios I, et al. ViralZone: a knowledge resource to understand virus diversity. *Nucleic Acids Res* 2011;39 (Database issue):D576-82.
  35. Pruitt KD, Tatusova T, Maglott DR. NCBI Reference Sequence (RefSeq): a curated non-redundant sequence database of genomes, transcripts and proteins. *Nucleic Acids Res* 2005;33(Database issue):D501-4.
  36. Haasnoot J, Berkhout B. RNAi and cellular miRNAs in infections by mammalian viruses. In: *Antiviral RNAi*. Springer; 2011. p. 23-41.
  37. Peele KA, Chandrasai P, Srihansa T, Krupanidhi S, Sai AV, Babu DJ, et al. Molecular docking and dynamic simulations for antiviral compounds against SARS-CoV-2: A computational study. *Inform Med Unlocked* 2020; 100345.
  38. Matsuyama S, Nagata N, Shirato K, Kawase M, Takeda M, Taguchi F. Efficient activation of the severe acute respiratory syndrome coronavirus spike protein by the transmembrane protease TMPRSS2. *J Virol* 2010;84(24): 12658-64.
  39. Shulla A, Heald-Sargent T, Subramanya G, Zhao J, Perlman S, Gallagher T. A transmembrane serine protease is linked to the severe acute respiratory syndrome coronavirus receptor and activates virus entry. *J Virol* 2011; 85(2):873-82.
  40. Filimonov DA, Lagunin AA, Glorizova TA, Rudik A V, Druzhilovskii DS, Pogodin P V, et al. Prediction of the biological activity spectra of organic compounds using the PASS online web resource. *Chem Heterocycl Compd* 2014;50(3):444-57.
  41. Maunz A, Gütlein M, Rautenberg M, Vorgrimmler D, Gebele D, Helma C. Lazar: a modular predictive toxicology framework. *Front Pharmacol* 2013;4:38.
  42. Folegatti PM, Bittaye M, Flaxman A, Lopez FR, Bellamy D, Kupke A, et al. Safety and immunogenicity of a candidate Middle East respiratory syndrome coronavirus viral-vectored vaccine: a dose-escalation, open-label, non-randomised, uncontrolled, phase 1 trial. *Lancet In-*



- fect Dis 2020;20(7):816-26.
43. Zhu FC, Guan XH, Li YH, Huang JY, Jiang T, Hou LH, et al. Immunogenicity and safety of a recombinant adenovirus type-5-vectored COVID-19 vaccine in healthy adults aged 18 years or older: a randomised, double-blind, placebo-controlled, phase 2 trial. *Lancet* 2020;396(10249):479-88.
  44. Zheng J. SARS-CoV-2: an emerging coronavirus that causes a global threat. *Int J Biol Sci* 2020;16(10):1678-85.
  45. Zheng L, Zhang L, Huang J, Nandakumar KS, Liu S, Cheng K. Potential treatment methods targeting 2019-nCoV infection. *Eur J Med Chem* 2020;112687.
  46. Li Y, Zhou W, Yang L, You R. Physiological and pathological regulation of ACE2, the SARS-CoV-2 receptor. *Pharmacol Res* 2020;157:104833.
  47. Bacha U, Barrila J, Velazquez-Campoy A, Leavitt SA, Freire E. Identification of novel inhibitors of the SARS coronavirus main protease 3CLpro. *Biochemistry* 2004;43(17):4906-12.
  48. Elfiky AA. SARS-CoV-2 RNA dependent RNA polymerase (RdRp) targeting: An in silico perspective. *J Biomol Struct Dyn* 2020;1-9.
  49. Balmeh N, Mahmoudi S, Mohammadi N, Karabedian-hajabadi A. Predicted therapeutic targets for COVID-19 disease by inhibiting SARS-CoV-2 and its related receptors. *Inform Med Unlocked* 2020;100407.
  50. Xu Z, Peng C, Shi Y, Zhu Z, Mu K, Wang X, et al. Nelfinavir was predicted to be a potential inhibitor of 2019-nCov main protease by an integrative approach combining homology modelling, molecular docking and binding free energy calculation. *BioRxiv*. 2020;
  51. Li Y, Zhang J, Wang N, Li H, Shi Y, Guo G, et al. Therapeutic drugs targeting 2019-nCoV main protease by high-throughput screening. *BioRxiv*. 2020;
  52. Khan RJ, Jha RK, Amera GM, Jain M, Singh E, Pathak A, et al. Targeting SARS-CoV-2: a systematic drug repurposing approach to identify promising inhibitors against 3C-like proteinase and 2'-O-ribose methyl-transferase. *J Biomol Struct Dyn* 2020;1-14.
  53. Khan SA, Zia K, Ashraf S, Uddin R, Ul-Haq Z. Identification of chymotrypsin-like protease inhibitors of SARS-CoV-2 via integrated computational approach. *J Biomol Struct Dyn* 2020;2607-16.
  54. Hendaus MA. Remdesivir in the treatment of Coronavirus Disease 2019 (COVID-19): A simplified summary. *J Biomol Struct Dyn* 2020;1-6.
  55. Food and Drug Administration. Remdesivir EUA Letter of Authorization - FDA [Internet]. Vol. 364 KB. 2020. p. 6. Available from: <https://www.fda.gov/media/137564/download>.
  56. Kadioglu O, Saeed M, Johannes Greten H, Efferth T. Identification of novel compounds against three targets of SARS CoV-2 coronavirus by combined virtual screening and supervised machine learning. *Comput Biol Med* 2021;133:104359.