



A New Specific DNA Target Sequence for Identification of *Staphylococcus epidermidis* using Modified Comparative Genomic Analysis

Reza Khoshbakht^{1,2†}, Hosna Zare^{1,2†}, Reza Kamali Kakhki^{1,2}, Alireza Neshani^{2,3},
and Maryam Arfaatabar^{4*}

1. Antimicrobial Resistance Research Center, Mashhad University of Medical Sciences, Mashhad, Iran

2. Student Research Committee, Mashhad University of Medical Sciences, Mashhad, Iran

3. Department of Laboratory Sciences, School of Paramedical Sciences, Mashhad University of Medical Sciences, Mashhad, Iran

4. Department of Medical Laboratory Sciences, Kashan Branch, Islamic Azad University, Kashan, Iran

† The first and the second authors have had equal contribution to this manuscript

Abstract

Background: *Staphylococcus epidermidis* (*S. epidermidis*) is the most frequently isolated pathogen from prostheses infections in the body. Therefore, improving its diagnostic methods, including rapid Nucleic Acid Amplification Tests (NAAT), seems necessary. Since the first step in designing a NAAT is to find a specific sequence and all DNA targets that have been introduced so far are not completely specific, we introduced a new 100% specific DNA target sequence to identify *S. epidermidis* in this study.

Methods: Modified comparative genomic analysis was used to find the best specific target sequence to detect *S. epidermidis*. A PCR method was designed for the evaluation of this target. To determine the detection limit and analytical specificity, pure genomic DNA of 18 bacteria include 12 standard strains (one *S. epidermidis* and 11 non-*S. epidermidis*) and six clinical isolates (five *S. epidermidis* and one non-*S. epidermidis*) were used.

Results: The 400 bp sequence of *S. epidermidis* ATCC 14990 was identified as the most specific sequence (Se400), having a 100% sequence similarity to *S. epidermidis* genomes but not with other bacteria. The detection limit of Se400-PCR was 10 fg, equal to about 4 copies of *S. epidermidis* genomic DNA/ μ l. All pure DNA templates from *S. epidermidis* generated a detectable amplicon by 264 bp length, but the PCR test was negative for the non-*S. epidermidis* group.

Conclusion: The Se400 sequence can be considered as a specific target for detecting *S. epidermidis*, based on our findings.

Avicenna J Med Biotech 2022; 14(3): 216-222

Keywords: Comparative genomic analysis, Detection, Pathogen, Polymerase chain reaction, Se400, *Staphylococcus epidermidis*

* Corresponding author:
Maryam Arfaatabar Ph.D.,
Department of Medical
Laboratory Sciences, Kashan
Branch, Islamic Azad University,
Kashan, Iran
Tel/Fax: +98 9133615486
E-mail:
arfahmara081@gmail.com
Received: 12 Jan 2022
Accepted: 9 Apr 2022

Introduction

One of the most isolated members of the coagulase-negative staphylococci (CoNS) group is *Staphylococcus epidermidis* (*S. epidermidis*). This bacterium colonizes mucous membranes and the skin, accounting for the majority of the bacterial flora in this environment¹⁻³. Genome study of *S. epidermidis* revealed that it is fully equipped with genes supposed to offer resistance from the severe circumstances faced in surrounding environment, allowing it to remain longer in dry conditions in hospitals⁴. *S. epidermidis* is the most commonly implicated pathogen in infections related to any form of an indwelling medical device⁵. This micro-organism

has been detected with a relatively high prevalence from the Central Nervous System (CNS) shunts, joint prostheses, and prosthetic valves^{6,7}. Also, the mentioned bacterium has been repeatedly isolated from different specimens, such as blood, skin, wound, urinary tract, soft tissue infections, endocarditis, bacteremia, and pneumonia^{8,9}. According to scientific documents, CoNS cause half of all cases of Prosthetic Valve Endocarditis (PVE)¹⁰. More than 20% of people with implanted cardiac devices are infected by *S. epidermidis*, which in turn causes pain and purulence at the infection site and sepsis⁶. The sepsis mortality rate resulting

from *S. epidermidis* in infants could be as high as 4.8 and 9.4%¹¹. The mortality rate due to endocarditis caused by CoNS is reported to be about 36%⁶, while it is estimated at 30% for septic shock¹².

Traditionally, *S. epidermidis* diagnosis has been performed according to the biochemical tests and morphological characteristics. Such methods are time-consuming (sometimes up to several days) and do not reliably distinguish *S. epidermidis* from other CoNS. Therefore, finding faster and more reliable methods has always been required. The development of Nucleic Acid Amplification Tests (NAATs) such as Polymerase Chain Reaction (PCR) in recent decades, has greatly increased the speed, sensitivity, and specificity of diagnostic tests¹³⁻¹⁵. One of the critical points for designing a NAAT is a completely specific DNA sequence for the desired micro-organism. The specific sequence should be present in all strains of such micro-organism but not found in any other micro-organism or has very low similarity¹⁶. In recent years, various PCR tests have been designed to detect *S. epidermidis* based on genes such as *serp0107*, *gseA*, *Staphostatin A*, and *sesC*¹⁷⁻²⁰. Our bioinformatics evaluation showed that all genes introduced so far as diagnostic targets, are not 100% specific, and have many similarities with other species of *Staphylococcus*. Therefore, finding a specific target sequence that can be applied to design a completely specific PCR is still needed. Modified comparative genomic analysis or modified genome comparison is among the methods for finding a specific target sequence for organisms and was introduced by our team in 2018^{16,21}. So, this study aimed to introduce a novel target gene that is specific for the *S. epidermidis* complex, as well as to design highly specific and sensitive primers for the rapid detection of *S. epidermidis* using modified comparative genomic analysis.

Materials and Methods

Staphylococcus epidermidis-specific target mining

According to the described method in our previous studies^{16,21}, the genomic sequence of *S. epidermidis* ATCC 14990 was compared with the available genomic sequences on the nucleotide collection database^{22,23}, and the most specific sequence was selected (Figure 1). The steps are described below:

1. Genomic sequences of *S. epidermidis* on nucleotide collection database were determined. Then, one case which was preferably the NCBI reference sequence, was regarded as the reference.
2. The sequence of *S. epidermidis* ATCC 14990 (NZ_CP035288.1) was selected as the reference, and the sequence was obtained and cut to 5000 bp independent fragments, producing about 493 fragments.
3. Separately, each fragment was compared with other available sequences of nucleotide collection database by Basic Local Alignment Search Tool (BLAST). BLAST discovers similar regions between DNA sequences. The nucleotide sequences is compared with




Figure 1. Method for mining *S. epidermidis*-specific nucleotide sequences.

available sequences on database and the statistical significance is calculated by the program²².

4. After each analysis, results were screened, and the best fragments were selected. Evaluation of results was performed by two criteria:

- a. Presence of all *S. epidermidis* NCBI reference sequences in search results, having both identity and query cover of 100%.
 - b. No other microorganism except *S. epidermidis* would appear with the query cover >90%. The selection of these two criteria was based on our experiences and evaluation of the first 200 fragments.
5. Selected fragments of the previous step were compared separately with non- *S. epidermidis* sequences of the nucleotide collection database, and conserved parts of each fragment were determined.
6. Finally, the longer specific part was selected, and we named it Se400.

Primer design and PCR

To evaluate the specificity of Se400, an end-point PCR was designed with Oligo7 software²⁴. The primers were then tested for secondary structure and anticipated melting temperature using Oligo Analyzer 3.1 (<https://eu.idtdna.com/calc/analyzer>) and were manufactured by DENAzist Asia Company. 264 bp amplicon and Primer sequences are provided in table 1. PCR reaction was prepared in 25 μ l containing 2.5 μ l of 10 \times PCR buffer (100 mM Tris-HCl [pH=8.3], 500 mM KCl), 1 μ l of each 10 μ M forward and reverse primers, 1 μ l of DNA sample, 0.5 μ l of 200 μ M (each) of the four dNTPs, 1.5 μ l of 25 mM MgCl₂, 0.625 U of Taq DNA polymerase, and PCR grade water. 10 ng of

A New Specific Target Gene for Identification of *Staphylococcus epidermidis*

Table 1. The Se400 sequence, the designed primers for PCR, and amplicon

Specific DNA target sequence (Se400)
>NZ_CP035288.1:242200-242600 <i>Staphylococcus epidermidis</i> strain ATCC 14990 chromosome, complete genome GAGAGCCTACACCAAAATTGTAATAAACACGAGGACAGTGATTCTAACATCTTGCCTCGTTTTGAAGACCAAAAATCC ATCATTATTCATCTCACATCTTTAAATTTTTGTTGATAAAAAAAGCAACTTAATCATTATAACGCTAATGGTAAGGTGGG TAAATAAACTGATATGATGATATTAATAAATGAAAAAGAAAATCTTAATATATTTTTATTTTTAAAAAGATAATGTAATTATT TATACTGTAAAGATATGAAAATGGCATGGTATTTCATTAATAAATAACTAACTTTGGAATCAATGACCAATAACC ATAAATACGTTATCCTTACATAGAAGGTTATTAATGAATTCGAATGGAGGATAATACGTTATGATGGATTTTTTACAT
Forward primer
5'-TTGTAATAAACACGAGGACA
Reverse primer
5'-TAATGAAATACCATGCCAT
264 bp amplicon
5'TTGTAATAAACACGAGGACAGTGATTCTAACATCTTGCCTCGTTTTGAAGACCAAAAATCCATCATTTATTCATCTCAC TCTTTAAATTTTTGTTGATAAAAAAAGCAACTTAATCATTATAACGCTAATGGTAAGGTGGGTAATAAACTGATATGATG ATATTAATAAATGAAAAAGAAAATCTTAATATATTTTTATTTTTAAAAAGATAATGTAATTATTTACTGTAAAGATATGA AAATGGCATGGTATTTTCATTA

S. epidermidis ATCC 14990 pure DNA was used as the positive control, and water as the negative control.

For DNA amplification, 5 min of initial denaturation at 95°C was followed by 30 cycles of (i) 45 s of denaturation at 95°C, (ii) 45 s of annealing at 49°C for, (iii) 60 s of extension at 72°C for, and (iv) final 5 min of extension at 72°C. Finally, 3 µl of PCR product was visualized using 1.5% agarose gel electrophoresis and DNA green viewer. The presence of a 264 bp amplicon specifies the positive result.

Bacterial isolates

In this study, pure genomic DNA of 18 bacteria, including 12 standard strains (one *S. epidermidis* and 11 non-*S. epidermidis*) and six clinical isolates (5 *S. epidermidis* and one non-*S. epidermidis*) were used. All standard strains were acquired from the microbial bank in the Antimicrobial Resistance Research Center of Mashhad University of Medical Sciences, and all clinical isolates were acquired from Imam Reza Hospital of Mashhad (Table 2).

Analytical sensitivity (limit of detection)

The pure DNA concentration of *S. epidermidis* ATCC 14990 was calculated by a spectrophotometer (Thermo Scientific). A serial dilution of pure DNA was then prepared in water (10 ng, 1 ng, 100 pg, 10 pg, 1 pg, 100 fg, 10 fg, 1 fg) and the volume of 1 µl was applied as the template. The process was repeated three times to ensure the results.

Analytical specificity

The primer specificity was investigated by Blastn software to determine cross-reactivity with other human or bacterial genomes. To determine the analytical specificity for Se400-PCR, pure DNA of six *S. epidermidis* and 12 non-*S. epidermidis* were used (Table 2). 10 ng of pure DNA was applied in each reaction. Finally, sequencing was performed on all positive PCR products.

Table 2. The list of bacteria used in this research

Groups	Bacterial names
<i>Staphylococcus epidermidis</i>	
	<i>S. epidermidis</i> ATCC 14990
	<i>S. epidermidis</i> (clinical isolate 1)
	<i>S. epidermidis</i> (clinical isolate 2)
	<i>S. epidermidis</i> (clinical isolate 3)
	<i>S. epidermidis</i> (clinical isolate 4)
	<i>S. epidermidis</i> (clinical isolate 5)
Non-<i>Staphylococcus</i>	
	<i>Staphylococcus aureus</i> ATCC 25923
Epidermidis bacteria	
	<i>Staphylococcus haemolyticus</i> ATCC 29970
	<i>Staphylococcus saprophyticus</i> ATCC 15305
	<i>Staphylococcus schleiferi</i> ATCC 43808
	<i>Streptococcus pyogenes</i> ATCC 12344
	<i>Streptococcus agalactiae</i> ATCC 12386
	<i>Enterococcus faecalis</i> ATCC 29212
	<i>Streptococcus oralis</i> (clinical isolate)
	<i>Listeria monocytogenes</i> ATCC 35152
	<i>Lactobacillus casei</i> ATCC 393
	<i>Pseudomonas aeruginosa</i> ATCC 27853
	<i>Escherichia coli</i> ATCC 25922

Results

Target mining

The 400 bp sequence containing nucleotides 242, 200 to 242,600 of *S. epidermidis* ATCC 14990 (NZ_CP035288.1) was recognized as the most specific sequence, having a 100% sequence similarity to *S. epidermidis* genomes but not with other bacteria. Se400 is a non-coding sequence located between *EQW00_01195* and *EQW00_01200* genes (Figure 2). The blastn search showed that Se400 could detect all strains of *S. epidermidis* among the registered complete genomes in the nucleotide collection database, and no similarity was observed with other micro-organisms.




Figure 2. Schematic location of Se400 between *EQW00_01195* and *EQW00_01200* genes.

Analytical sensitivity (limit of detection)

The detection limit is defined as the lowest analyte concentration that can be reliably detected. Consequently, the Se400-PCR detection limit was 10 fg, equal to about 4 copies of *S. epidermidis* genomic DNA/μl (Figure 3).

The analytical specificity

PCR amplification using Se400-specific primers was performed with 18 bacteria using pure genomic DNA as the template to consider the analytical specificity of the Se400-PCR. As presented in figure 4, all genomic DNA templates from *S. epidermidis* generated a detectable amplicon by 264 bp length, but the PCR test was negative for the non-*S. epidermidis* group; subsequently, the Se400 primer was specific for detecting *S. epidermidis*. Finally, analysis of sequencing results for PCR products from 6 positive samples showed that the produced amplicon is related to the Se400 sequence of *S. epidermidis*.




Figure 3. Gel electrophoresis of Se400-PCR product on 1.5% agarose gel. 264 bp amplicons are shown at various concentrations of *S. epidermidis* pure DNA as template. Amplicons using 10 ng, 1 ng, 100 pg, 10 pg, 1 pg, 100 fg, 10 fg, and 1 fg are shown in lanes.




Figure 4. Gel electrophoresis of Se400-PCR products. The amplicon size is 264 bp. M: 100-bp molecular DNA ladder (Pars Tous, Iran).

Discussion

Before the discovery of molecular methods, phenotypic and biochemical tests were the only powerful methods for the detection and differentiation of various bacteria, including *S. epidermidis*. With the development of molecular methods, especially PCR in recent decades, the problems of traditional methods such as being slow and lacking reliability were solved^{13,14}. One of the challenges for designing a PCR test to detect *S. epidermidis* is the lack of a completely specific DNA target. According to the literature, most NAATs in the 1990s used genus-specific targets due to the unavailability of species-specific DNA targets. The amplified sequences based on such genes could only differentiate *S. epidermidis* from other close species when they were analyzed by confirmatory methods. The most famous method for analyzing these fragments was sequencing and then comparing the results with available sequences on GenBank. *16S rDNA*, *sodA*, *hsp60*, and *tuf* were among the most used genes in such methods²⁵⁻²⁸. Although the addition of the sequencing step reduced the speed and increased the costs, some cases were still seen in which the sequenced fragment was quite similar in several species^{25,29}. For example, although the *16S rDNA* gene has been suggested as a target gene, it cannot be used as a distinct target to detect *S. epidermidis* in clinical specimens due to the significant similarity to *Staphylococcus aureus* (*S. aureus*)³⁰⁻³².

With an increased detection of *S. epidermidis* infections in the last two decades, the need to find species-specific targets for the detection of this bacterium has been increased significantly. Efforts eventually led to the introduction of *serp0107*, *gseA*, *ecpB*, and *SesC* genes as species-specific targets¹⁷⁻²⁰. However, our bioinformatics evaluation before starting this project showed that none of these diagnostic targets was 100% specific (Table 3).

In 2004, the *gseA* gene (GenBank acc. No. AB096695), responsible for the production of glutamic acid-specific 27-kDa serine protease (GluSE), was introduced by Ikeda *et al* as a species-specific target of *S. epidermidis*. This protease is involved in degrading human fibronectin, collagen, the complement protein C5, and slime formation. Thus, the protease may be associated with the pathogenesis of *S. epidermidis*^{17,33}. Despite an appropriate length of this gene (1214 bp), a comparison of its sequence with other available sequences at nucleotide collection database showed that this gene has similarity with some parts of the *Staphylococcus saccharolyticus* (*S. saccharolyticus*) genome with >68% query cover and >75% identity. This bacterium is a normal flora of the skin, and the pathogenesis is not apparent yet. Furthermore, very similar sequences to the *gseA* gene are present in other bacteria, reducing the specificity of this gene for the detection *S. epidermidis*. These bacteria include *Staphylococcus capitis* (*S. capitis*), *Staphylococcus caprae* (*S. caprae*), *S.*

A New Specific Target Gene for Identification of *Staphylococcus epidermidis*

Table 3. Comparison of Se400 sequence specificity with other introduced genes for detecting *S. epidermidis* in species level

Target	Length	Similarity to non- <i>S. epidermidis</i> genomes with >35% query cover and >60% identity
Se400	400 bp	No similarity to any non- <i>S. epidermidis</i> microbes
gseA (2004)	1214 bp	<ul style="list-style-type: none"> • 3 strains of <i>S. saccharolyticus</i> with >68% query cover and >75% identity • 5 strains of <i>S. caprae</i> with >58% query cover and >69% identity • 11 strains of <i>S. warneri</i> with 55% query cover and 67% identity • >1300 strain of <i>S. aureus</i> with 50% query cover and 67-69% identity • 6 strains of <i>S. pasteur</i> with 46-51% query cover and 67-69% identity
serp0107 (2006)	882 bp	<ul style="list-style-type: none"> • 3 strains of <i>S. saccharolyticus</i> with 98% query cover and 69% identity • 7 strains of <i>S. capitis</i> with 95% query cover and 68% identity • 5 strains of <i>S. caprae</i> with 66% query cover and 70% identity • 3 strains of <i>S. haemolyticus</i> with 42% query cover and >68% identity • Other microorganisms: <i>S. warneri</i>, <i>S. hominis</i>, <i>S. argenteus</i>
sesC (2016)	2031 bp	<ul style="list-style-type: none"> • 3 strains of <i>S. saccharolyticus</i> with 83-100% query cover and >69% identity • 7 strains of <i>S. capitis</i> with 42% query cover and >67% identity • 5 strains of <i>S. caprae</i> with 52% query cover and 66% identity
ecpB (2019)	318 bp	<ul style="list-style-type: none"> • 3 strains of <i>S. saccharolyticus</i> with 100% query cover and >77% identity • 19 strain of <i>S. aureus</i> with 65% query cover and 67% identity

The specificity of the genes was tested bioinformatically using Blastn software.

aureus, *Staphylococcus haemolyticus* (*S. haemolyticus*), *Staphylococcus saprophyticus* (*S. saprophyticus*), *Staphylococcus hominis* (*S. hominis*), *Staphylococcus equorum* (*S. equorum*), *Staphylococcus simulans* (*S. simulans*), *Staphylococcus lugdunensis* (*S. lugdunensis*), and *Staphylococcus schleiferi* (*S. schleiferi*).

serp0107, which is a putative transcriptional regulator gene, was introduced by Liu *et al* in 2006 as the species-specific diagnostic target to detect *S. epidermidis*. The sequence of this gene with 882 bp length was obtained from the (nt 91276–92157) of *S. epidermidis* RP62a (GenBank Accession No. CP000029) ¹⁸. Our assessment of this sequence using the BLASTN search tool showed that a highly similar sequence is also present in *S. saccharolyticus* and *S. capitis*. Also, similar points to the *serp0107* gene are found in other *Staphylococcus* species, including *S. caprae*, *S. haemolyticus*, *Staphylococcus warneri* (*S. warneri*), *S. hominis*, and *Staphylococcus argenteus* (*S. argenteus*) leads to a lack of complete specificity of this gene for the detection of *S. epidermidis*.

S. epidermidis surface protein C (*SesC*) gene was introduced by Khodaparast *et al* in 2016 as a specific diagnostic target of *S. epidermidis*. *SesC* protein is expressed more in *S. epidermidis* biofilm-associated cells than planktonic ones. Also, this target is appropriate to design various NAATs due to the suitable length (2031 bp) ¹⁹. Nevertheless, our evaluation of this gene showed that similar sequences are found in three other species of the *Staphylococcus* genus, which reduces its specificity. The highest similarity was seen for the *S. saccharolyticus* with 83-100% query cover and >69% identity, followed by *S. capitis* and *S. caprae*.

Finally, the last introduced species-specific gene for this bacterium was the *ecpB* gene, encoding the Staphostatin A protein. This 318 bp gene was initially

introduced to differentiate *S. aureus* and *S. epidermidis* ²⁰. However, the high similarity with a sequence in *S. aureus* with 65% query cover and 67% identity makes it unspecific. The assessment of available sequences in NCBI also showed that this gene is also found in *S. saccharolyticus* with 100% query cover and >77% identity.

According to the results obtained using the BLASTN search tool, the Se400 sequence is completely specific to the *S. epidermidis* and is not found in any other micro-organisms, unlike the other sequences having nonspecific regions in several points. Therefore, using this sequence seems to solve the unavailability of a completely specific target for *S. epidermidis*.

In this study, the specificity of the Se400 target sequence was confirmed by PCR. The specificity of Se400 in all *S. epidermidis* strains highlighted both the efficacy of the comparative genomic analysis for finding possible targets and the significance of experimental research. Also, the Se400-PCR test was able to detect very low levels of genomic DNA template. One of our limitations was the small number of bacterial strains tested, and it is recommended that further research be conducted with a large sample size. Future studies might include comparing the previously known target gene with Se400 sequence in clinical samples.

Conclusion

In conclusion, *S. epidermidis*-specific target sequences were identified using a new comparative genomics method for finding species-specific nucleotide sequences. Many nucleotide targets were assessed, and a target sequence was applied to design a PCR test to detect *S. epidermidis* in clinical samples. Further surveys are being planned to include more bacterial strains for the evaluation of the particular targets. Unique tar-

gets may be found using this method for the detection of any micro-organism, for which a genome sequence is available.

Conflict of Interest

None declared.

References

- Morse SA, Mietzner TA, Miller S, Riedel S. *Jawetz Melnick & Adelbergs Medical Microbiology* 28 E: McGraw-Hill Education; 2019.
- Stacy A, Belkaid Y. *Microbial guardians of skin health. Science* 2019;363(6424):227-8.
- Rendboe AK, Johannesen TB, Ingham AC, Månsson E, Iversen S, Baig S, et al. The Epidome - a species-specific approach to assess the population structure and heterogeneity of *Staphylococcus epidermidis* colonization and infection. *BMC Microbiol* 2020;20(1):362.
- Chu V, Miro JM, Hoen B, Cabell CH, Pappas PA, Jones P, et al. Coagulase-negative staphylococcal prosthetic valve endocarditis—a contemporary update based on the International Collaboration on Endocarditis: prospective cohort study. *Heart* 2009;95(7):570-6.
- Sabaté Brescó M, Harris LG, Thompson K, Stanic B, Morgenstern M, O'Mahony L, et al. Pathogenic mechanisms and host interactions in *Staphylococcus epidermidis* device-related infection. *Front Microbiol* 2017;8(1401).
- Rogers KL, Fey PD, Rupp ME. Coagulase-negative staphylococcal infections. *Infect Dis Clin North Am* 2009;23(1):73-98.
- McCann MT, Gilmore BF, Gorman SP. *Staphylococcus epidermidis* device-related infections: pathogenesis and clinical management. *J Pharm Pharmacol* 2008;60(12):1551-71.
- Bennett JE, Dolin R, Blaser MJ. *Mandell, Douglas, and Bennett's Principles and Practice of Infectious Diseases E-Book*: Elsevier Health Sciences; 2019.
- Namvar AE, Bastarahang S, Abbasi N, Ghehi GS, Farhadbakhtrian S, Arezi P, et al. Clinical characteristics of *Staphylococcus epidermidis*: a systematic review. *GMS Hyg Infect Control* 2014;9(3):Doc23.
- Lalani T, Kanafani ZA, Chu VH, Moore L, Corey GR, Pappas P, et al. Prosthetic valve endocarditis due to coagulase-negative staphylococci: findings from the International Collaboration on Endocarditis Merged Database. *Eur J Clin Microbiol Infect Dis* 2006;25(6):365-8.
- Dong Y, Speer CP, Glaser K. Beyond sepsis: *Staphylococcus epidermidis* is an underestimated but significant contributor to neonatal morbidity. *Virulence* 2018;9(1):621-33.
- Kumar G, Kumar N, Taneja A, Kaleekal T, Tarima S, McGinley E, et al. Nationwide trends of severe sepsis in the 21st century (2000-2007). *Chest* 2011;140(5):1223-31.
- Sah S, Bordoloi P, Vijaya D, Amarnath SK, Devi CS, Indumathi V, et al. Simple and economical method for identification and speciation of *Staphylococcus epidermidis* and other coagulase negative *Staphylococci* and its validation by molecular methods. *J Microbiol Methods* 2018;149:106-19.
- Roberts AL. Identification of *Staphylococcus epidermidis* in the clinical microbiology laboratory by molecular methods. *Methods Mol Biol* 2014;1106:33-53.
- Safdari H, Neshani A, Sadeghian A, Ebrahimi M, Iranshahi M, Sadeghian H. Potent and selective inhibitors of class A β -lactamase: 7-prenyloxy coumarins. *J Antibiot (Tokyo)* 2014;67(5):373-7.
- Neshani A, Kakhki RK, Sankian M, Zare H, Chichaklu AH, Sayyadi M, et al. Modified genome comparison method: a new approach for identification of specific targets in molecular diagnostic tests using *Mycobacterium tuberculosis* complex as an example. *BMC Infect Dis* 2018;18(1):517.
- Ikeda Y, Ohara-Nemoto Y, Kimura S, Ishibashi K, Kikuchi K. PCR-based identification of *Staphylococcus epidermidis* targeting *gseA* encoding the glutamic-acid-specific protease. *Can J Microbiol* 2004;50(7):493-8.
- Liu D, Swiatlo E, Austin F, Lawrence M. Use of a putative transcriptional regulator gene as target for specific identification of *Staphylococcus epidermidis*. *Lett Appl Microbiol* 2006;43(3):325-30.
- Khodaparast L, Khodaparast L, Van Mellaert L, Shahrooei M, Van Ranst M, Van Eldere J. *sesC* as a genetic marker for easy identification of *Staphylococcus epidermidis* from other isolates. *Infect Genet Evol* 2016;43:222-4.
- Ghatts MZ, ElRakaiby MT, Aziz RK, Zedan HH. A novel PCR method targeting staphostatin genes differentiates *Staphylococcus aureus* from *Staphylococcus epidermidis* in clinical isolates and nasal microbiome samples. *Research square* 2019. Unpublished.
- Kakhki RK, Neshani A, Sankian M, Ghazvini K, Hooshyar A, Sayadi M. The short-chain dehydrogenases/reductases (SDR) gene: a new specific target for rapid detection of *Mycobacterium tuberculosis* complex by modified comparative genomic analysis. *Infect Genet Evol* 2019;70:158-64.
- Sayers EW, Beck J, Bolton EE, Bourexis D, Brister JR, Canese K, et al. Database resources of the national center for biotechnology information. *Nucleic Acids Res* 2021;49(D1):D10.
- Sharma S, Ciuffo S, Starchenko E, Darji D, Chlumsky L, Karsch-Mizrachi I, et al. The NCBI biocollections database. *Database (Oxford)* 2018;2018:bay006.
- Rychlik W. OLIGO 7 primer analysis software. *Methods Mol Biol* 2007;402:35-60.
- Stackebrandt E, GOEBEL BM. Taxonomic note: a place for DNA-DNA reassociation and 16S rRNA sequence analysis in the present species definition in bacteriology. *Int J Systematic Evolutionary Microbiology* 1994;44(4):846-9.
- Goh SH, Potter S, Wood JO, Hemmingsen SM, Reynolds RP, Chow AW. HSP60 gene sequences as universal targets for microbial species identification: studies with coagulase-negative staphylococci. *J Clin Microbiol* 1996;34(4):818-23.

27. Martineau F, Picard FJ, Ke D, Paradis S, Roy PH, Ouellette M, et al. Development of a PCR assay for identification of staphylococci at genus and species levels. *J Clin Microbiol* 2001;39(7):2541-7.
28. Poyart C, Quesne G, Boumaila C, Trieu-Cuot P. Rapid and accurate species-level identification of coagulase-negative staphylococci by using the *sodA* gene as a target. *J Clin Microbiol* 2001;39(12):4296-301.
29. Sivadon V, Rottman M, Quincampoix JC, Avettand V, Chaverot S, de Mazancourt P, et al. Use of *sodA* sequencing for the identification of clinical isolates of coagulase-negative staphylococci. *Clin Microbiol Infect* 2004;10(10):939-42.
30. Zakrzewska-Czerwińska J, Gaszewska-Mastalarz A, Pulverer G, Mordarski M. Identification of *Staphylococcus epidermidis* using a 16S rRNA-directed oligo-nucleotide probe. *FEMS Microbiol Lett* 1992;100(1-3):51-8.
31. Bahador A, Esmaili D, Khaledi A, Ghorbanzadeh R. An in vitro assessment of the antibacterial properties of nanosilver Iranian MTA against *Porphyromonas gingivalis*. *J Chem Pharmaceut Res* 2013;5(10):65-71.
32. Khaledi A, Khademi F, Esmaili D, Esmaili SA, Rostami H. The role of HPaA protein as candidate vaccine against *Helicobacter pylori*. *Der Pharma Chemica* 2016; 8(3):235-7.
33. Hosseini SMJ, Naeini NS, Khaledi A, Daymad SF, Esmaili D. Evaluate the relationship between class 1 integrons and drug resistance genes in clinical isolates of *Pseudomonas aeruginosa*. *Open Microbiol J* 2016;10: 188-96.