

Social support and anxiety correlation with resilience in patients with multiple sclerosis during COVID-19 pandemic

Received: 13 June 2023
Accepted: 09 Aug. 2023

Vahid Shaygannejad^{1,2}, Omid Mirmosayyeb^{1,2}, Aysa Shaygannejad², Saeed Vaheb², Sara Bagherieh², Mozhdeh Askari², Mahsa Ghajarzadeh³

¹ Department of Neurology, School of Medicine, Isfahan University of Medical Sciences, Isfahan, Iran

² Isfahan Neurosciences Research Center, Isfahan University of Medical Sciences, Isfahan, Iran

³ Universal Council of Epidemiology (UCE), Universal Scientific Education and Research Network (USERN), University of Tehran, Tehran, Iran

Keywords

Multiple Sclerosis; COVID-19; Resilience; Anxiety; Social Support

Abstract

Background: Social support and anxiety are essential for patients with chronic diseases such as multiple sclerosis (MS). During coronavirus disease 2019 (COVID-19) pandemic, the psychological well-being of subjects with MS was an important issue, and we designed this study to assess anxiety, resilience, and social support in these patients during COVID-19 pandemic stage.

Methods: We used convenience sampling for this study. Inclusion criteria were definite diagnosis of MS based on McDonald criteria and age more than 18 years. Two hundred patients with MS were enrolled. Subjects were asked to fill out valid and reliable Persian versions of Connor-Davidson Resilience Scale (CD-RISC), Beck Anxiety Inventory (BAI), and Perceived Social Support Scale. We also collected demographic

data (age, sex, marital status, and occupation), disease duration, and disability level [Expanded Disability Status Scale (EDSS)]. Continuous variables were presented as mean \pm standard deviation (SD) (except for EDSS, as its distribution was not normal), and categorical variables were presented as frequencies. Correlation coefficients were calculated. We did a subgroup analysis and compared patients with BAI less than 30 and more than 30 (severe anxiety). A P-value less than 0.05 was considered significant.

Results: Mean age and mean duration of the disease were 36.5 ± 9.3 and 6.2 ± 5.4 years, respectively. Mean BAI, social support, and resilience scores were 33.8 ± 11.4 , 65.7 ± 16.7 , and 62.5 ± 19.4 , respectively.

How to cite this article: Shaygannejad V, Mirmosayyeb O, Shaygannejad A, Vaheb S, Bagherieh S, Askari M, et al. Social support and anxiety correlation with resilience in patients with multiple sclerosis during COVID-19 pandemic. Curr J Neurol 2023; 22(4): 217-20.

There was a significant positive correlation between resilience and social support scales ($r = 0.44$, $P < 0.001$), and also a significant negative correlation between resilience and BAI ($r = -0.31$, $P < 0.001$). Patients with severe anxiety (BAI > 30) had lower social support scores and resilience (social support: 70.3 ± 13.1 vs. 61.5 ± 18.6 , $P < 0.001$; resilience: 57.3 ± 17.0 vs. 68.2 ± 19.6 , $P < 0.001$) compared to patients with BAI ≤ 30 . By considering resilience score as the dependent variable and other variables as independent variables, we found that BAI and social support scores were independent predictors.

Conclusion: Social support and anxiety are independent predictors of resilience during COVID-19 pandemic in patients with MS.

Introduction

Multiple sclerosis (MS), an autoimmune disease of the central nervous system (CNS), is a disabling condition which could be controlled after the administration of immunomodulatory or immunosuppressive agents.^{1,2} Administration of immunosuppressive agents in these patients was suggested to increase the risk of any infections such as coronavirus disease 2019 (COVID-19), while the literature showed that the infection rate was not higher in the MS population.¹ A recent systematic review and meta-analysis showed that the pooled prevalence of COVID-19 disease in subjects with MS was estimated as 4%.³

Psychological problems such as anxiety are among the prevalent complications in MS.^{4,5} Anxiety in subjects with MS is related to worsening of neurological symptoms and disability.^{5,6} During the pandemic period, subjects with MS may have fears of infection and its consequences such as relapses which interfere with daily activities and quality of life (QOL).⁷ Social support will help them adjust with stressful life events and support themselves from negative outcomes.⁸ Protection against mental disorders and adaptation with life events is defined as resilience.^{9,10}

During the COVID-19 pandemic, the psychological well-being of subjects with MS was an important issue and we designed this study to assess anxiety, resilience, and social support in these patients during the COVID-19 pandemic stage.

Materials and Methods

This cross-sectional study was done in Hakim Private Clinic in Isfahan, Iran, Between March and June 2022. We used convenience sampling for this study.

Inclusion criteria were definite diagnosis of MS based on McDonald criteria¹¹ and age of more than 18 years.

All participants were asked to fill out informed consent forms, although the study had been approved by the Ethics Committee of Isfahan University of Medical Sciences (IR.ARI.MUI.REC.1400.124).

Subjects were also asked to fill valid and reliable Persian version of Connor-Davidson Resilience Scale (CD-RISC), Beck Anxiety Inventory (BAI), and Perceived Social Support Scale.¹²⁻¹⁴

CD-RISC includes 25 questions, evaluating the feelings of the last month. Each question could be answered based on a Likert scale: not true at all (0), rarely true (1), sometimes true (2), often true (3), and true nearly all of the time (4). The sum of the answers is the total score. The higher the score, the greater the resilience of the subject.

BAI includes 21 self-reported questions; each should be answered on a Likert scale (from 0 to 3). The higher the score, the higher the anxiety level.

Perceived Social Support Scale includes 12 questions; each is graded from 1 (very strongly disagree) to 7 (very strongly agree), evaluating different aspects of support. The sum of the scores leads to the total score. The higher the score, the better the perceived social support.

We also collected demographic data (age, sex, marital status, occupation), disease duration, and disability level [Expanded Disability Status Scale (EDSS)].

All data were analyzed using Stata software (version 13.0, Stata Corporation, College Station, TX, USA).

Continuous variables were presented as mean \pm standard deviation (SD) (except for EDSS, as its distribution was not normal), and categorical variables were presented as frequencies. Correlation coefficients were calculated.

We did a subgroup analysis and compared patients with BAI less than 30 and more than 30 (severe anxiety). A P-value less than 0.05 was considered significant.

Results

Two hundred patients were enrolled in this study. One hundred sixty-two were women (81%) and 38 (19%) were men. Mean age and mean duration of the disease were 36.5 ± 9.3 and 6.2 ± 5.4 years, respectively (Table 1).

Mean BAI, social support, and resilience scores

were 33.8 ± 11.4 , 65.7 ± 16.7 , and 62.5 ± 19.4 , respectively.

Table 1. Basic characteristics of enrolled subjects in the study

Variable	Value
Sex [n (%)]	
Women	162 (81.0)
Men	38 (19.0)
Age (year) (mean \pm SD)	36.5 ± 9.3
Marital status [n (%)]	
Single	57 (28.5)
Married	143 (71.5)
Occupation [n (%)]	
Employed	81 (42.0)
Unemployed	118 (59.0)
EDSS [median (IQR)]	0 (2)

SD: Standard deviation; IQR: Interquartile range

There was a significant positive correlation between resilience and social support scales ($r = 0.44$, $P < 0.001$), and also a significant negative correlation between resilience and BAI ($r = -0.31$, $P < 0.001$).

Patients with severe anxiety (BAI > 30) had lower social support score and resilience (social support: 70.3 ± 13.1 vs. 61.5 ± 18.6 , $P < 0.001$; resilience: 57.3 ± 17.0 vs. 68.2 ± 19.6 , $P < 0.001$) compared to patients with BAI ≤ 30 .

By considering resilience score as dependent variable and other variables as independent variables, we found that BAI and social support scores were independent predictors (Table 2).

Table 2. Linear regression analysis by considering resilience score as dependent variable and other variables as independent variables

Variables	Unadjusted regression coefficient	95% CI	P
BAI	-0.37	-0.60, -0.14	0.003
Social support	0.45	0.30-0.60	< 0.001
EDSS	0.49	-0.40, 1.40	0.070
Age	0.16	-0.11, -0.45	0.040
Sex	1.90	-0.46, 8.40	0.200

BAI: Beck Anxiety Inventory; EDSS: Expanded Disability Status Scale; CI: Confidence interval

Discussion

To our knowledge, this is the first study evaluating resilience, social support, and anxiety during the COVID-19 pandemic in patients with MS.

The results show that social support predicts resilience during the COVID-19 in patients with

MS. We found that there was a significant positive correlation between resilience and social support, and social support and anxiety were significant predictors of the resilience.

This finding is not surprising as positive social interactions lead to better daily activities and improved QOL. A recent study showed that better social support was related to better quality of sleep.¹⁵ They found that this relationship was independent of age, education, disability, and psychological status.

Social support is characterized by emotional support from other persons¹⁶ that results in better self-care, improved QOL, and encouraging better behavior.^{17,18} During the pandemic, social support was limited and patients with MS were not excepted.

Resilience in patients with chronic diseases such as MS is dependent on self-efficacy, optimism, hardiness, coping strategies, benefit finding, and social support.¹⁹ Resilience leads to better treatment response, and is negatively related to depression and anxiety.²⁰⁻²⁴

Patients with MS have a higher level of anxiety, and management of stress will help them have better self-care and health. Anxiety could exacerbate the severity of MS disease. In a previous study in Iran, Naser Moghadasi found that patients with MS had moderate to severe anxiety level²⁵ while in a study in Serbia,²⁶ patients with MS showed a higher level of anxiety compared to one year ago (before the pandemic). This indicates the influence of public health on the level of anxiety in patients with MS. They also investigated worse results when compared with healthy subjects.

In another study in Iran, in which patients with MS were enrolled, the prevalence of anxiety during COVID-19 pandemic was reported as 39%.

It should be considered that long isolation due to severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) and a sense of uncertainty related to the pandemic led to a higher level of anxiety in subjects with MS. They avoided public attendance, and tried to do all public health guidelines. Our results show that stress had a significant negative correlation with resilience, and BAI score was a significant negative predictor of resilience score.

Patients with comorbidities have been considered to have a higher risk of infection, and they experience more anxiety and fear regarding this issue.

In this case, social support which could be through different ways such as telephone calls, chats, social media, group meetings, etc. will help

patients pass this period with less complications.

This study had some strengths. First, we evaluated resilience, social support, and anxiety simultaneously. Second, the sample size was large.

The study had also limitations. First, we did not assess the depression and other psychological status in patients. Second, we did not evaluate fear of relapse.

Further multi-centric studies evaluating psychological status and other risk factors of resilience comprehensively are recommended.

References

1. Ghajarzadeh M, Bonavita S. Are patients with multiple sclerosis (MS) at higher risk of COVID-19 infection? *Neurol Sci* 2020; 41(9): 2315-6.
2. Mirmosayyeb O, Shaygannejad V, Bagherieh S, Hosseinabadi AM, Ghajarzadeh M. Prevalence of multiple sclerosis (MS) in Iran: A systematic review and meta-analysis. *Neurol Sci* 2022; 43(1): 233-41.
3. Moghadasi AN, Mirmosayyeb O, Barzegar M, Sahraian MA, Ghajarzadeh M. The prevalence of COVID-19 infection in patients with multiple sclerosis (MS): A systematic review and meta-analysis. *Neurol Sci* 2021; 42(8): 3093-9.
4. Santangelo G, Sacco R, Siciliano M, Bisecco A, Muzzo G, Docimo R, et al. Anxiety in multiple sclerosis: Psychometric properties of the State-Trait Anxiety Inventory. *Acta Neurol Scand* 2016; 134(6): 458-66.
5. Askari F, Ghajarzadeh M, Mohammadifar M, Azimi A, Sahraian MA, Owji M. Anxiety in patients with multiple sclerosis: Association with disability, depression, disease type and sex. *Acta Med Iran* 2014; 52(12): 889-92.
6. Bonavita S, Sparaco M, Russo A, Borriello G, Lavorgna L. Perceived stress and social support in a large population of people with multiple sclerosis recruited online through the COVID-19 pandemic. *Eur J Neurol* 2021; 28(10): 3396-402.
7. Shaygannejad V, Mirmosayyeb O, Nehzat N, Ghajarzadeh M. Fear of relapse, social support, and psychological well-being (depression, anxiety, and stress level) of patients with multiple sclerosis (MS) during the COVID-19 pandemic stage. *Neurol Sci* 2021; 42(7): 2615-8.
8. Dennison L, Moss-Morris R, Chalder T. A review of psychological correlates of adjustment in patients with multiple sclerosis. *Clin Psychol Rev* 2009; 29(2): 141-53.
9. Nishi D, Uehara R, Kondo M, Matsuoka Y. Reliability and validity of the Japanese version of the Resilience Scale and its short version. *BMC Res Notes* 2010; 3: 310.
10. Luthar SS, Cicchetti D, Becker B. The construct of resilience: A critical evaluation and guidelines for future work. *Child Dev* 2000; 71(3): 543-62.
11. Thompson AJ, Banwell BL, Barkhof F, Carroll WM, Coetzee T, Comi G, et al. Diagnosis of multiple sclerosis: 2017 revisions of the McDonald criteria. *Lancet Neurol* 2018; 17(2): 162-73.
12. Connor KM, Davidson JR. Development of a new resilience scale: The Connor-Davidson Resilience Scale (CD-RISC). *Depress Anxiety* 2003; 18(2): 76-82.
13. Bagherian-Sararoudi R, Hajian A, Ehsan HB, Sarafraz MR, Zimet GD. Psychometric properties of the Persian version of the Multidimensional Scale Of Perceived Social Support in Iran. *Int J Prev Med* 2013; 4(11): 1277-81.
14. Kaviani H, Mousavi AS. Psychometric properties of the Persian version of Beck Anxiety Inventory (BAI). *Tehran Univ Med J* 2008; 66(2): 136-40. [In Persian].
15. Harris TP, Zaeske LM, Ludwig R, Baker S, Lynch S, Thuringer A, et al. Social support predicts sleep quality in people with multiple sclerosis during the COVID-19 pandemic. *Mult Scler Relat Disord* 2022; 64: 103970.
16. Taylor SE. Social support: A review. In: Friedman HS, editor. *The Oxford handbook of health psychology*. New York, NY: Oxford University Press; 2011. p. 189-214.
17. Pearson JE. The definition and measurement of social support. *J Couns Dev* 1986; 64(6): 390-5.
18. Graven LJ, Grant JS. Social support and self-care behaviors in individuals with heart failure: An integrative review. *Int J Nurs Stud* 2014; 51(2): 320-33.
19. Stewart DE, Yuen T. A systematic review of resilience in the physically ill. *Psychosomatics* 2011; 52(3): 199-209.
20. Min JA, Lee CU, Chae JH. Resilience moderates the risk of depression and anxiety symptoms on suicidal ideation in patients with depression and/or anxiety disorders. *Compr Psychiatry* 2015; 56: 103-11.
21. Min JA, Lee NB, Lee CU, Lee C, Chae JH. Low trait anxiety, high resilience, and their interaction as possible predictors for treatment response in patients with depression. *J Affect Disord* 2012; 137(1-3): 61-9.
22. Min JA, Yu JJ, Lee CU, Chae JH. Cognitive emotion regulation strategies contributing to resilience in patients with depression and/or anxiety disorders. *Compr Psychiatry* 2013; 54(8): 1190-7.
23. Wagnild GM, Young HM. Development and psychometric evaluation of the Resilience Scale. *J Nurs Meas* 1993; 1(2): 165-78.
24. Herrmann D, Scherg H, Verres R, von HC, Strowitzki T, Wischmann T. Resilience in infertile couples acts as a protective factor against infertility-specific distress and impaired quality of life. *J Assist Reprod Genet* 2011; 28(11): 1111-7.
25. Naser MA. One Aspect of Coronavirus disease (COVID-19) Outbreak in Iran: High anxiety among MS patients. *Mult Scler Relat Disord* 2020; 41: 102138.
26. Stojanov A, Malobabic M, Milosevic V, Stojanov J, Vojinovic S, Stanojevic G, et al. Psychological status of patients with relapsing-remitting multiple sclerosis during coronavirus disease-2019 outbreak. *Mult Scler Relat Disord* 2020; 45: 102407.

Conclusion

Social support and anxiety are independent predictors of resilience during COVID-19 pandemic in patients with MS.

Conflict of Interests

The authors declare no conflict of interest in this study.

Acknowledgments

None.