

# Effect of Stress Management Training on the Mental Health of Medical Students and Interns Training Centers During the Pandemic: An Intervention Study

Sobhan Abbasi<sup>1</sup>, Yahya Mohammadi<sup>2</sup>, Mohammadreza Raeisoon<sup>3, 4\*</sup>

<sup>1</sup>School of Medicine, Birjand University of Medical Sciences, Birjand, Iran.

<sup>2</sup>Education Development Center, Birjand University of Medical Sciences, Birjand, Iran.

<sup>3</sup>Department of Community Medicine, School of Medicine, Birjand University of Medical Sciences, Birjand, Iran.

<sup>4</sup>Cardiovascular Research Center, Birjand University of Medical Sciences, Birjand, Iran.

Corresponding Author: Mohammadreza Raeisoon, Department of Community Medicine, School of Medicine, Birjand University of Medical Sciences, Birjand, Iran. Cardiovascular Research Center, Birjand University of Medical Sciences, Birjand, Iran. Email:raeisoon49@gmail.com; ORCID: 0009-0000-8652-5171

Received 2023 April 28; Accepted 2023 May 22.

## Abstract

**Background:** Medical students, as a part of the first line of combating COVID-19, have been exposed to harm caused by mental disorders. Objectives: This study investigated how stress management training affects the mental health of medical students and interns during the COVID-19 pandemic.

**Methods:** sixteen training and internship students of Birjand University of Medical Sciences from 2020-2021 were included in this intervention study. The intervention group completed the stress management training course in 6 sessions. Mental health dimensions were collected with the SCL90 questionnaire before and after intervention in both groups. SPSS 22 software and Mann-Whitney U-test, Wilcoxon, Chi-square, and Fisher's exact test were utilized for data analysis.

**Results:** The basic data showed no significant differences between groups ( $P > 0.05$ ). After the intervention, the mean scores of physical complaint, obsessive-compulsive, interpersonal relations, depression, anxiety, aggression, fear, paranoid and psychotic dimensions in the intervention group significantly decreased ( $P < 0.05$ ). In controls, the obsessive-compulsive, depression, anxiety, and aggression dimensions significantly decreased ( $P < 0.05$ ). However, in controls, there was no significant difference in the mean score of other dimensions before and after intervention ( $P > 0.05$ ). Before the intervention, the frequency distribution of various levels of sensitivity to interpersonal relationships in the two studied groups was the same ( $P > 0.05$ ); after the intervention, these levels were significantly different between groups ( $P < 0.05$ ). After intervention, the control group had a higher frequency distribution of individuals with degrees of depression ( $P < 0.05$ ).

**Conclusions:** The mental health of medical internship students during a crisis can be improved by stress management training, which reduces anxiety and depression. Therefore, stress management courses are recommended in stressful situations.

**Keywords:** Education; Stress Management; Mental Health; Student; Corona

## 1. Background

An emergency in global health in all countries of the world was caused by the unique spread of COVID-19 and its high transmission speed within a few months (1). This disease causes public health concerns and the occurrence of various psychological diseases (2, 3). Healthcare workers are the front line of fighting infectious diseases and COVID-19; they are the first to be exposed to this virus. So that, anxiety related to COVID-19 was experienced by 42.5% of Iranian dentists (4). During the COVID-19 epidemic in China, medical health workers had a higher risk of mental disorders than non-medical health workers (5). A significant proportion of health workers reported symp-

toms of anxiety, depression, and insomnia, as revealed by study findings in Nepal (6). In a systematic review in 2020, moderate to intense COVID-19-associated mental disorders were reported by a significant proportion of dentists (7). It was found that there is a significant correlation between physical illness and psychiatric disorders related to anxiety associated with COVID-19 (4). Also, the quality of healthcare workers' activities and services can be severely affected by psychological disorders like anxiety, fear, stress, and stigma (8, 9).

In addition to being concerned about the COVID-19 impact on reducing the learning of job skills, medical



students, as active participants in the fight against COVID-19, along with other healthcare workers, were exposed to various mental disorders (10-13), and were less likely to seek support (14, 15).

For all people influenced by the highly contagious COVID-19 virus, it is better to use appropriate psychotherapy methods such as stress management training using modern facilities such as videoconferencing and online programs. According to research, these methods in the treatment of anxiety, depression, and PTSD are effective (16-18).

Given that in a communicable disease epidemic, health workers and medical students are located at the first line of facing the infection, and they suffer high mental pressure. It's important to prioritize medical students' mental health and devise a plan to enhance it.

## 2. Objectives

This research assessed the effect of stress management training on the mental health of medical students and interns at Birjand City Medical Training Center during the COVID-19 pandemic in the academic year 2021.

## 3. Methods

### 3.1. Study Type and Statistical Population

The population of this randomized intervention study was the students of the training and internship levels studying at Birjand University of Medical Sciences throughout the 2020 - 2021 academic year.

### 3.2. Sample Size and Sampling Method

Shahraviet al.'s study (19) results were used to determine the sample size, and using Cochran's mean difference formula, the study population consisted of 60 people (30 in each group). It was then split into intervention and control groups randomly.

### 3.3. Inclusion and Exclusion Criteria

Internship students of Birjand University of Medical Sciences who had no history of depression and were accepted to take part in the study included in the study and uncomplete questionnaire, guest or transfer students from other universities, re-studying in other fields and students who had taken study leave, they were omitted from the study.

### 3.4. Gathering Data Tools and Methods

Students' mental health was evaluated using the SCL90 questionnaire. Dragotiset et al. introduced the SCL90 questionnaire. (20) in 1973 and was finalized in 1976.

Each SCL90 questionnaire item's answer is measured using a scale of 5 points of discomfort (none to severe). Scoring and interpretation of the test are based on three indicators of the general quotient of morbid symptoms, the measure of discomfort quotient, and the sum of morbid symptoms. The reliability of the 9 dimensions of the SCL90 questionnaire was measured using two reliability calculation methods, the retest method.

To calculate the internal consistency of the test, which was conducted on 219 people in the United States to measure the stability and uniformity of the questions, alpha coefficients and Coder Richardson 200 were used. The results of all the obtained coefficients for 9 dimensions were completely satisfactory (21).

To calculate the nine dimensions of the questionnaire, the scores of the questions of different dimensions were added together: Physical complaints dimension (12 questions), obsessive-compulsive dimension (10 questions), sensitivity in interpersonal relationships dimension (9 questions), depression dimension (13 questions), anxiety dimension (10 questions), aggression dimension (6 questions), fear and morbidity dimension (7 questions), paranoid dimension (6 questions) and psychotic dimension (10 questions).

In addition, to obtain the levels of different dimensions of the SCL90 questionnaire, the average score for each dimension is calculated by multiplying the number of questions. If the average score is below 1, there is no mental disorder; between 1 and 2, with a mild mental disorder; between 2 and 3, with a moderate mental disorder or significant and between 3 and 4 were considered serious or severe mental disorders (22).

The validity of this test has been determined through several studies. In the context of concurrent validity, Dragotis and Wikelzorak (1976) conducted this test together with the MMPI on 19 volunteer subjects (20).

The Persian form of the questionnaire mentioned was validated in Iran for its validity and reliability by Akhavan Abeiri et al. in 2018, and all dimensions of the questionnaire had a Cronbach's alpha coefficient above 0.75 (21).

### 3.5. Method of Conducting the Study

After justifying the students and completing the informed consent form, 60 eligible individuals were selected and were randomly divided into two study groups. Both groups completed the demographic checklist (age and gender) and the SCL90 questionnaire.

Then, the experimental group underwent a stress management training course of 6 sessions 45 minutes, 1 session per week (Table 1), taught by a specialized doctorate in education management and a master's degree in psychology. There was no intervention given to the control group during this period. Seven days after the

last session in the Corona department, the SCL90 questionnaire was completed again by both groups.

**Table 1.** Workshop Schedule for Students Participating in the Intervention Group

Session Number	Topic
1	Offering an introduction to the necessity and importance of training stress control skills, as well as defining stress, addressing the differences between people in how they deal with it, and the reasons behind them.
2	The physical, psychological, and behavioural effects of stress are analyzed in order to discuss how stress affects various body systems as a whole.
3	Finding mental and rational ways to handle stress, introducing problem-oriented and emotion-oriented methods as coping strategies, and exploring people's coping strategies in stressful situations.
4	Introducing the stages of coping with stress, focusing on the initial step of understanding your emotions, as well as developing study skills, test preparation, and time management.
5	Enhancing self-esteem, self-confidence, and coping with depression and anxiety.
6	Dealing with the 2nd step of stress control skills and remembering both long-term and short-term strategies.

### 3.6. Data Analysis

SPSS 22 software was utilized for data analysis. Central and dispersion indicators were used to report descriptive results. To verify the data's normality, the Kolmogorov-Smirnov test was employed. Analytical analysis was done by a 2-sided Mann-Whitney U-test, Paired Wilcoxon test, Chi-square test or Fisher's exact test at a significance level of  $\alpha = 0.05$ .

### 3.7. Ethical Considerations

This study was approved by the ethics committee of Birjand University of Medical Sciences (IR.BUMS.REC.1400.236). Furthermore, informed consent was obtained.

## 4. Results

This study included 60 medical students with  $25 \pm 0.83$  (intervention) and  $25.4 \pm 1.06$  (control) years of age, without significant differences between groups ( $P = 0.112$ ). Also, the gender frequency distribution was not significantly different ( $P = 0.435$ ) (Table 2).

**Table 2.** Comparison of Gender Frequency Distribution of People in the Two Studied Groups a

Group	Intervention	Control	Chi-square Test
Sex			
Male	19 (63.3)	15 (50)	$X^2 = 1.086, P = 0.435$
Female	11 (36.7)	15 (50)	

<sup>a</sup> Values are presented as No. (%).

The intervention did not significantly alter the average scores of physical complaints, obsessive-compulsive disorder, interpersonal relationships, depression, anxiety, aggression, and fear ( $P > 0.05$ ). After the intervention, the average scores of the paranoid and psychotic dimensions were not significant ( $P > 0.05$ ). However, before the intervention, the intervention group had a significantly higher level than the control group ( $P < 0.05$ ). In addition, after the intervention, the average dimensions of physical complaint, obsessive-compulsive, interpersonal

relationships, depression, anxiety, aggression, fear, paranoid and psychosis in the intervention group and the dimensions of obsessive-compulsive, depression, anxiety and aggression compared to the beginning of the study significantly decreased in the control group ( $P < 0.05$ ).

Furthermore, after intervention, the mean score of physical complaints, interpersonal relationships, fear, paranoia, and psychotic dimensions did not differ significantly compared to the beginning of the study in the control group ( $P > 0.05$ ) (Table 3).

**Table 3.** Comparison of the Average of Different Dimensions of Mental Health Before and After the Intervention in the Two Studied Groups

Group and Dimension	Intervention	Control	P-Value, (Mann-Whitney U test)
Physical complaints (Somatization)			
Before intervention	$7.16 \pm 6.68$	$5.7 \pm 4.41$	0.393
After intervention	$4.46 \pm 5.08$	$5.56 \pm 4.34$	0.352

<b>Wilcoxon signed-rank test</b>	< 0.001	0.612	
<b>Obsessive-compulsive</b>			
Before intervention	11.03 ± 6.00	9.00 ± 4.33	0.247
After intervention	8.00 ± 3.88	8.00 ± 3.96	0.835
<b>Wilcoxon signed-rank test</b>	< 0.001	0.007	
<b>Sensitivity in interpersonal relationships, (Interpersonal sensibility)</b>			
Before intervention	8.33 ± 5.07	5.76 ± 4.40	0.095
After intervention	6.16 ± 4.62	5.76 ± 4.40	0.835
<b>Wilcoxon signed-rank test</b>	< 0.001	0.204	
<b>Depression</b>			
Before intervention	12.16 ± 9.84	10.96 ± 7.36	0.824
After intervention	8.10 ± 7.20	9.80 ± 6.92	0.235
<b>Wilcoxon signed-rank test</b>	< 0.001	0.013	
<b>Anxiety</b>			
Before intervention	6.13 ± 5.51	5.30 ± 3.08	0.829
After intervention	4.50 ± 4.38	4.56 ± 3.23	0.526
<b>Wilcoxon signed-rank test</b>	< 0.001	0.013	
<b>Aggression (Anger-hostility)</b>			
Before intervention	4.26 ± 3.89	2.70 ± 2.05	0.125
After intervention	3.10 ± 2.95	0.50 ± 1.91	0.161
<b>Wilcoxon signed-rank test</b>	< 0.001	0.006	
<b>Fear (Phobic-anxiety)</b>			
Before intervention	3.70 ± 4.81	2.80 ± 2.44	0.928
After intervention	2.76 ± 3.61	2.36 ± 2.51	0.769
<b>Wilcoxon signed-rank test</b>	0.006	0.176	
<b>Paranoid ideation</b>			
Before intervention	7.56 ± 3.82	5.10 ± 2.68	0.011
After intervention	5.06 ± 2.93	4.53 ± 2.55	0.617
<b>Wilcoxon signed-rank test</b>	< 0.001	0.066	
<b>Psychoticism</b>			
Before intervention	7.76 ± 5.39	5.36 ± 2.52	0.045
After intervention	8.33 ± 4.35	5.40 ± 2.44	0.665
<b>Wilcoxon signed-rank test</b>	0.001	0.974	

a Values are presented as Mean ± SD.

The frequency distribution of different levels of physical complaint, obsessive-compulsive, anxiety, morbid fear, aggression, paranoia and psychosis before and after

the intervention in the studied groups was not significantly different ( $P > 0.05$ ). (Table 4).

**Table 4.** Comparing the distribution of the frequency distribution of physical complaints of obsessive-compulsive before and after the intervention in the two studied groups a

Group, Dimension			Intervention	Control	Fishers exact test
<b>Physical complaints (Somatization)</b>	Before intervention	Normal	28 (93.4)	28 (93.4)	$X^2 = 1.347, P = 1.000$
		Mild disorder	1 (3.3)	2 (6.6)	
		Significant disruption	1 (3.3)	0 (0.0)	
	After intervention	Normal	28 (93.4)	27 (90)	$X^2 = 0.218, P = 1.000$
		Mild disorder	2 (6.6)	3 (10)	

<b>Obsessive-compulsive</b>	Before interven- tion	Normal	15 (50)	15 (50)	X <sup>2</sup> = 2.034, P = 0.796		
		Mild disorder	13 (43.3)	15 (50)			
		Significant disruption	1 (3.3)	0 (0)			
		severe disorder	1 (3.3)	0 (0)			
<b>Sensitivity in interpersonal relation- ships, (Interpersonal sensibility)</b>	After interven- tion	Normal	19 (63.3)	18 (60)	X <sup>2</sup> = 0.071, P = 1.000		
		Mild disorder	11 (36.7)	12 (40)			
		Before interven- tion	Normal	17 (56.7)		19 (63.3)	X <sup>2</sup> = 1.764, P = 0.581,
			Mild disorder	11 (36.7)		11 (36.7)	
<b>Depression</b>	After interven- tion	Significant disruption	2 (6.7)	0 (0)	X <sup>2</sup> = 9.429, P = 0.004		
		Normal	25 (83.3)	17 (56.7)			
		Mild disorder	3 (10)	13 (43.3)			
		Before interven- tion	Normal	22 (73.3)		18 (60)	X <sup>2</sup> = 3.976, P = 0.102
Mild disorder	6 (20)		12 (40)				
Serious disorder	2 (6.7)		0 (0)				
Normal	26 (86.7)		18 (60)				
<b>Anxiety</b>	After interven- tion	Mild disorder	2 (6.7)	125 (40)	X <sup>2</sup> = 10.356, P = 0.003		
		Significant disruption	2 (6.7)	0 (0)			
		Before interven- tion	Normal	27 (90)		28 (93.3)	X <sup>2</sup> = 2.058, P = 0.612
			Mild disorder	1 (3.3)		2 (6.7)	
Significant disruption	2 (6.7)		0 (0)				
Normal	28 (93.3)		29 (96.7)				
<b>Aggression (Anger-hostility)</b>	After interven- tion	Mild disorder	2 (6.7)	1 (3.3)	X <sup>2</sup> = 0.357, P = 1.000		
		Significant disruption	2 (6.7)	0 (0)			
		Before inter- vention	Normal	22 (73.2)		27 (90)	X <sup>2</sup> = 3.097, P = 0.216
			Mild disorder	6 (20)		3 (10)	
Significant disruption	2 (6.7)		0 (0)				
Normal	26 (86.7)		29 (96.7)				
<b>Fear (Phobic-anxiety)</b>	After interven- tion	Mild disorder	2 (6.7)	1 (3.3)	X <sup>2</sup> = 2.201, P = 0.418		
		Significant disruption	2 (6.7)	0 (0)			
		Before inter- vention	Normal	25 (83.3)		28 (93.3)	X <sup>2</sup> = 2.061, P = 0.506
			Mild disorder	3 (10)		2 (6.7)	
Significant disruption	2 (6.7)		0 (0)				
Normal	26 (86.7)		28 (93.3)				
<b>Paranoid ideation</b>	After interven- tion	Mild disorder	2 (6.7)	1 (6.7)	X <sup>2</sup> = 1.825, P = 0.548		
		Significant disruption	2 (7.9)	0 (0)			
		Before inter- vention	Normal	9 (30)		18 (60)	X <sup>2</sup> = 5.873, P = 0.050
			Mild disorder	17 (56.7)		11 (36.7)	
Significant disruption	4 (13.3)		1 (3.3)				
Normal	17 (56.7)		18 (60)				
<b>Psychoticism</b>	After interven- tion	Mild disorder	11 (36.7)	12 (40)	X <sup>2</sup> = 1.725, P = 0.635		
		Significant disruption	2 (6.7)	0 (0)			
		Before inter- vention	Normal	25 (83.3)		29 (96.7)	X <sup>2</sup> = 2.878, P = 0.195
			Mild disorder	4 (13.3)		1 (3.3)	
Significant disruption	1 (3.3)		0 (0)				
Normal	28 (93.3)		29 (96.7)				
	After interven- tion	Mild disorder	2 (6.7)	1 (3.3)	X <sup>2</sup> = 2.580, P = 0.492		
		Significant disruption	2 (6.7)	0 (0)			

normal, Mild disorder, Significant disruption, severe disorder, Serious disorder

<sup>a</sup> Values are presented as No. (%).

Before intervention, no significant difference in the frequency distribution of different levels of sensitivity in interpersonal relationships in the studied groups was shown ( $P > 0.05$ ); however, after intervention, different levels of sensitivity in interpersonal relationships were significantly different between the groups.

The frequency distribution of students with degrees of depression did not differ significantly between groups before intervention ( $P > 0.05$ ). However, following the intervention, the intervention group had a significantly higher level ( $P < 0.05$ ).

## 5. Discussion

Study results showed that after the intervention, the average scores of the dimensions of physical complaints, obsessive compulsiveness, interpersonal relationships, depression, anxiety, aggression, fear, paranoid and psychotic in the intervention group, and the average scores of dimensions of obsessive-compulsive disorder, depression, anxiety, and aggression in the control group decreased significantly ( $P < 0.05$ ).

In addition, after intervention with different levels of interpersonal relationship sensitivity, the two groups showed a significant difference ( $P < 0.05$ ). In contrast, the intervention led to a significant increase in the frequency distribution of individuals with depression degrees in the control group ( $P < 0.05$ ).

In agreement with our study, Narimani et al. (23), Dijkstra et al. (24), Tejad Gelardo et al. (25), and Anuri et al. (26) showed a significant decrease in mental health scores vs. the study beginning ( $P < 0.05$ ). Also, Kordani et al. found that the average anxiety score after the intervention in the nurses decreased significantly (27).

Dincer et al. reported that educational intervention improved the state of nurses' mental health by reducing stress, anxiety, and job burnout during the coronavirus pandemic, which was in agreement with our findings (28). According to Xu et al., the intervention significantly improved mental health and its dimensions in health-care workers through the SCL90 questionnaire (29).

Similar to Hamsour, Lerardi et al. in 2021 found that after the intervention, the average score of depression, interpersonal sensitivity, anxiety, and obsessive-compulsive dimensions in the studied students decreased significantly (30).

In addition to the hard work and the stressful nature of health workers' jobs, the chances of contracting COVID-19 infection are higher (6). The mental and psychological conditions of health workers are more affected by the absence of adequate personal protection facilities due to the fact that many people infected with COVID-19 are symptomatic (6, 9, 31-36)

Mental health outcomes impact health workers' work performance, and specialized mental health services are necessary to address this (32, 37, 38). It is important to prioritize the mental well-being of health workers involved

in the COVID-19 response (6).

According to global studies, psychological support is necessary to enhance resiliency against negative mental health outcomes (38-41). Therefore, governments should consider stress management courses as one of the preventive measures (6).

According to the review conducted by Farzani et al., therapists around the world used various psychological interventions such as cognitive behavioural therapy, especially online, emotional release techniques, mindfulness, emotion regulation techniques, and neurofeedback to treat the mentioned psychological injuries (42). By managing anxiety, medical students at the university can improve their hardiness and self-efficacy to handle various life challenges (43). An issue that should be considered is the difference in background conditions (individual, social, cultural and economic). Also, many people's reluctance to express their psychological problems and problems has affected their responses (18).

In general, the increase in anxiety and public concern following the pandemic of infectious diseases, especially the unknown genital infection, leads to a disturbance in the understanding of disease-related issues and other psycho-social challenges, including social stigma and discrimination, which require more attention (44, 45).

People experienced a high level of fear and anxiety due to the Coronavirus pandemic and lack of definitive treatment. On the other hand, this point should be mentioned even though the Coronavirus disease is a deadly disease with a very high prevalence. For several years, the pandemic imposed many economic, social and psychological consequences on the people.

### 5.1. Strengths and Limitations

This study's strength is its implementation of an interventional study for high-risk groups in the COVID-19 crisis period. The Coronavirus pandemic has led to the study's main limitations, which include the inability to provide complete and appropriate access to the students studied and the study's focus on medical students.

### 5.2. Conclusions

Stress management training improves the mental health of medical students during a crisis by decreasing anxiety and depression in them. Improving mental health and reducing anxiety in health workers and medical students during critical situations like COVID-19 can be obtained through stress management training as an effective intervention.

Authors' Contribution:

Conceptualization: Sobhan Abbasi, Yahya Mohammadi, Mohammadreza Raeisoon; Formal analysis: Yahya Mohammadi, Mohammadreza Raeisoon; Data curation: Sobhan Abbasi, Yahya Mohammadi; Project administration: Mohammadreza Raeisoon; Original draft writing: all authors; Review writing and editing: all authors.

**Conflict of Interests:**

There is no competition.

**Data Reproducibility:**

Data requests can be made to the corresponding author.

**Ethical Approval:**

Birjand University of Medical Sciences ethics committee approved this study (IR.BUMS.REC.1400.236, <https://ethics.research.ac.ir/ProposalCertificateEn.php?id=229864>). Furthermore, all participants in the study were given informed consent.

**Funding/Support:**

It was not declared by the authors. This study has no funding support.

**References**

- Zangrillo A, Beretta L, Silvani P, Colombo S, Scandroglio AM, Dell'Acqua A, et al. Fast reshaping of intensive care unit facilities in a large metropolitan hospital in Milan, Italy: Facing the COVID-19 pandemic emergency. *Crit Care Resusc.* 2020;**22**(2):91-4. [PubMed ID:32227819].
- Yang L, Wu D, Hou Y, Wang X, Dai N, Wang G, et al. Analysis of psychological state and clinical psychological intervention model of patients with COVID-19. 2020. <https://doi.org/10.1101/2020.03.22.20040899>.
- Liu S, Yang L, Zhang C, Xiang YT, Liu Z, Hu S, et al. Online mental health services in China during the COVID-19 outbreak. *Lancet Psychiatry.* 2020;**7**(4):e17-e8. [PubMed ID:32085841]. [PubMed Central ID:PMC7129099]. [https://doi.org/10.1016/S2215-0366\(20\)30077-8](https://doi.org/10.1016/S2215-0366(20)30077-8).
- Salehiniya H, Abbaszadeh H. Prevalence of corona-associated anxiety and mental health disorder among dentists during the COVID-19 pandemic. *Neuropsychopharmacol Rep.* 2021;**41**(2):223-9. [PubMed ID:33825340]. [PubMed Central ID:PMC8250041]. <https://doi.org/10.1002/npr2.12179>.
- Zhang WR, Wang K, Yin L, Zhao WF, Xue Q, Peng M, et al. Mental Health and Psychosocial Problems of Medical Health Workers during the COVID-19 Epidemic in China. *Psychother Psychosom.* 2020;**89**(4):242-50. [PubMed ID:32272480]. [PubMed Central ID:PMC7206349]. <https://doi.org/10.1159/000507639>.
- Khanal P, Devkota N, Dahal M, Paudel K, Joshi D. Mental health impacts among health workers during COVID-19 in a low resource setting: a cross-sectional survey from Nepal. *Global Health.* 2020;**16**(1):89. [PubMed ID:32977818]. [PubMed Central ID:PMC7517059]. <https://doi.org/10.1186/s12992-020-00621-z>.
- Salehiniya H, Hatamian S, Abbaszadeh H. Mental health status of dentists during COVID-19 pandemic: A systematic review and meta-analysis. *Health Sci Rep.* 2022;**5**(3):e617. [PubMed ID:35509394]. [PubMed Central ID:PMC9059210]. <https://doi.org/10.1002/hsr2.617>.
- Li L, Wan C, Ding R, Liu Y, Chen J, Wu Z, et al. Mental distress among Liberian medical staff working at the China Ebola Treatment Unit: a cross sectional study. *Health Qual Life Outcomes.* 2015;**13**:156. [PubMed ID:26409446]. [PubMed Central ID:PMC4583730]. <https://doi.org/10.1186/s12955-015-0341-2>.
- Wu P, Fang Y, Guan Z, Fan B, Kong J, Yao Z, et al. The psychological impact of the SARS epidemic on hospital employees in China: exposure, risk perception, and altruistic acceptance of risk. *Can J Psychiatry.* 2009;**54**(5):302-11. [PubMed ID:19497162]. [PubMed Central ID:PMC3780353]. <https://doi.org/10.1177/070674370905400504>.
- Eyni S, Ebadi M, Torabi N. Developing a model of corona anxiety in students based on optimism and resilience: The mediating role of the perceived social support. *Counseling Culture and Psychotherapy.* 2020;**11**(43):1-32.
- Wu Z, McGoogan JM. Characteristics of and Important Lessons From the Coronavirus Disease 2019 (COVID-19) Outbreak in China: Summary of a Report of 72 314 Cases From the Chinese Center for Disease Control and Prevention. *JAMA.* 2020;**323**(13):1239-42. [PubMed ID:32091533]. <https://doi.org/10.1001/jama.2020.2648>.
- Mittal R, Su L, Jain R. COVID-19 mental health consequences on medical students worldwide. *J Community Hosp Intern Med Perspect.* 2021;**11**(3):296-8. [PubMed ID:34234896]. [PubMed Central ID:PMC8118449]. <https://doi.org/10.1080/20009666.2021.1918475>.
- Chandratre S. Medical Students and COVID-19: Challenges and Supportive Strategies. *J Med Educ Curric Dev.* 2020;**7**:2382120520935059. [PubMed ID:32637642]. [PubMed Central ID:PMC7315659]. <https://doi.org/10.1177/2382120520935059>.
- Xiong P, Ming WK, Zhang C, Bai J, Luo C, Cao W, et al. Factors Influencing Mental Health Among Chinese Medical and Non-medical Students in the Early Stage of the COVID-19 Pandemic. *Front Public Health.* 2021;**9**:603331. [PubMed ID:34095044]. [PubMed Central ID:PMC8172592]. <https://doi.org/10.3389/fpubh.2021.603331>.
- Essangri H, Sabir M, Benkabbou A, Majbar MA, Amrani L, Ghanam A, et al. Predictive Factors for Impaired Mental Health among Medical Students during the Early Stage of the COVID-19 Pandemic in Morocco. *Am J Trop Med Hyg.* 2021;**104**(1):95-102. [PubMed ID:33205748]. [PubMed Central ID:PMC7790070]. <https://doi.org/10.4269/ajtmh.20-1302>.
- Rees CS, MacLaine E. A Systematic Review of Videoconference-Delivered Psychological Treatment for Anxiety Disorders. *Australian Psychologist.* 2020;**50**(4):259-64. <https://doi.org/10.1111/ap.12122>.
- García-Lizana F, Muñoz-Mayorga I. Telemedicine for depression: a systematic review. *Perspect Psychiatr Care.* 2010;**46**(2):19-26. [PubMed ID:20377799]. <https://doi.org/10.1111/j.1744-6163.2010.00247.x>.
- Turgoose D, Ashwick R, Murphy D. Systematic review of lessons learned from delivering teletherapy to veterans with post-traumatic stress disorder. *J Telemed Telecare.* 2018;**24**(9):575-85. [PubMed ID:28958211]. <https://doi.org/10.1177/1357633X17730443>.
- Shahroie A, Mehrabizadeh Honarmand M, Mohammad Hassan Adel S, Shahroie M, Shahroey S. [The effectiveness of cognitive-behavioral stress management intervention on general health patients with cardiovascular disease]. *Jundi Sci Med J.* 2017;**15**(6):757-66. Persian.
- Eivors A, Button E, Warner S, Turner K, Fassino S, Abbate-Daga G, et al. Hillsdale, NJ: Erlbaum. Derogatis, LR, Lipman, RS, & Covi, L.(1973). Scl-90: An outpatient psychiatric rating scale—preliminary report. *Psychopharmacology Bulletin*, 9, 13-28. Di Pietro, G., Valoroso, L., Fichelle, M., Bruno, C., & Sorge, F.(2002). What happens to eating disorder outpatients who withdrew from therapy? *Eating & Weight. Measuring Eating Disorder Outcome.* 2008;**81**:29-35.
- Akhavan Abiri F, Shairi MR. [Validity and reliability of symptom checklist-90-revised (SCL-90-R) and brief symptom inventory-53 (BSI-53)]. *Clinical Psychology and Personality.* 2020;**17**(2):169-95. Persian.
- Tabrizizadeh M, Yasini Ardakani SM, Rostamzade P, Zare M. [The Mental Health Status of Students of Medicine and Dentistry A Study in Shahid Sadoughi University of Medical Sciences Yazd Iran]. *Strides in Develop Med Edu.* 2013;**9**(2):153-61. Persian.
- Narimani F, Azmoudeh M, Livar Jani S, MeserAabadi J. [Comparison of the Effectiveness of Positivist Psychology group education and Metacognition group education on the academic Achievement Motivation of Quarantined female Students with their families]. *Women Family Studies.* 2023;**16**(60). Persian.
- Schotanus-Dijkstra M, Keyes CLM, de Graaf R, Ten Have M. Recovery from mood and anxiety disorders: The influence of positive mental health. *J Affect Disord.* 2019;**252**:107-13. [PubMed ID:30981053]. <https://doi.org/10.1016/j.jad.2019.04.051>.
- Tejada-Gallardo C, Blasco-Belled A, Torrelles-Nadal C, Alsinet C. Effects of School-based Multicomponent Positive Psychology Interventions on Well-being and Distress in Adolescents: A Systematic Review and Meta-analysis. *J Youth Adolesc.* 2020;**49**(10):1943-60. [PubMed ID:32683592]. <https://doi.org/10.1007/s10964-020-01289-9>.
- Anvari J, Sardary B. [The Effectiveness of positive psychotherapy on improving cognitive emotion regulation strategies in male secondary high school students with anxiety symptoms]. *Positive Psychol Res.* 2019;**5**(2):31-46. Persian.
- Kardani S, DashtBozorgi Z. [The Effect of Positive Mindfulness Education on Health Anxiety, Feeling of Happiness and Health

- Adjustment of Nurses in During the Covid-19 Pandemic]. *Iran J Psychiatric Nurs.* 2022;**10**(3):53-64. Persian.
28. Dincer B, Inangil D. The effect of Emotional Freedom Techniques on nurses' stress, anxiety, and burnout levels during the COVID-19 pandemic: A randomized controlled trial. *Explore (NY).* 2021;**17**(2):109-14. [PubMed ID:33293201]. [PubMed Central ID:PMC7834511]. <https://doi.org/10.1016/j.explore.2020.11.012>.
  29. Xu J, Liu X, Xiao Y, Fang X, Cheng Y, Zhang J. Effect of EAP Psychological Intervention on Improving the Mental Health of Medical Workers Under the Novel Coronavirus Epidemic in China. *Front Public Health.* 2021;**9**:649157. [PubMed ID:34395355]. [PubMed Central ID:PMC8355514]. <https://doi.org/10.3389/fpubh.2021.649157>.
  30. Ierardi E, Bottini M, Riva Crugnola C. Effectiveness of an online versus face-to-face psychodynamic counselling intervention for university students before and during the COVID-19 period. *BMC Psychol.* 2022;**10**(1):35. [PubMed ID:35193671]. [PubMed Central ID:PMC8861610]. <https://doi.org/10.1186/s40359-022-00742-7>.
  31. Lai J, Ma S, Wang Y, Cai Z, Hu J, Wei N, et al. Factors Associated With Mental Health Outcomes Among Health Care Workers Exposed to Coronavirus Disease 2019. *JAMA Netw Open.* 2020;**3**(3):e203976. [PubMed ID:32202646]. [PubMed Central ID:PMC7090843]. <https://doi.org/10.1001/jamanetworkopen.2020.3976>.
  32. Kang L, Ma S, Chen M, Yang J, Wang Y, Li R, et al. Impact on mental health and perceptions of psychological care among medical and nursing staff in Wuhan during the 2019 novel coronavirus disease outbreak: A cross-sectional study. *Brain Behav Immun.* 2020;**87**:11-7. [PubMed ID:32240764]. [PubMed Central ID:PMC7118532]. <https://doi.org/10.1016/j.bbi.2020.03.028>.
  33. Liu Q, Luo D, Haase JE, Guo Q, Wang XQ, Liu S, et al. The experiences of health-care providers during the COVID-19 crisis in China: a qualitative study. *Lancet Glob Health.* 2020;**8**(6):e790-e8. [PubMed ID:32573443]. [PubMed Central ID:PMC7190296]. [https://doi.org/10.1016/S2214-109X\(20\)30204-7](https://doi.org/10.1016/S2214-109X(20)30204-7).
  34. Liu X, Kakade M, Fuller CJ, Fan B, Fang Y, Kong J, et al. Depression after exposure to stressful events: lessons learned from the severe acute respiratory syndrome epidemic. *Compr Psychiatry.* 2012;**53**(1):15-23. [PubMed ID:21489421]. [PubMed Central ID:PMC3176950]. <https://doi.org/10.1016/j.compsych.2011.02.003>.
  35. Lee SM, Kang WS, Cho AR, Kim T, Park JK. Psychological impact of the 2015 MERS outbreak on hospital workers and quarantined hemodialysis patients. *Compr Psychiatry.* 2018;**87**:123-7. [PubMed ID:30343247]. [PubMed Central ID:PMC7094631]. <https://doi.org/10.1016/j.comppsy.2018.10.003>.
  36. Day M. Covid-19: four fifths of cases are asymptomatic, China figures indicate. *BMJ.* 2020;**369**:m1375. [PubMed ID:32241884]. <https://doi.org/10.1136/bmj.m1375>.
  37. Weaver MD, Vetter C, Rajaratnam SMW, O'Brien CS, Qadri S, Benca RM, et al. Sleep disorders, depression and anxiety are associated with adverse safety outcomes in healthcare workers: A prospective cohort study. *J Sleep Res.* 2018;**27**(6):e12722. [PubMed ID:30069960]. [PubMed Central ID:PMC6314290]. <https://doi.org/10.1111/jsr.12722>.
  38. Du J, Dong L, Wang T, Yuan C, Fu R, Zhang L, et al. Psychological symptoms among frontline healthcare workers during COVID-19 outbreak in Wuhan. *Gen Hosp Psychiatry.* 2020;**67**:144-5. [PubMed ID:32381270]. [PubMed Central ID:PMC7194721]. <https://doi.org/10.1016/j.genhosppsych.2020.03.011>.
  39. Chen Q, Liang M, Li Y, Guo J, Fei D, Wang L, et al. Mental health care for medical staff in China during the COVID-19 outbreak. *Lancet Psychiatry.* 2020;**7**(4):e15-e6. [PubMed ID:32085839]. [PubMed Central ID:PMC7129426]. [https://doi.org/10.1016/S2215-0366\(20\)30078-X](https://doi.org/10.1016/S2215-0366(20)30078-X).
  40. Liu CY, Yang YZ, Zhang XM, Xu X, Dou QL, Zhang WW, et al. The prevalence and influencing factors in anxiety in medical workers fighting COVID-19 in China: a cross-sectional survey. *Epidemiol Infect.* 2020;**148**:e98. [PubMed ID:32430088]. [PubMed Central ID:PMC7251286]. <https://doi.org/10.1017/S0950268820001107>.
  41. Park SC, Park YC. Mental Health Care Measures in Response to the 2019 Novel Coronavirus Outbreak in Korea. *Psychiatry Investig.* 2020;**17**(2):85-6. [PubMed ID:32093458]. [PubMed Central ID:PMC7047003]. <https://doi.org/10.30773/pi.2020.0058>.
  42. Farzani M, Shalbfaf ZKH, Zandkarimi G. [Evaluation of the effectiveness of psychological interventions on anxiety and depression during the COVID-19 pandemic: A systematic review study]. *Rooyesh-e-Ravanshenasi J (RRJ).* 2022;**11**(2):181-90. Persian.
  43. Sahranavard S, Esmaili A, Dastjerdi R, Salehiniya H. The effectiveness of stress-management-based cognitive-behavioral treatments on anxiety sensitivity, positive and negative affect and hope. *Biomedicine (Taipei).* 2018;**8**(4):23. [PubMed ID:30474604]. [PubMed Central ID:PMC6254097]. <https://doi.org/10.1051/bm-dcn/2018080423>.
  44. Wang D, Hu B, Hu C, Zhu F, Liu X, Zhang J, et al. Clinical Characteristics of 138 Hospitalized Patients With 2019 Novel Coronavirus-Infected Pneumonia in Wuhan, China. *JAMA.* 2020;**323**(11):1061-9. [PubMed ID:32031570]. [PubMed Central ID:PMC7042881]. <https://doi.org/10.1001/jama.2020.1585>.
  45. Pfefferbaum B, North CS. Mental Health and the Covid-19 Pandemic. *N Engl J Med.* 2020;**383**(6):510-2. [PubMed ID:32283003]. <https://doi.org/10.1056/NEJMp2008017>.