

## The Ameliorating Effects of Garlic (*Allium Sativum*) on Blood Glucose Levels and Lipid-Related Indices

Fatemeh Samimi<sup>1,2</sup>, Ali Sharifi-Rigi<sup>2,3</sup>, Sanaz Dastghaib<sup>4</sup>, Narjes Hazar<sup>1</sup>, Fatemeh Zal<sup>2</sup>, Morvarid Siri<sup>5</sup>, Nasim Namiranian<sup>1</sup>, Mohammad Afkhami-Ardekani<sup>1\*</sup>

<sup>1</sup>Diabetes Research Center, Shahid Sadoughi University of Medical Sciences, Yazd, Iran.

<sup>2</sup>Department of Biochemistry, School of Medicine, Shiraz University of Medical Sciences, Shiraz, Iran.

<sup>3</sup>Student Research Committee, Shiraz University of Medical Sciences, Shiraz, Iran.

<sup>4</sup>Endocrinology and Metabolism Research Center, Shiraz University of Medical Sciences, Shiraz, Iran.

<sup>5</sup>Autophagy Research Center, Department of Clinical Biochemistry, School of Medicine, Shiraz University of Medical Sciences, Shiraz, Iran.

### Abstract

Garlic (*Allium sativum* L.), a culinary plant with medicinal properties, has been utilized as a traditional remedy by people worldwide for centuries. Garlic possesses a variety of health benefits, such as its antibacterial and anticancer properties, its capacity to slow the aging process and prevent obesity, its antihypertensive, antioxidative, and cardioprotective properties, and its capacity to reduce blood sugar and cholesterol. Garlic possesses organosulfur compounds, which have been associated with advantageous and favorable impacts on health. This review specifically examines the therapeutic potential and molecular mechanisms of Garlic in influencing important processes related to the control of glucose and lipid metabolism. These effects have been observed in both laboratory studies and clinical trials. The processes encompassed within this category entail the excretion of insulin, the absorption of glucose, the accumulation of glycogen, and the synthesis of lipids in the liver. The results of this study investigating the impact of Garlic on glucose and lipid regulation will deepen our comprehension of this fascinating natural compound. This information will be invaluable for healthcare professionals, researchers, and individuals seeking to prevent and treat metabolic disorders.


**Keywords:** Garlic, Organic sulfides, Blood glucose, Lipid profile, Health benefits

### QR Code



**Citation:** Samimi F, Sharifi-Rigi A, Dastghaib S, Hazar N, Zal F, Siri M, et al . The Ameliorating Effects of Garlic (*Allium Sativum*) on Blood Glucose Levels and Lipid-Related Indices. IJDO 2025; 17 (2) :141-149

**URL:** <http://ijdo.ssu.ac.ir/article-1-956-en.html>

 10.18502/ijdo.v17i2.18852

### Article info:

**Received:** 3 January 2025

**Accepted:** 20 April 2025

**Published in May 2025**



This is an open access article under the (CC BY 4.0)

### Corresponding Author:

**Mohammad Afkhami-Ardekani**, Professor of Endocrinology, Diabetes Research Center, Shahid Sadoughi University of Medical Sciences and Health Services, Yazd, Iran.

**Email:** Afkhamiam@yahoo.com

**Orcid ID:** 0000-0001-9768-5904

**Tell:** (98) 353 728 0215

## Introduction

One of the main risk factors for the development of complications in diabetes is hyperglycemia, which can lead to retinopathy, neuropathy, nephropathy, and cardiovascular disease (1). In addition, dyslipidemia is a prevalent feature of several metabolic disorders, including obesity, diabetes, and fatty liver disease (NAFLD) (2,3). High triglycerides (TG) and total cholesterol (TC), a reduction in high-density lipoprotein cholesterol (HDL-c), a rise in the plasma level of low-density lipoprotein cholesterol (LDL-c), and very low-density lipoprotein (VLDL) are the most noticeable signs of dyslipidemia (4). An independent risk factor for atherosclerosis disease is elevated LDL-c levels (5). Time will reveal if hyperlipidemia can lead to catastrophic conditions including heart issues and stroke if left untreated (6).

As a result, several of these chemicals are used in the treatment and management of individuals with high blood sugar. Unfortunately, there aren't many therapy choices for these kinds of illnesses, and the majority of medications only address one specific issue. As a result, a number of medications that lower blood sugar and reduce cholesterol have been introduced to the market; however, these benefits are only momentary (7). These medications also cause adverse responses, which raise the possibility of hypoglycemic reactions and the accumulation of extra fat just for the purpose of temporarily relieving episodes related to diabetes and other metabolic problems (8). They have been studied and have received some notoriety lately because of their potential therapeutic efficacy when used to treat various illnesses. Modern traditional medicine uses the anti-inflammatory and antioxidant properties of garlic to treat a variety of ailments, including heart disease, cancer, diabetes mellitus, and infections like the common cold (9-11). This article provides a thorough analysis of these effects and the mechanisms involved in order to determine

how garlic may affect the treatment of diabetes and/or metabolic syndrome.

### Garlic as a therapeutic agent: unveiling the healing power

The *Allium* genus is well known for its garlic (*Allium sativum*). It is a flowering plant with several uses. Renowned for its numerous medicinal benefits in addition to its unique flavor and scent, which are used in preparing food (12). A range of organosulfur compounds have been increasingly credited by the pharmaceutical industry with their therapeutic effectiveness. Garlic contains a variety of naturally occurring therapeutic qualities due to its sulfur and non-sulfur components (13).

S-allyl-cysteine sulfoxide, or alliin, is referred to as the active ingredient in garlic by one of the leading bioactive scientists (14). This chemical goes through several stages of processing. Alliinase catalyzes a series of transformational processes including chewing, crushing, cutting, and extracting. The enzymatic process known as alliinase converts alliin into allicin (diallyl thiosulfinate), a highly active compound that serves as a foundation for several health benefits (15). Ajoene, allyl sulfides, and vinyl dithiines are among the other sulfur compounds produced during the transformation that are beneficial in medicine and together they contribute to the variety of biological activities of garlic (16).

Garlic (*Allium sativum*) contains a chemical called allicin, which gives it its distinct and strong smell. Attention is drawn to allicin because of its potential medicinal benefits and health benefits. Because of its potential medicinal benefits—such as its anti-oxidative (17), anti-microbial (18), and possible cardiovascular (19) and anticancer effects (20) researchers have shown interest in it. However, further clinical research has to be done to prove that Allicin can only be extensively utilized after patients are aware of the dosage, mode of administration, and potential adverse effects that are advised as a therapeutic drug.

Numerous health advantages of garlic include its antibacterial and anticancer properties, its ability to prevent obesity and aging, its antihypertensive, antioxidative, and cardioprotective properties, and its ability to decrease blood sugar and cholesterol. As evidenced by the literature, they have been extensively researched to determine these impacts. It could occur from a diet high in garlic, which has anti-obesity properties (21-23). However, Mohammad et al. (2014) shown that garlic can inhibit pre-adipocyte maturation, which would slow the formation of adipose tissue and thus fight obesity (24). Moreover, administering garlic to humans orally also works well. Rats fed a high-fat diet and exhibiting signs of hyperlipidemia and epididymal obesity lost a significant amount of body weight. These findings thus suggest that garlic offers an option in the battle against obesity (25). Because it prevents angiotensin from being converted to angiotensin II, it further improves heart health (26). It decreases blood pressure since it is an antihypertensive agent (22). Garlic's antioxidants protect the blood vessels and the heart, avoiding atherosclerosis (27). Numerous studies highlight the benefits of reducing diabetes and associated consequences as well as the function that garlic plays in nutrition in preventing atherosclerosis (28-30). S-allyl cysteine sulfoxide, which has been identified in garlic, has been shown to act as an insulin secreting agents in diabetic rats (31). Additionally, garlic indirectly lowers the risk of atherosclerosis by lowering blood pressure, cholesterol, and maybe even diabetes mellitus. It also inhibits the production of thrombi (32, 33). Moreover, garlic may help decrease cholesterol and blood sugar levels. It may be able to help control hyperlipidemia and diabetes (28,34). Although the effects of garlic on blood lipids and glucose have been well studied, the origins of garlic and the evolution of animal trials are better understood. Our analysis looks at the data that is currently available on gap closure. These results demonstrate that garlic has several health benefits in addition to being a dietary

ingredient. Thus, in order to provide a thorough understanding of the therapeutic implications of utilizing garlic in the treatment of metabolic health, this review incorporates findings from both animal and clinical investigations (Tables 1&2).

### **Underlying Mechanisms for the glucose and Lipid-Regulating Effects of Garlic**

Rats with normal health showed significant changes in serum concentrations of total cholesterol (TC), triglycerides (TG), and blood glucose after receiving 0.5 g/kg/day of raw garlic for 4 weeks (34). Black garlic (1.5, 3, and 6 mg/200 g Bw/day) for 25 days substantially decreased blood glucose, TG, TC, and LDL-c levels in Alloxan-induced diabetic rats, according to Isnaini et al.'s validation (35). In rats fed a fructose diet, Gargouri et al. observed that 8 weeks of exposure to raw garlic (250 mg/kg/day) significantly decreased plasma glucose, HbA1c, insulin resistance, and TG (36).

In Supakul L's experiment, rats that were obese and showed signs of insulin resistance were given dosages of 250 and 500 mg/kg/day of garlic extract for four weeks. Insulin levels, HOMA-IR (Homeostatic Model Assessment of Insulin Resistance), and total cholesterol (TC) were all significantly reduced by both garlic dosages (37). Another study showed that giving rats given high-cholesterol diets 5 grams of garlic per kilogram per day for 10 weeks significantly reduced their levels of TG, TC, and LDL-c (38). All of these results point to the effectiveness of garlic in raising insulin sensitivity as well as in lipid and glucose metabolism in animal models. Garlic significantly lowers fasting blood glucose (FBG) and HbA1c in diabetic individuals, according to research by Ebadi et al. The study's findings demonstrated the potential of garlic as a dietary supplement for the management and treatment of diabetes (39).

**Table 1. Animal studies of the glucose and lipid management of Garlic**

Animal	Interventions	Dose of Garlic	Duration (Week or day)	Results	Ref.
<b>High-fat fed rats</b>	Black Garlic extract	(15g /kg/day)	5 w	↓TG, TC ↓mRNA of SREBP-1c, ACC, FAS, G6PDH, HMG-CoA reductase, ACAT	(55)
<b>Normal rats</b>	Raw Garlic	0.5 g/kg/day	4 w	↓Glucose, TG, TC	(34)
<b>Alloxan induced diabetic rats</b>	Black Garlic	1.5, 3, 6 mg/200 g Bw/day	25 d	↓Blood glucose ↓TG, TC, LDL-c	(35)
<b>Alloxan induced diabetic rats</b>	Garlic extract	300 mg/kg/day	6 w	↓Blood glucose ↓Total lipid, TC	(57)
<b>Rats fed a high fructose</b>	Raw Garlic	250 mg/kg/day	8 w	↓Serum glucose, HbA1c, insulin resistance ↓TG	(36)
<b>Insulin-resistant obese rats</b>	Garlic extract	250 or 500 mg/kg/day	4 w	↓Insulin, HOMA-IR ↓TC	(37)
<b>Rats fed high cholesterol diets</b>	High temperature- and high pressure-processed Garlic	5g /kg/day	10 w	TC, LDL-c, TG	(38)
<b>NAFLD rats induced long-term high-fat diet</b>	Diallyl disulfide (DADS)	20 mg/kg/day	12 w	↓Body weight, Adipose tissue weight Down-regulation of SREBP-1c, ACC, FAS, and HMG-CoA reductase, Stimulation of PPAR $\alpha$ and CPT-1	(56)
<b>High-fat fed rats</b>	Garlic oil	92.6 mg/kg/day	60 d	↓Body weight, Adipose tissue weight ↓TC, TG and LDL-c ↑HDL-c	(58)
<b>High-fat/sucrose fed rats</b>	Aged black Garlic	250 mg/kg/day	8 w	↓Body weight, TG, LDL-c, ↑HDL-c ↓PPAR- $\gamma$ , LPL, HSL expression ↑INSR, GLUT-4 expression	(59)

↓: significantly decrease, ↑: significantly increase in the experimental group compared to the control group)

**Table 2. Clinical trials about Garlic and its beneficial effects on glucose and lipid parameters**

Participants	N sample (Case: control)	Interventions	Dose of Garlic (mg/day)	Duration (week)	Results	Ref.
<b>Subjects with type 2 diabetes</b>	30: 30	Garlic + Allicin tablet Control: (Vit B1 tablet)	2406	12	↓FBS ↓HbA1c	(39)
<b>Subjects with type 2 diabetes and obesity</b>	30: 30	Metformin + Garlic capsul Control: (Metformin)	500	12	↓FBS, TG, TC, LDL ↑HDL	(30)
<b>Subjects with type 2 diabetes</b>	30: 30	Garlic + Metformin Control: (Placebo+Metformin)	900	24	↓FBS, TG, TC, LDL ↑HDL	(40)
<b>Women with GDM</b>	26: 23	Garlic pill Control: Placebo	400	8	↓FBS	(41)
<b>Obese patients</b>	46: 46	Garlicin capsule Control: Placebo	400	12	↓TC, LDL	(60)
<b>Subjects with type 2 diabetes and hyperlipidemia</b>	40: 0	Garsin tablet Control: -	900	4	↓TC ↓LDL-c	(42)
<b>Subjects with type 2 diabetes</b>	20: 20	Allicore (Garlic powder) Control: placebo	600	1	↓FBS ↓TG	(43)
<b>Patients with dyslipidemia</b>	35:35	Garlic tablet Control: placebo	600	12	↓TC, LDL ↑HDL	(61)
<b>Patients with NAFLD</b>	45: 43	Garlic tablet Control: placebo	1600	12	↓TG, TC, LDL ↑HDL	(44)

↓: significantly decrease, ↑: significantly increase in the intervention group compared to the control group)

In comparison to the control group, Kumar and Ashraf showed that garlic dramatically decreased the blood levels of FBG, TG, TC, and LDL-c and increased the levels of HDL-c in people with T2DM. According to the findings of these research, people with diabetes can better control their blood sugar and hyperlipidemia by taking garlic with their

regular anti-diabetic medications (30,40). According to research by Faroughi et al., giving women with gestational diabetes 400 mg of garlic daily for eight weeks lowers their diastolic blood pressure, prediabetes symptoms, and FBS levels (41). According to a research by Afkhami et al., supplementing with garlic lowers LDL-c and total cholesterol in



those with Type 2 diabetes (T2DM). The results of this investigation showed no discernible variation between the levels of FBG and 2hpp glucose before and after garlic consumption. After consuming garlic, HDL-c levels increased, but not significantly (42). In the Sobenin research, FBG, serum triglyceride, and fructosamine levels decreased after a week of therapy with 600 mg/day of Allicore powder tablets. In addition to diet therapy, the study's findings suggest that Allicore Garlic powder pills can be helpful in the management and treatment of type 2 diabetes (43). A different investigation on NAFLD patients found that HDL-c increased while TG, TC, and LDL-c decreased (44).

Garlic could reduce blood sugar levels and enhance lipid profiles through a number of mechanisms, according to a number of pre-clinical and trial investigations, even if the exact mechanism behind its anti-glycemic and anti-lipidemic effects is unknown. Organosulfur compounds (OSC) from garlic may function like insulin, reduce insulinase activity, and enhance insulin sensitivity (45). Moreover, it boosts insulin release and promotes the growth and regeneration of pancreatic beta cells. One theory for the mechanism of action is that the allicin in garlic prevents insulin inactivation by reacting with molecules that include sulfhydryl groups that deactivate insulin (46,47). Additionally, it has been shown that garlic extract increases the amount of liver glycogen storage, stimulates insulin and Glut-4 expression, absorbs carbs, and lowers blood glucose levels (48). There hasn't been enough research done on the mechanism by which garlic decreases plasma lipids. However, studies on animals have shown that taking supplements containing garlic lowers the activity of hepatic lipogenic and cholesterogenic enzymes (49). Sterol Regulatory Element Binding Protein-1c (SREBP-1c), an essential transcription factor in lipid metabolism, is activated and blood lipid levels are raised by excessive fat ingestion. SREBP-1c suppresses the expression of lipolytic genes, such as Carnitine

Palmitoyltransferase-1 (CPT-1) and Peroxisome Proliferator-Activated Receptor Alpha (PPAR $\alpha$ ), and activates a multitude of lipogenic genes involved in the synthesis of fatty acids, triglycerides, and cholesterol (3-Hydroxy-3-Methylglutaryl-CoA Reductase - HMG-CoA reductase and Acyl-CoA: Cholesterol Acyltransferase - ACAT)(50,51). Key adipogenic transcription factor peroxisome proliferator activated receptor  $\gamma$  (PPAR $\gamma$ ) increases the transcription of genes involved in glucose and fatty acid absorption (52). Research has revealed that mice fed a high-fat diet (HFD) have elevated levels of PPAR $\gamma$ 2 mRNA, which may indicate that HFD-induced adipocyte hypertrophy is related to it. Studies have shown that supplementing with garlic significantly reduces the expression of genes that accumulate fat, such as PPAR $\gamma$ , ACC, and adipocyte Protein 2 (aP2) mRNA (53,54). The expression of lipogenic enzymes, including ACC and glucose-6-phosphate dehydrogenase (G6PDH), was likewise decreased by garlic administration, indicating a possible function in preventing lipogenesis and thereby enhancing fatty acid oxidation. The decrease in plasma triglyceride and cholesterol levels that has been reported may be attributed to this mechanism, underscoring the medicinal potential of garlic in the management of hyperlipidemia (54).

A specific investigation found that giving rats black garlic extract reduced the expression of liver SREBP-1c and the target genes it is linked to (FAS, ACC, and G6PDH). As a consequence, the liver's ability to synthesize fat was decreased due to this downregulation. Moreover, a noteworthy reduction in the expression of ACAT and HMG-CoA reductase was seen in rats given black garlic supplements, which helped to significantly lower the blood cholesterol levels (55). A different investigation on the garlic component diallyl disulfide (DADS) showed a reduction in the mRNA expression of SREBP-1c, ACC, FAS, and HMG-CoA reductase. Concurrently, there was a rise in PPAR $\alpha$  and CPT-1 mRNA expression. These results show that in rats given a high-fat diet (HFD), DADS

substantially decreased the production of hepatic fatty acids and cholesterol while modulating lipid metabolism (56).

## Conclusion

The findings of this study demonstrate that the compounds found in Garlic have a beneficial impact on lowering blood glucose and cholesterol in both animal and human subjects. Garlic components enhance blood glucose control by promoting the release of insulin and the absorption of glucose. Moreover, the protection it provides against changes in lipid metabolism mostly occurs through the SREBP-1c dependent route. Garlic components have no significant adverse effects, making it suitable for use as an herbal remedy to enhance glucose and lipid metabolism, particularly in those with diabetes and dyslipidemia. Garlic and its bioactive components show great potential as dietary supplements for preventing and treating many ailments. Further investigation is required to uncover the intricacies of the processes by which Garlic operates and its molecular pathways. Furthermore, it is imperative to conduct additional clinical trials in order to

substantiate the physiological advantages of Garlic in the human population.

## Acknowledgments

We would like to express our gratitude to those who have helped us in Clinical Biochemistry Research Center of Shiraz University of Medical Sciences

## Funding

This research received no specific grant from funding agencies in the public, commercial, or not-for-profit sectors

## Conflict of Interest

The authors have stated no conflict of interest.

## Authors' contributions

Conceptualization: F.S, A.SR, F.Z and S.D. Data gathering and curation: N.H and M.S. Writing original draft preparation: F.S, A.SR, N.H, and N.N. Writing review and final editing: A.SR, S.D and M.AA. Visualization: M.S. Supervision: A.SR and M.AA. All authors have read and agreed to the published version of the manuscript.

## References

1. Babel RA, Dandekar MP. A review on cellular and molecular mechanisms linked to the development of diabetes complications. *Current Diabetes Reviews*. 2021;17(4):457-73.
2. Matsuzaka T, Shimano H. New perspective on type 2 diabetes, dyslipidemia and non-alcoholic fatty liver disease. *Journal of diabetes investigation*. 2020;11(3):532-4.
3. Vekic J, Zeljkovic A, Stefanovic A, Jelic-Ivanovic Z, Spasojevic-Kalimanovska V. Obesity and dyslipidemia. *Metabolism*. 2019;92:71-81.
4. Hermans MP, Valensi P. Elevated triglycerides and low high-density lipoprotein cholesterol level as marker of very high risk in type 2 diabetes. *Current Opinion in Endocrinology, Diabetes and Obesity*. 2018;25(2):118-29.
5. Sone H, Tanaka S, Tanaka S, Iimuro S, Oida K, Yamasaki Y, et al. Serum level of triglycerides is a potent risk factor comparable to LDL cholesterol for coronary heart disease in Japanese patients with type 2 diabetes: subanalysis of the Japan Diabetes Complications Study (JDCS). *The Journal of Clinical Endocrinology & Metabolism*. 2011;96(11):3448-56.
6. Alloubani A, Nimer R, Samara R. Relationship between hyperlipidemia, cardiovascular disease and stroke: a systematic review. *Current Cardiology Reviews*. 2021;17(6):52-66.
7. Birajdar Z. Effect of Cassia Glauca Lamk Leaves Extract on Cardiovascular and Renal Complications Complicated with Streptozotocin-Induced Diabetic Rats: Rajiv Gandhi University of Health Sciences (India). *International Journal of Pharma Sciences and Research (IJPSR)*. 2015; 6(2):322-33.
8. Asadi-Samani M, Moradi MT, Mahmoodnia L, Alaei S, Asadi-Samani F, Luther T. Traditional uses of medicinal plants to prevent and treat diabetes; an updated review of ethnobotanical studies in Iran. *Journal of nephropathology*. 2017;6(3):118.
9. Sanie-Jahromi F, Zia Z, Afarid M. A review on the effect of garlic on diabetes, BDNF, and VEGF as a potential treatment for diabetic retinopathy. *Chinese medicine*. 2023;18(1):18.

10. Suleria HA, Butt MS, Khalid N, Sultan S, Raza A, Aleem M, et al. Garlic (*Allium sativum*): diet based therapy of 21st century—a review. *Asian Pacific journal of tropical disease*. 2015;5(4):271-8.
11. Foroozand R, Sabokroo M, Firouzi A, Mahalik G, Bouyahya A, Amiri-Ardekani E. Clinical Trials on Role of Garlic (*Allium sativum*) in Managing Diabetes: Systematic Review. *Systematic Reviews in Pharmacy*. 2023;14(2):77-81.
12. Wilson A, Pandya D, Mankad A, Solanki H. A Review on *Allium sativum* (L.) as medicinal plant. *Int. J. Res. Anal. Rev. (IJRAR)*. 2019;6:140-7.
13. Bazaraliyeva A, Moldashov D, Turgumbayeva A, Kartbayeva E, Kalykova A, Sarsenova L, et al. Chemical and biological properties of bio-active compounds from garlic (*Allium sativum*). *Pharmacia*. 2022;69(4):955-64.
14. Lawson LD, Hunsaker SM. Allicin bioavailability and bioequivalence from garlic supplements and garlic foods. *Nutrients*. 2018;10(7):812.
15. Farhat Z, Hershberger PA, Freudenheim JL, Mammen MJ, Hageman Blair R, Aga DS, et al. Types of garlic and their anticancer and antioxidant activity: A review of the epidemiologic and experimental evidence. *European Journal of Nutrition*. 2021;60(7):3585-3609.
16. Ozma MA, Abbasi A, Ahangarzadeh Rezaee M, Hosseini H, Hosseinzadeh N, Sabahi S, et al. A critical review on the nutritional and medicinal profiles of garlic's (*Allium sativum* L.) bioactive compounds. *Food Reviews International*. 2023;39(9):6324-61.
17. Salehi B, Zucca P, Orhan IE, Azzini E, Adetunji CO, Mohammed SA, et al. Allicin and health: A comprehensive review. *Trends in Food Science & Technology*. 2019;86:502-16.
18. Reiter J, Hübbers AM, Albrecht F, Leichert LIO, Slusarenko AJ. Allicin, a natural antimicrobial defence substance from garlic, inhibits DNA gyrase activity in bacteria. *International Journal of Medical Microbiology*. 2020;310(1):151359.
19. Sánchez-Gloria JL, Arellano-Buendía AS, Juárez-Rojas JG, García-Arroyo FE, Argüello-García R, Sánchez-Muñoz F, et al. Cellular mechanisms underlying the cardioprotective role of allicin on cardiovascular diseases. *International journal of molecular sciences*. 2022;23(16):9082.
20. Maitisha G, Aimaiti M, An Z, Li X. Allicin induces cell cycle arrest and apoptosis of breast cancer cells in vitro via modulating the p53 pathway. *Molecular Biology Reports*. 2021;48(11):7261-72.
21. Ezeorba TP, Chukwudozie KI, Ezema CA, Anaduaka EG, Nweze EJ, Okeke ES. Potentials for health and therapeutic benefits of garlic essential oils: Recent findings and future prospects. *Pharmacological Research-Modern Chinese Medicine*. 2022;3:100075.
22. El-Saber Batiha G, Magdy Beshbishy A, G. Wasef L, Elewa YH, A. Al-Sagan A, Abd El-Hack ME, et al. Chemical constituents and pharmacological activities of garlic (*Allium sativum* L.): A review. *Nutrients*. 2020;12(3):872.
23. Pangastuti A, Indriwati SE, Amin M. Investigation of the anti-aging properties of allicin from *Allium sativum* L bulb extracts by a reverse docking approach. *Tropical Journal of Pharmaceutical Research*. 2018;17(4):635-9.
24. Mohamed GA, Ibrahim SR, Elkhayat ES, El Dine RS. Natural anti-obesity agents. *Bulletin of faculty of pharmacy, Cairo University*. 2014;52(2):269-84.
25. Zhang Y, Xu L, Ding M, Su G, Zhao Y. Anti-obesity effect of garlic oil on obese rats via Shenque point administration. *Journal of ethnopharmacology*. 2019;231:486-93.
26. Mahdavi-Roshan M, Nasrollahzadeh J, Zadeh AM, Zahedmehr A. Does garlic supplementation control blood pressure in patients with severe coronary artery disease? A clinical trial study. *Iranian Red Crescent Medical Journal*. 2016;18(11):e23871.
27. Hussein HJ, Hameed IH, Hadi MY. A Review: Anti-microbial, Anti-inflammatory effect and Cardiovascular effects of Garlic: *Allium sativum*. *Research Journal of Pharmacy and Technology*. 2017;10(11):4069-78.
28. Shabani E, Sayemiri K, Mohammadpour M. The effect of garlic on lipid profile and glucose parameters in diabetic patients: A systematic review and meta-analysis. *Primary care diabetes*. 2019;13(1):28-42.
29. Zhu Y, Anand R, Geng X, Ding Y. A mini review: garlic extract and vascular diseases. *Neurological research*. 2018;40(6):421-5.
30. Kumar R, Chhatwal S, Arora S, Sharma S, Singh J, Singh N, et al. Antihyperglycemic, antihyperlipidemic, anti-inflammatory and adenosine deaminase-lowering effects of garlic in patients with type 2 diabetes mellitus with obesity. *Diabetes, metabolic syndrome and obesity: targets and therapy*. 2013:49-56.
31. Ansari MA, Arain AA, Phull QZ, Memon AR. Effects of S-allyl cysteine on insulin secretion: a proposed mechanism for its anti-hyperglycemic effects. *Biomedical Journal Scientific & Technical Research*. 2018;6(3):1-3.
32. Li M, Yun W, Wang G, Li A, Gao J, He Q. Roles and mechanisms of garlic and its extracts on atherosclerosis: A review. *Frontiers in Pharmacology*. 2022;13:954938.
33. Shaikh K, Kinninger A, Cherukuri L, Birudaraju D, Nakanishi R, Almeida S, et al. Aged garlic extract reduces low attenuation plaque in coronary arteries of patients with diabetes: a randomized, double-blind, placebo-controlled study. *Experimental and therapeutic medicine*. 2020;19(2):1457-61.

34. Thomson M, Al-Qattan KK, Bordia T, Ali M. Including garlic in the diet may help lower blood glucose, cholesterol, and triglycerides. *The Journal of nutrition*. 2006;136(3):800S-2S.
35. Prihanti GS, Isnaini F, Yudistia R, Faradilla A, Rahman M. Effect of black garlic extract on blood glucose, lipid profile, and sgpt-sgot of wistar rats diabetes mellitus model. *Majalah Kedokteran Bandung*. 2019;51(2):82-7.
36. Padiya R, Khatua TN, Bagul PK, Kuncha M, Banerjee SK. Garlic improves insulin sensitivity and associated metabolic syndromes in fructose fed rats. *Nutrition & metabolism*. 2011;8:53.
37. Supakul L, Pintana H, Apaijai N, Chattipakorn S, Shinlapawittayatorn K, Chattipakorn N. Protective effects of garlic extract on cardiac function, heart rate variability, and cardiac mitochondria in obese insulin-resistant rats. *European journal of nutrition*. 2014;53:919-28.
38. Sohn CW, Kim H, You BR, Kim MJ, Kim HJ, Lee JY, et al. High temperature-and high pressure-processed garlic improves lipid profiles in rats fed high cholesterol diets. *Journal of Medicinal Food*. 2012;15(5):435-40.
39. Ebadi A, RahimiLanji E, Taghdosi M, Khorshidi A, Akbari H. Effect of garlic tablet on blood glucose in type 2 diabetic patients. *Feyz Medical Sciences Journal*. 2007;11(1):20-5.(in Persian)
40. Ashraf R, Khan RA, Ashraf I. Garlic (*Allium sativum*) supplementation with standard antidiabetic agent provides better diabetic control in type 2 diabetes patients. *Pakistan Journal of Pharmaceutical Sciences*. 2011;24(4):565-70.
41. Faroughi F, Charandabi S, Javadzadeh Y, Mirghafourvand M. Effects of garlic pill on blood glucose level in borderline gestational diabetes mellitus: a randomized controlled trial. *Iranian Red Crescent Medical Journal*. 2018;20(5):1-9.
42. Afkhami-Ardekani M, Kamali-Ardekani A, Shojaoddiny-Ardekani A. Effects of garlic on serum lipids and blood glucose of type 2 diabetic patients. *International Journal of Diabetes in Developing Countries*. 2006;26(2):86-8.
43. Sobenin IA, Nedosugova LV, Filatova LV, Balabolkin MI, Gorchakova TV, Orekhov AN. Metabolic effects of time-released garlic powder tablets in type 2 diabetes mellitus: the results of double-blinded placebo-controlled study. *Acta Diabetologica*. 2008;45(11):1-6.
44. Sangouni AA, Azar MRMH, Alizadeh M. Effect of garlic powder supplementation on hepatic steatosis, liver enzymes and lipid profile in patients with non-alcoholic fatty liver disease: a double-blind randomised controlled clinical trial. *British Journal of Nutrition*. 2020;124(4):450-6.
45. Saikat ASM, Hossain R, Mina FB, Das S, Khan IN, Mubarak MS, et al. Antidiabetic effect of garlic. *Revista Brasileira de Farmacognosia*. 2021;32(1):1-11.
46. Trio PZ, You S, He X, He J, Sakao K, Hou D-X. Chemopreventive functions and molecular mechanisms of garlic organosulfur compounds. *Food & function*. 2014;5(5):833-44.
47. Obih P, Obih J-C, Arome O. Is alpha-glucosidase inhibition a mechanism of the antidiabetic action of garlic (*Allium sativum*)? *Journal of Biosciences and Medicines*. 2019;7(10):42-9.
48. Bhattacharya S, Maji U, Khan GA, Das R, Sinha AK, Ghosh C, et al. Antidiabetic role of a novel protein from garlic via NO in expression of Glut-4/insulin in liver of alloxan induced diabetic mice. *Biomedicine & Pharmacotherapy*. 2019;111:1302-14.
49. Zeng T, Zhang C-L, Zhao X-L, Xie K-Q. The roles of garlic on the lipid parameters: a systematic review of the literature. *Critical reviews in food science and nutrition*. 2013;53(3):215-30.
50. Moon Y-A. The SCAP/SREBP pathway: a mediator of hepatic steatosis. *Endocrinology and Metabolism*. 2017;32(1):6-10.
51. Serviddio G, Bellanti F, Vendemiale G. Free radical biology for medicine: learning from nonalcoholic fatty liver disease. *Free Radical Biology and Medicine*. 2013;65:952-68.
52. Wang J, Zhou M, Wu T, Fang L, Liu C, Min W. Novel anti-obesity peptide (RLLPH) derived from hazelnut (*Corylus heterophylla* Fisch) protein hydrolysates inhibits adipogenesis in 3T3-L1 adipocytes by regulating adipogenic transcription factors and adenosine monophosphate-activated protein kinase (AMPK) activation. *Journal of Bioscience and bioengineering*. 2020;129(3):259-68.
53. Hsu W-TC, Lim JL. Inhibitory Effect of Methanolic Extract of Black Garlic on Adipogenesis in 3T3-L1 Adipocytes. *Zhongshan Medical Journal*. 2018;29:P113-21.
54. Kim M, Kim H. Effect of garlic on high fat induced obesity. *Acta Biologica Hungarica*. 2011;62(3):244-54.
55. Ha AW, Ying T, Kim WK. The effects of black garlic (*Allium sativum*) extracts on lipid metabolism in rats fed a high fat diet. *Nutrition research and practice*. 2015;9(1):30-6.
56. Lai Y-S, Chen W-C, Ho C-T, Lu K-H, Lin S-H, Tseng H-C, et al. Garlic essential oil protects against obesity-triggered nonalcoholic fatty liver disease through modulation of lipid metabolism and oxidative stress. *Journal of agricultural and food chemistry*. 2014;62(25):5897-906.
57. Ozougwu J, Eyo J. Studies on the anti-diabetic activity of *Allium sativum* (garlic) aqueous extracts on alloxan-induced diabetic albino rat. *Pharmacologyonline*. 2010;2:1079-88.



58. Yang C, Li L, Yang L, Lü H, Wang S, Sun G. Anti-obesity and Hypolipidemic effects of garlic oil and onion oil in rats fed a high-fat diet. *Nutrition & metabolism*. 2018;15(1):1-8.
59. Amor S, González-Hedström D, Martín-Carro B, Inarejos-García AM, Almodóvar P, Prodanov M, et al. Beneficial effects of an aged black garlic extract in the metabolic and vascular alterations induced by a high fat/sucrose diet in male rats. *Nutrients*. 2019;11(1):153.
60. Szulińska M, Kręgielska-Narozna M, Świątek J, Styś P, Kuźnar-Kamińska B, Jakubowski H, et al. Garlic extract favorably modifies markers of endothelial function in obese patients—randomized double blind placebo-controlled nutritional intervention. *Biomedicine & Pharmacotherapy*. 2018;102:792-7.
61. Ashraf R, Aamir K, Shaikh AR, Ahmed T. Effects of garlic on dyslipidemia in patients with type 2 diabetes mellitus. *Journal of Ayub Medical College, Abbottabad: JAMC*. 2005;17(3):60-4.