



Tehran University of Medical
Sciences Publication
<https://tums.ac.ir>

Iran J Parasitol

Open access Journal at
<https://ijpa.tums.ac.ir>



Iranian Society of Parasitology
<https://isp.tums.ac.ir>

Original Article

Synergistic Effects of Combined Caryophyllene and Albendazole against Invaginated and Evaginated Protoscoleces of *Echinococcus granulosus*

Maryam Khodadadi ¹, Mohammad Javad Pour Maram ², Hossein Amini Khoei ³,
Hadi Raeisi Shahraki ⁴, *Ebrahim Saedi Dezaki ³

1. School of Medicine, Shahrekord University of Medical Sciences, Shahrekord, Iran

2. Department of Parasitology, School of Medicine, Shahrekord University of Medical Sciences, Shahrekord, Iran

3. Medical Plants Research Center, Basic Health Sciences Institute, Shahrekord University of Medical Sciences, Shahrekord, Iran

4. Department of Biostatistics, Shahrekord University of Medical Sciences, Shahrekord, Iran

Received 10 Feb 2026
Accepted 15 May 2026

Keywords:
Natural products;
Scolicidal;
Apoptosis;
Protoscoleces;
Echinococcus granulosus

***Correspondence**
Email:
saedi1358@gmail.com

Abstract

Background: Natural compounds have garnered increasing attention as candidate antiparasitic therapeutics. We aimed to evaluate the protoscolicidal efficacy of caryophyllene (CAP), alone and in combination with albendazole (ALZ), against hydatid cyst protoscoleces.

Methods: Suspensions of 10,000 invaginated and evaginated protoscoleces (400 μ L normal saline) were exposed to CAP (62.5–500 μ g/mL), ALZ (250 μ g/mL), or CAP+ALZ combinations in 24-well culture plates. Protoscoleces viability was assessed by eosin exclusion test. Reactive oxygen species (ROS) induction was quantified using the DCFH-DA assay, and pre-apoptotic gene expression was analyzed by real-time PCR.

Results: CAP, particularly in combination with ALZ, significantly reduced the viability of both invaginated and evaginated protoscoleces in a concentration- and time-dependent manner. Combination therapy achieved 100% mortality of invaginated protoscoleces within 10–30 minutes, demonstrating significant synergism over either monotherapy. Evaginated protoscoleces viability was also markedly diminished. The observed scolicidal activity was mechanistically correlated with apoptosis induction, evidenced by elevated ROS production and significant upregulation of caspase-3 and P53 gene expression ($P<0.001$).

Conclusion: CAP, particularly in combination with ALZ, exhibits potent *in vitro* protoscolicidal activity against hydatid cyst protoscoleces, with apoptosis induction and ROS generation implicated as underlying mechanisms. Further pharmacodynamic investigations and clinical trials are warranted to elucidate the precise mechanisms of action and evaluate the therapeutic potential of CAP-based combination therapy in the management of cystic echinococcosis.



Copyright © 2026 Khodadadi et al. Published by Tehran University of Medical Sciences.

This work is licensed under a Creative Commons Attribution-NonCommercial 4.0 International license.

(<https://creativecommons.org/licenses/by-nc/4.0/>). Non-commercial uses of the work are permitted, provided the original work is properly cited
<https://doi.org/10.18502/ijpa.v21i2.22078>

Introduction

Hydatid disease, recognized as a "neglected parasitic infection," is caused by the larval stage of the cestode *Echinococcus granulosus* (1). This zoonotic infection is transmitted from animals to humans and is prevalent in numerous regions worldwide, particularly in areas with extensive livestock farming (1). Human infection occurs incidentally through the ingestion of water or vegetables contaminated with *E. granulosus* eggs shed by dogs, or via direct contact with infected dogs (2). Although hydatid cysts are typically asymptomatic during the initial stages, progressive cyst growth exerts mechanical pressure on the involved organ, leading to complications (3). The disease severity and clinical presentation depend on the organ affected, the infection intensity, and the organ's susceptibility to cyst development (3).

Pharmacological treatment with benzimidazoles, such as albendazole or mebendazole, is advised for patients who are not candidates for surgery, those with cysts in multiple organs, and for postoperative management (4). Surgical intervention, while a primary treatment modality, is complicated by frequent recurrences due to the potential spillage of viable protoscoleces during cyst excision, underscoring the complexity of hydatid cyst surgery (4). To mitigate recurrence risk, surgeons administer scolical agents into the cyst cavity intraoperatively. Various chemical scolicides have been employed, including 2% formalin, 20%-30% hypertonic saline, 0.54% silver nitrate, and betadine solutions (5). However, the use of these agents is constrained by their associated adverse effects, such as hepatic and biliary fibrosis, allergic and hypersensitivity reactions, and damage or necrosis of adjacent tissues.

Now, compounds derived from natural supplies are acknowledged for their promising potential as applicants for antiparasitic therapeutics. Likewise, phytomedicines are progressively recognized as economical and safe alternatives to conventional treatment modalities (6). Caryophyllene ($C_{15}H_{24}$, CAP) is a

prominent natural compound within the terpene class, particularly classified as a sesquiterpene, and is commonly present in the essential oils of various plants including cloves, cinnamon, rosemary, and species within the cannabis genus (6). Its distinctive chemical structure, notably the incorporation of a cyclobutane ring, underpins a range of biological activities such as anti-inflammatory, analgesic, antioxidant, and antimicrobial effects (7). In addition, CAP-rich herbs displayed efficacy in inhibiting the growth and proliferation of parasites like *Leishmania* spp., and *Trypanosoma* spp. (8). The proposed mechanisms underlying these effects involve disruption of parasite cell membranes, impairment of cellular energy production via mitochondrial inhibition, and enhancement of host immune responses targeting the parasites (9). Additionally, the antioxidant capacity of caryophyllene may contribute to mitigating oxidative stress-induced damage commonly associated with parasitic infections (10).

We aimed to evaluate the protoscolicidal effects of caryophyllene (CAP) alone and along with albendazole (ALZ) against protoscoleces of hydatid cysts.

Materials and Methods

Ethics

This work was approved by the ethical committee of the Shahrekord University of Medical Sciences, Shahrekord, Iran (No. IR.SKUMS.REC.1403.019).

Chemicals

Caryophyllene (≥ 98.0 purity), albendazole (≥ 98.0 purity), and eosin dye were purchased from Sigma-Aldrich, Germany.

Collecting the protoscoleces

Sheep livers harboring hydatid cysts were obtained from Junaqan slaughterhouse (Iran) and transported to the Parasitology Laboratory,

Shahrekord Medical School, Iran. Large, unilocular cysts containing adequate protoscoleces were selected for analysis. Under sterile conditions, cyst surfaces were decontaminated sequentially with 70% ethanol and betadine solution. Cyst fluid was aspirated via syringe into 50 mL Falcon tubes; cysts were then incised with a sterile scalpel, and protoscoleces were aspirated, sedimented, and washed three times with phosphate-buffered saline (PBS) (11).

Viability examination

Protoscolex viability was assessed by mixing a drop of sediment with 0.1% eosin solution (Sigma-Aldrich, Germany) and examining under a 10× objective; protoscoleces absorbing the stain (orange-to-pink coloration) were classified as non-viable (Fig. 1). Only cysts with >90% viability, determined by evaluating 100 protoscoleces, were selected for further analysis. The protoscolex suspension was subsequently standardized to 2×10^3 /mL (12).




Fig. 1: Hydatid cyst protoscoleces stained with eosin dye (present study). Viable and non-viable protoscoleces are indicated by white and black arrow, respectively.

Antiparasitic Efficacy on invaginated protoscoleces

Following previously established protocols (13), suspensions of 10,000 protoscoleces in 400 μ L normal saline were exposed to CAP (62.5–500 μ g/mL), ALZ (250 μ g/mL), or

CAP+ALZ combinations in 24-well culture plates. Viability was assessed at 5–60 min intervals by eosin exclusion test: protoscoleces absorbing 0.01% eosin solution with orange-to-pink uniform staining of cellular contents were classified as non-viable. All experiments were performed in triplicate, and IC₅₀ values were calculated using GraphPad Prism v8.4.3.

Antiparasitic Efficacy on evaginated protoscoleces

Biphasic culture

The biphasic culture medium consisted of a liquid phase comprising CMRL 1066 (Gibco), yeast extract, penicillin/streptomycin (Biosera), and 0.5% dog bile in PBS, and a solid phase of fetal bovine serum (Gibco). Approximately 10,000 protoscoleces (100 μ L sediment) were initially transferred into CMRL 1066 supplemented with FBS and dog bile for 48-hour acclimatization, then transferred to solid-phase flasks with ~45 mL liquid medium and incubated at 37°C in a CO₂ incubator. To prevent aggregation, flasks were gently agitated every 4–6 hours, followed by continuous shaking via an in-incubator device. The liquid phase was renewed every 3–5 days, guided by protoscolex density, developmental stage, and medium color change. Only evaginated protoscoleces were employed for pharmacological assessment (22).

Effects on the viability of evaginated protoscoleces

Protoscolex viability was monitored daily by inverted microscopy throughout culture. Following 48-hour incubation, evaginated protoscoleces were exposed to CN concentrations of 25–200 μ g/mL, with viability assessed at 5–60 minute intervals. All experiments were performed in triplicate, and IC₅₀ values were determined using GraphPad Prism v8.4.3.

Inducing reactive oxygen species (ROS) generation

The induction of ROS by CAP alone and in combination with ALZ was evaluated employing the dichlorodihydrofluorescein diacetate (DCFH-DA) assay. Specifically, protoscoleces were incubated with 10 μ M DCFH-DA (Sigma-Aldrich, Germany) at 37°C for one hour. Following this incubation, the protoscoleces were exposed to CAP alone and in combination with ALZ for 180 minutes. Fluorescence intensity, corresponding to ROS levels, was measured at excitation/emission wavelengths of 488/525 nm over a four-hour timeframe using a microplate reader (14).

Expression level of the pre-apoptotic genes

mRNA extraction

Total RNA was extracted from CAP-, ALZ-, and combination-treated protoscoleces using the SinaPure™ RNA kit (SinaClon, Iran),

followed by DNase I treatment to eliminate DNA contamination. RNA quantity and integrity were assessed by Nanodrop spectrophotometry and agarose gel electrophoresis, respectively.

Complementary DNA (cDNA) synthesis

cDNA synthesis was performed using the Yekta Tehiz kit (Iran) per manufacturer's instructions, with successful synthesis confirmed by target gene amplification and gel visualization; cDNA was stored at -80°C until use.

Primer design

Primers for caspase-3, P53, and β -actin were designed based on Ensembl sequence data using Gene Runner software, with secondary structure and primer-dimer analysis conducted via Oligo Analyzer, in silico PCR validation using the UCSC tool, and specificity confirmed by NCBI Primer-BLAST (Table 1) (15, 16).

Table 1: Primer sequences utilized for real-time polymerase chain reaction (PCR) analysis

Primer	Sequence (5'–3')
P53	F: AACCACCGAACTCACAAC R: AACCGACACAACATCATCA
Caspase-3	F: TGGTTCATCCAGTCGCTTTG R: CATTCTGTTGCCACCTTTTCG
β-actin	F: GTGACGTGTGACATCCGTAAAGA R: GCCGGACTCATCGTACTCC

qRT-PCR analysis

qRT-PCR was performed on a Rotor-Gene system (Qiagen, Germany) using SYBR Green master mix (Yekta Tajhiz Azma) in a 25 μ L reaction containing 7 μ L master mix, 1 μ L each primer (10 pM), 3 μ L cDNA, and 13 μ L sterile water. Thermal cycling comprised initial denaturation at 96°C for 10 min, followed by 39 cycles of 96°C denaturation, 61°C annealing (30 sec), and 73°C extension (30 sec). Relative gene expression was quantified using the $2^{-\Delta\Delta C_t}$ method with β -actin as the reference gene (15, 16).

Statistical analysis

Following data collection, the dataset was input into the statistical software SPSS version 26 (IBM Corp., Armonk, NY, USA) for analysis, primarily utilizing one-way ANOVA. Additionally, post hoc tests

were conducted to evaluate comparative effects among groups. The findings were presented through relevant tables and statistical graphs. Where correlation analyses were warranted, suitable statistical tests, including the chi-square test with a significance threshold of $P < 0.05$, were employed.

Results

In vitro Scolicidal Activity Against Invaginated Protoscoleces

CAP significantly reduced invaginated protoscolex viability in a concentration- and time-dependent manner (Fig. 2). At 60 min, mortality rates of 39.7%, 46.8%, 78.9%, and 100% were recorded at 62.5, 125, 250, and 500 μ g/mL, respectively, with IC₅₀ values of >500, >500, 370.9, 260.9, and 181.4 μ g/mL at 5, 10, 20, 30, and 60

min, respectively. CAP+ALZ (250 µg/mL) combination therapy achieved 100% mortality at 62.5, 125, 250, and 500 µg/mL after 30, 20, 10, and 10

min, respectively, with IC₅₀ values of 171.4, 75.9, 34.8, and <30 µg/mL at 5, 10, 20, 30, and 60 min, respectively.




Fig. 2: The effect of caryophyllene (CAP) and albendazole (ALZ) alone (A) and in combination (B) on the viability of hydatid cyst invaginated protoscoleces was evaluated using the eosin exclusion test. Results are presented as mean $\pm$ standard deviation (N=3). Statistical significance was determined via one-way ANOVA with Tukey's post-hoc analysis

In vitro Scolicidal Activity Against Evaginated Protoscoleces

Microscopic analysis revealed that CAP reduced evaginated protoscolex viability in a dose- and time-dependent manner (Fig. 3). After 60 min, mortality rates of 30.2%, 41.3%, 59.6%, and 100% were recorded at 62.5, 125, 250, and 500 µg/mL, respectively, with IC₅₀ values of >500, >500, 486.4, 376.5, and

209.7 µg/mL at 5, 10, 20, 30, and 60 min, respectively. CAP+ALZ (250 µg/mL) combination therapy markedly enhanced scolicidal efficacy, achieving mortality rates of 38.6%, 66.6%, 79.6%, and 100% at 62.5, 125, 250, and 500 µg/mL CAP after only 10 min of exposure, with corresponding IC₅₀ values of 207.9, 93.8, 37.4, 65.3, and 33.7 µg/mL at 5, 10, 20, 30, and 60 min, respectively.




Fig. 3: The effect of caryophyllene (CAP) and albendazole (ALZ) alone (A) and in combination (B) on the viability of hydatid cyst evaginated protoscoleces was evaluated using the eosin exclusion test. Results are presented as mean $\pm$ standard deviation (N=3). Statistical significance was determined via one-way ANOVA with Tukey's post-hoc analysis

Inducing ROS generation

The findings from the DCFH-DA assay demonstrated that the mean fluorescence intensities were 7.3, 13.9, 25.3, and 29.3 arbitrary units following exposure of protoscoleces to CAP at half the IC₅₀ concentration, CAP at the IC₅₀ concentration, a combination of ALZ and CAP at half the IC₅₀ concentration, and a combination of ALZ and CAP at the IC₅₀ concentration, respectively. ANOVA with Tukey's post-hoc analysis revealed that these increases in fluorescence intensity were statistically significant ($P<0.001$) when compared to the control group treated with normal saline (Fig. 4).




Fig. 4: The impact of caryophyllene (CAP) alone, as well as in combination with albendazole (ALZ, 250 µg/mL), on the induction of reactive oxygen species (ROS) in hydatid cyst protoscoleces was evaluated using the dichlorodihydrofluorescein diacetate (DCFH-DA) assay. Results are presented as mean $\pm$ standard deviation (N=3). * $P<0.05$; ** $P<0.01$; and *** $P<0.001$ compared to normal saline. ϕ $P<0.05$ compared to AL; Statistical significance was determined via one-way ANOVA with Tukey's post-hoc analysis

Expression level of the pre-apoptotic genes

Real-time PCR analysis demonstrated that following exposure of protoscoleces to CAP and the combination of CAP+ALZ, the expression levels of the *caspase-3* and *P53* genes were significantly ($P<0.001$) elevated (Fig. 5). ANOVA followed by Tukey's post-hoc test

revealed that the expression levels of the *caspase-3* gene were significantly ($P<0.05$) up-regulated in protoscoleces exposed to CAP at half the IC₅₀ concentration, CAP at the IC₅₀ concentration, the combination of ALZ and CAP at half the IC₅₀ concentration, and the combination of ALZ and CAP at the IC₅₀ concentration. Specifically, the fold changes in gene expression were -1.78, -2.12, -3.11, and -4.32, respectively, relative to the control group. Conversely, the expression levels of the *P53* gene after exposure to CAP at 1/2 IC₅₀, CAP at IC₅₀, ALZ+CAP at 1/2 IC₅₀, and ALZ+CAP at IC₅₀ were 1.56, -1.98, -4.11, and -5.39-fold changes, respectively, compared to the control group.




Fig. 5: The impact of caryophyllene (CAP) alone, as well as in combination with albendazole (ALZ, 250 µg/mL), on the expression of pre-apoptotic genes in hydatid cyst protoscoleces was evaluated using Real-time PCR. Results are presented as mean $\pm$ standard deviation (N=3). * $P<0.05$; ** $P<0.01$; and *** $P<0.001$ compared to normal saline. ϕ $P<0.05$; $\phi\phi$ $P<0.001$ compared to ALZ. Statistical significance was determined via one-way ANOVA with Tukey's post-hoc analysis

Discussion

The present study sought to assess the protoscolicidal efficacy of CAP and to investigate its underlying cellular mechanisms, both independently and in conjunction with ALZ, against hydatid cyst protoscoleces. Our results

showed the potent protoscolicidal effects of CAP mainly combined with ALZ against invaginated and evaginated hydatid cyst protoscoleces. Recent studies have demonstrated that combination therapy involving synthetic pharmaceuticals and natural compounds has garnered significant interest as an innovative strategy for addressing parasitic diseases (17). Natural products, through their enhancing, synergistic, or complementary effects, have the potential to augment the efficacy of antiparasitic agents, thereby enabling dose reduction of synthetic drugs and minimizing associated adverse effects (18). By targeting distinct metabolic pathways or modulating the host immune response, these natural compounds not only enhance therapeutic outcomes but also contribute to the prevention of drug resistance development in parasites (19). Consequently, the integration of chemical drugs with herbal constituents is regarded as a promising approach for the advancement of more effective and sustainable antiparasitic treatments (19).

Regarding the antiparasitic properties of CAP, investigations involving leaf extracts of *Pelargonium quercetorum*, which contain CAP, have demonstrated inhibitory effects against *Leishmania major* parasites; however, these effects were not observed to be dose-dependent (20). Additionally, another study reported that the essential oil derived from *Protium ovatum* leaves, also containing CAP, exhibited notable activity against the trypomastigote forms of *Trypanosoma cruzi* (21). The presence of CAP in certain plant extracts has been associated with their traditional application in the treatment of parasitic infections, suggesting a potential synergistic antiparasitic effect when combined with other bioactive compounds (22). The antimicrobial activity of caryophyllene is primarily attributed to its ability to disrupt the cell membrane, thereby enhancing membrane permeability and inducing the leakage of intracellular constituents. This compromise of membrane integrity constitutes a fundamental mechanism

underlying its antibacterial and antifungal properties (23,24).

Concerning the involvement of ROS in the regulation and death of hydatid cyst protoscoleces, research indicates that ROS play a significant role in eliminating these protoscoleces by inducing cellular damage and death, thereby representing a potential therapeutic target (24). Nonetheless, *Echinococcus granulosus* parasites possess intrinsic antioxidant defense mechanisms that mitigate host-generated ROS, helping to maintain the host-parasite equilibrium (24). When ROS concentrations exceed the parasite's antioxidant capacity, oxidative damage occurs to the parasite's DNA, lipids, and proteins, ultimately resulting in loss of viability and cell death (24). It has been proven that CAP can activate MAPK signaling pathways, leading to an increase in reactive oxygen species (ROS) in eukaryotic cells (25). Our results reported the CAP mainly combined with ALZ markedly induced the ROS generation against hydatid cyst protoscoleces, proposed that CAP primarily showed its protoscolicidal effects by increasing the generation of ROS.

Apoptosis serves a dual function in the inhibition of parasites: it operates as a host defense strategy by eliminating infected cells, thereby restricting the propagation of the parasite (27). The results of the present study indicate that the induction of apoptosis serves as an effective mechanism for the elimination of hydatid cyst protoscoleces. Apoptosis offers several advantages over necrosis, notably the mitigation of inflammation and the avoidance of extensive immune responses (28). These characteristics render apoptosis a viable therapeutic target for managing hydatid cysts (28). In this investigation, molecular markers indicative of apoptosis was significantly upregulated in protoscoleces following treatment with CAP alone or in combination with ALZ. Specifically, the activation of caspases, which are critical apoptotic markers, was prominently observed in treated specimens. These findings align with those reported in prior research. One hypothesized

mechanism underlying apoptosis induction in protoscoleces involves the elevated generation of ROS and consequent mitochondrial damage. This pathway may facilitate rapid and effective cell death, particularly in parasitic cells that lack sophisticated antioxidant defense mechanisms (29). Collectively, the data suggest that apoptosis induction represents a promising and targeted approach for hydatidosis therapy. These insights contribute to the advancement of novel pharmacological agents and non-surgical interventions characterized by enhanced efficacy and reduced adverse effects. Previous studies have demonstrated that ROS generation induced by CAP can initiate apoptosis, a regulated cell death mechanism, as indicated by elevated Annexin V binding, disruption of mitochondrial membrane potential, and activation of caspases (30). Therefore, it is reasonable to propose that the use of CAP can effectively trigger apoptosis, thus facilitating the removal of protoscoleces.

Notwithstanding the encouraging results obtained, this study is subject to certain limitations. Primarily, its exclusively *in vitro* design does not account for the physiological complexities inherent to living organisms, nor does it consider pharmacokinetic factors such as absorption, distribution, metabolism, and excretion (ADME). Additionally, the elevated concentrations of CAP employed in the experiments prompt concerns regarding the therapeutic viability and potential toxicity of the treatment. Therefore, it is imperative to conduct preclinical investigations using *in vivo* models to validate the efficacy, establish a safe dosage range, and thoroughly assess the toxicity profile of this therapeutic regimen.

Conclusion

Recent investigation revealed promising *in vitro* effects of CAP, particularly when combined with ALZ, on invaginated and evaginated hydatid cyst protoscoleces. Furthermore, the study highlighted potential underlying

mechanisms, including the induction of apoptosis and the generation of ROS, in the control of hydatid cyst protoscoleces. Nonetheless, additional research is necessary to elucidate the precise mechanisms of action and to evaluate the efficacy of this approach in clinical trials, which could support the integration of CAP into therapeutic strategies for the treatment and management of cystic echinococcosis.

Declaration of interest

The authors declare that there is no conflict of interest.

Acknowledgements

The authors would like to thank the staff of the Medical Plants Research Center, Shahrekord University of Medical Sciences, Shahrekord, Iran for providing laboratory facilities.

References

1. Gessese AT. Review on epidemiology and public health significance of hydatidosis. *Vet Med Int* 2020;2020(1):8859116.
2. Khalili N, Iranpour P, Khalili N, Haseli S. Hydatid disease: a pictorial review of uncommon locations. *Iran J Med Sci*. 2023;48(2):118-129.
3. Dietrich CF, Douira-Khomsy W, Gharbi H, et al. Cystic echinococcosis, review and illustration of non-hepatic manifestations. *Med Ultrason*. 2020;22(3):319-24.
4. Ün M, Yaman SS, Erbaş O. Hydatid cyst and treatment. *Demiroglu Science University Florence Nightingale Journal of Transplantation*. 2020;5(2):070-8.
5. Cheraghipour K, Rouzbahani AK, Fallahi S, et al. Recent advances in therapeutic strategies against hydatid cysts using nanomaterials: A systematic review. *Letters in Drug Design & Discovery*. 2023 1;20(9):1185-93.
6. Abdelmohsen UR, Balasubramanian S, Oelschlaeger TA, et al. Potential of marine

- natural products against drug-resistant fungal, viral, and parasitic infections. *Lancet Infect Dis.* 2017;17(2):e30-41.
7. Gyrdymova YV, Rubtsova SA. Caryophyllene and caryophyllene oxide: A variety of chemical transformations and biological activities. *Chemical Papers.* 2022;76(1):1-39.
 8. Blake K. Beta-Caryophyllene: A review of current research. *Alternative and Complementary Therapies.* 2021;27(5):222-6.
 9. Francomano F, Caruso A, Barbarossa A, et al. β -Caryophyllene: a sesquiterpene with countless biological properties. *Applied Sci.* 2019;9(24):5420.
 10. Sharma C, M. Al Kaabi J, M. Nurulain S, N. Goyal S, Amjad Kamal M, Ojha S. Polypharmacological properties and therapeutic potential of β -caryophyllene: a dietary phytocannabinoid of pharmaceutical promise. *Curr Pharm Des.* 2016;22(21):3237-64.
 11. Mahmoudvand H, Kheirandish F, Dezaki ES, Shamsaddini S, Harandi MF. Chemical composition, efficacy and safety of *Pistacia vera* (var. Fandoghi) to inactivate protoscoleces during hydatid cyst surgery. *Biomed Pharmacother.* 2016;82:393-8.
 12. Mahmoudvand H, Kheirandish F, Ghasemi Kia M, Tavakoli Kareshk A, Yarahmadi M. Chemical composition, protoscolicidal effects and acute toxicity of *Pistacia atlantica* Desf. fruit extract. *Nat Prod Res.* 2016;30(10):1208-11.
 13. Mahmoudvand H, Asadi A, Harandi MF, Sharififar F, Jahanbakhsh S, Dezaki ES. *In vitro* lethal effects of various extracts of *Nigella sativa* seed on hydatid cyst protoscoleces. *Iran J Basic Med Sci.* 2014;17(12):1001-1006.
 14. Marhoon IA, Alwan MS, Al-Ibrahimi LA. Promising Anthelmintic Effects of Camphene Against Protoscoleces and Hydatid Cysts of *Echinococcus granulosus*. *Acta Parasitol.* 2025;70(3):113.
 15. Lu S, Wen L, Gong Y, Tian C, et al. *In vitro* effects of harmine against *Echinococcus granulosus* protoscoleces by stimulating DNA damage. *Exp Parasitol.* 2021;226-227:108121.
 16. Mahmoudvand H, Khalaf AK, Moghadam EF, Yadegari JG, Adineh A, Jalili A. Investigating the Lethal Effects of Carvone on the Protoscoleces of Hydatid Cyst, *In vitro* and *Ex Vivo*. *Iran J Parasitol.* 2025;20(1):91-9.
 17. Shang X, Dai L, Cao X, Ma Y, Gulnaz I, Miao X, Li X, Yang X. Natural products in antiparasitic drug discovery: advances, opportunities and challenges. *Natural Product Reports.* 2025; 42(9):1419-58.
 18. Boğa M, Vlaia L, Jijie AR, et al. Exploring synergistic interactions between natural compounds and conventional chemotherapeutic drugs in preclinical models of lung cancer. *Pharmaceuticals.* 2024;17(5):598.
 19. Maroufi Y, Hoseini SR, Alavi M. Antiparasitic effect of leaf extract and major metabolites of *Pelargonium quercetorum* Agnew. against *Leishmania major*: *in vitro* and *in silico* studies. *Journal of Applied Biotechnology Reports.* 2022;9(4):817-30.
 20. Estevam EB, Deus IP, Silva VP, et al. *In vitro* antiparasitic activity and chemical composition of the essential oil from *Protium ovatum* leaves (Burceraceae). *Anais da Academia Brasileira de Ciências.* 2017;89(04):3005-13.
 21. Ahmed EA. The Potential Therapeutic Role of Beta-Caryophyllene as a Chemosensitizer and an Inhibitor of Angiogenesis in Cancer. *Molecules.* 2025;30(8):1751.
 22. Bonincontro G, Scuderi SA, Marino A, Simonetti G. Synergistic Effect of Plant Compounds in Combination with Conventional Antimicrobials against Biofilm of *Staphylococcus aureus*, *Pseudomonas aeruginosa*, and *Candida* spp. *Pharmaceuticals.* 2023;16(11):1531.
 23. Lim CR, Ogawa S, Kumari Y. β -Caryophyllene: a comprehensive review of its pharmacological activities, mechanisms of action, and therapeutic potential. *Archives of Pharmacol Research.* 2025;48(1):1-42. doi: 10.1007/s12272-024-01523-z.
 24. Moo CL, Yang SK, Osman MA, et al. Antibacterial activity and mode of action of β -caryophyllene on *Bacillus cereus*. *Pol J Microbiol.* 2020;69(1):1-6.
 25. Cancela M, Paes JA, Moura H, Barr JR, Zaha A, Ferreira HB. Unraveling oxidative stress

- response in the cestode parasite *Echinococcus granulosus*. Sci Rep. 2019;9(1):15876.
26. Fidyk K, Fiedorowicz A, Strzdała L, Szumny A. β -caryophyllene and β -caryophyllene oxide—natural compounds of anticancer and analgesic properties. Cancer Med. 2016;5(10):3007-17.
27. Buret AG, Bhargava A. Modulatory mechanisms of enterocyte apoptosis by viral, bacterial and parasitic pathogens. Crit Rev Microbiol. 2014;40(1):1-17.
28. Barcinski MA, DosReis GA. Apoptosis in parasites and parasite-induced apoptosis in the host immune system: a new approach to parasitic diseases. Braz J Med Biol Res. 1999;32:395-401.
29. Pawłowska M, Mila-Kierzenkowska C, Szczegielniak J, Woźniak A. Oxidative stress in parasitic diseases—Reactive oxygen species as mediators of interactions between the host and the parasites. Antioxidants. 2023;13(1):38.
30. Ullah H, Di Minno A, Santarcangelo C, Khan H, Daglia M. Improvement of oxidative stress and mitochondrial dysfunction by β -caryophyllene: A focus on the nervous system. Antioxidants. 2021;10(4):546.