

Original Article

Relationship between Biological and Qualitative Indices in Surface Waters Receiving the Effluent of Fish Farms in the Northwest of Iran

Farhad Pourfaraj¹, Malek Abazari², *Eslam Moradi-Asl², *S. Amad Mokhtari¹

¹Department of Environment Health, Ardabil University of Medical Sciences, Ardabil, Iran

²Arthropod-Borne Diseases Research Center, Ardabil University of Medical Sciences, Ardabil, Iran

*Corresponding authors: Dr Eslam Moradi-Asl, E-mail: Moradiasl83@yahoo.com, Dr S. Amad Mokhtari, E-mail: s.a.mokhtari@gmail.com

(Received 04 Aug 2020; accepted 01 Jun 2024)

Abstract

Background: Water quality is usually measured using various indicators based on physical, chemical and biological parameters. By using the biological index that is based on the identification of the arthropod families, it is possible to make a logical judgment about the ecosystem condition. The aim of this study was measuring correlation coefficients between qualitative and biological Indices.

Methods: Water samples were collected 27 samples in northwest of Iran and aquatic insects' samples 54 in 2019. The NSFWQI and IRWQISC as the most important indices of physical and chemical quality of water ranged from 54.45–76.21 and from 41.32 to 77.40, respectively.

Results: A total of 2,953 aquatic insects were collected, and biological Index ranged from 6.26 to 3.38. It can be stated that increasing in the concentration of pollutants in the source and end of the river could lead to a sharp decrease in biological index. IRWQISC index, the effluent stations of fish farms can fit into 'fairly bad quality' and 'moderate quality' categories.

Conclusion: The linear regression analysis revealed a significant relationship between the Hilsenhoff biological Index and the physiochemical parameters of pH, DO (Dissolved Oxygen) and total dissolved solids. The activity of fish farms and discharging their effluents into water sources, can change the physical, chemical and biological parameters of receiving waters, therefore it is recommended that the location of these units be reviewed and also the appropriate treatment for such effluents should be considered, so that the health risks caused by them can be effectively reduced.

Keywords: Water quality index; Biological index; Aquatic insects; Water quality; Iran

Introduction

Rivers and running waters have been always regarded as essential resources for human beings throughout history providing them with necessary water for drinking, farming and industrial purposes (1). One of the main reasons for the location and establishment of urban, agricultural and industrial centers is the presence of abundant water resources (2). These resources, however, are under grave threat by consumers (3). Hence, it is necessary to identify the resources of physiochemical and biological pollutants such as BOD (Biological oxygen demand), fecal coliform, nitrate, COD (Chemical

oxygen demand), ammonium, phosphate, turbidity, TDS (Total dissolved solids), and etc. Raise awareness regarding their changing trends and devise effective strategies to hinder the deterioration of water quality and find ways to improve it (4–7). Several countries have provided their special guidelines to monitor the quality of water resources or have tried to follow the standard guidelines suggested by international organizations (8, 9). The water quality index is measured via physical, chemical and biological parameters (10, 11) and is regarded as a criterion for the categorization of waters

based on using standard parameters (12). These Indices are mathematical tools to quantify the descriptive data on the quality level of waters (13, 14). The National Sanitation Foundation Water Quality Index (NSFWQI) and Iran Surface Water Quality Index (IRWQI_{SC}) Indices are two of the most complete and comprehensive water quality Indices (8, 9, 15, 16) used to study waters, monitor and detect pollutants in waters and qualitatively evaluate the zoning of surface waters by measuring physical and chemical features (17, 18). These two Indices are less problematic compared to other models and because of their simplicity and availability, they are frequently employed by researchers around the world (19–22). After coding and measuring the qualitative information, the sub-index amount of each parameter is measured based on the characteristic curve for quality score classification and weighting factor of each parameter listed in the tables (23–25). The above indicators present water quality in simple terms for experts and the public by presenting numbers in certain classifications. Nevertheless, in terms of ecology, one of the best and cheapest scientific methods, compared to chemical methods, is to determine the biological health of waters and determine the effects of human activities on water qualities by biological evaluations and monitoring (26–28). The biological index works based on the detection of arthropods in families (29, 30) and can pronounce logical judgments on ecosystems (31–33). Various national and international studies have been conducted on the investigation of water qualities in line with the two above-mentioned Indices (34, 35). For example, Shukrisarvey et al. studied the quality of the Tajan river in Mazandaran Province based on the biological Indices of the Hilsenhoff biological Index, Shannon-Wiener index, physicochemical parameters and environmental Indices (36). In another study, Heidari et al. evaluated the biological features of the Kashkan river in terms of the diversity and Macro Benthic population structure (37). Moreover, Ibiwumi et al. investigated the effects of

physicochemical parameters on Nigeria Macro Benthics via various Indices (38).

The purpose of this study, which was conducted for the first time in northwest Iran, was to investigate the effects of the physicochemical parameters of surface waters receiving effluents from fish farms on aquatic insects, determine the relationship between physicochemical parameters and biological indices, and also measure the correlation coefficients between qualitative and biological indices and provide basic information for the study and management of water resources in Bolaghlar area of Nair city, Ardabil Province.

Materials and Methods

This study is a cross-sectional descriptive study conducted to investigate the effects of physicochemical parameters in surface waters (Aglagan River) receiving the effluent of fish farms on biological Indices, determine the relationship between physicochemical parameters and aquatic insects and measure correlation coefficients between qualitative and biological Indices in Boulaghlar region, Nir City, over the period from March to August 2019. According to the topography of the region and the areas where the fish farms are located, as well as the status of other land uses in the region, nine stations were selected along the Aglagan river (Fig. 1). Six series sampling of aquatic insects were done from each station throughout the study period (54 samples collected in total), and they were detected down to family level by valid entomological keys (39, 40). To determine physicochemical parameters, three series of samples were collected from each station (27 samples in total). Based on the guidelines of the Standard Methods v. 20, the measured parameters in this study included fecal coliform (MPN/100 mL), biochemical oxygen (mg/mL), nitrate (mg/L), DO (mg/L), electrical conductivity, chemical oxygen demand (mg/L), ammonium (mg/L), phosphate (mg/L), turbidity (NTU), total hardness (mg/L.CaCO₃, total

dissolved solids (mg/L), temperature (°C) and pH. In order to analyze the correlation and relationship between the parameters and the obtained indicators, the collected data were statistically analyzed by SPSS v.22 software. To assess relationships among physicochemical parameters, Pearson's correlation coefficient was used; to determine the effects of physicochemical parameters on qualitative Indices and the effects of qualitative Indices on Hilsenhoff biological Index, regression correlational analysis was employed. In this study, NSFQI, IRWQISC and HBI Indices were used and calculated based on standard, relevant formulas (41, 42).

Results

Physicochemical parameters

The mean of measured physicochemical parameters in the samples used to measure NSFQI and IRWQISC Indices and also the amounts of NSFQI, IRWQISC and biological Indices are depicted in Table 1. As can be seen in Table 1, the minimum amount of electrical conductivity (0.38 μ S/cm) was observed in the upper stations of fish farms. The maximum amount of pH was detected in Station 2 (7.59) and the minimum was detected in Stations 7 and 8 (7.27). The minimum amounts of turbidity (1.03 NTU), COD (4.33 mg/L), nitrate (3 mg/L), phosphate (0.09 mg/L) and fecal coliform (4 colonies/100 ml, 7.67 MPN/100ml) belonged to Station 1. The maximum amounts of electrical conductivity (0.47 μ S/cm), hardness (189.67 mg/L CaCO₃), BOD (12.67 mg/L), ammonium (0.35 mg/L), nitrate (16.33 mg/L), phosphate (0.55 mg/L) and fecal coliform (27.33 colonies/100ml, 53.67 MPN/100ml) were detected in Station 3. The highest temperature parameter (17.53 °C) was reported in Station 5. The maximum amount of oxygen parameter (5.97 mg/L), and the minimum amounts of dissolved solids (284.33 mg/L), hardness parameter (94.27 mg/L CaCO₃), BOD (1.33 mg/L), ammonium (0.09 mg/L) and temperature (16.51 °C) were detected in Sta-

tion 6. The maximum amounts of turbidity (12.78 NTU) and COD (17 mg/L) were observed in Station 7. The highest amount of dissolved solids parameter (342 mg/L) belonged to Station 8, and the lowest amount of DO (4.74 mg/L) belonged to Station 9. The NSFQI index ranged from 54.45 to 76.21, which according to the results of this index, the studied stations can be put into 'good quality' and 'moderate quality' categories. The maximum (76.21) and minimum (54.45) amounts of this index belonged to Station 6 and based on the IRWQISC index, the studied stations ranged from 41.32 to 77.40 which puts them in the three categories of 'good quality', 'moderate quality' and 'fairly bad quality'. The maximum amount of this index (77.40) was observed in Station 6, and the minimum amount (41.32) was observed in Station 3.

Biological index

A total of 2,593 aquatic insects were collected from all the studied stations belonging to 9 families, 9 orders and 4 classes. The highest and lowest frequencies of the families of aquatic insects belonged to Gammarida (78.63%) and Enidae (0.11%), respectively. The Hilsenhoff biological Index ranged from 6.26 to 3.56; the highest index was observed in Station 9 with the biological quality of 'fairly poor', and the lowest index was observed in Station 1 whose biological quality was in accordance with the Hilsenhoff Index (Table 2).

Relationships among physicochemical parameters, qualitative index and biological index

The results of Pearson's correlation analysis (Table 3) revealed that there was not a significant relationship between temperature and other parameters. Investigating relationship between turbidity, TDS, phosphate, fecal coliform, nitrate, BOD, COD, electrical conductivity, total hardness and ammonium parameters showed a negative correlation with pH, DO and its saturation percentage parameters;

however, there were positive correlations between the above-mentioned parameters and the other parameters. The pH index had a significantly positive correlation with DO and its saturation ratio. The saturation percentage of DO had no significant relationships with fecal coliform. On the other hand, the investigation of the relationship between physicochemical parameters and the Hilsenhoff Index using linear regression showed that there were signifi-

cant relationships between this index and pH, DO and TDS parameters. Nevertheless, this effect was positive on TDS (Sig=0.046, B 0.027) yet negative on DO (Sig=0.037, B=-1.588) and pH (Sig=0.029, B=-4.837). Finally, there were no significant relationships between the Hilsenhoff Index and NSFQI (Sig= 0.151, B= 0.047) and IRQQI_{sc} (Sig= 0.179, B= -0.027) Indices (Table 4).




Fig. 1. Study area and location of samples sites in northwest of Iran, Ardabil Province, 2019




Fig. 2. Relationship between pH, Dissolved Oxygen (DO), Total Dissolved Solids (TDS) and Biological Index (BI) in studied stations (Northwest of Iran, Ardabil Province, 2019)

Table 1. Physicochemical parameters result in studies stations (Northwest of Iran, Ardabil Province, 2019)

Parameters	Unite	Stations								
		1	2	3	4	5	6	7	8	9
Turbidity	NTU	1.03	1.60	11.63	11.05	10.63	1.40	12.78	12.69	12.11
pH	-	7.58	7.59	7.29	7.43	7.45	7.55	7.27	7.27	7.28
BOD	mg/L	1.67	2.33	12.67	12	10	1.33	12	11.33	10.67
Fecal Coli-form	MPN/100 mL	7.67	9.67	53.67	41.67	29.67	12.67	53.67	45.67	29.67
Total Hardness	mg/L.CaCO ₃	97.27	101.23	189.67	169.30	174.73	94.27	191.23	188.87	187
DO	mg/L	5.77	5.69	5.19	5.24	5.14	5.97	5.11	4.94	4.74
Oxygen (Sat. Per.)	%	72.83	71.43	63.52	59.99	59.26	73.22	65.12	61.76	57.56
Temperature	°C	17.07	17.06	17.19	17.20	17.53	16.51	16.61	16.62	16.76
TDS	mg/L	285.67	290	328	317.67	324.33	284.33	334.33	342	339.33
Phosphate	mg/L	0.09	0.1	0.55	0.51	0.46	0.13	0.46	0.45	0.43
Nitrate	mg/L	3	3.67	16.33	15	13	3.33	16	15	12.67
Ammonium	mg/L	0.16	0.17	0.35	0.31	0.27	0.09	0.31	0.30	0.27
Colonies	Count	4	5.33	27.33	24.67	21.67	5	26.67	26.33	24
EC	µS/cm	0.38	0.38	0.47	0.46	0.46	0.38	0.45	0.46	0.46
COD	mg/L	4.33	5	15.33	14.67	12.67	3.67	17	16	13.67
NSFWQI	-	72.99	74.58	55.86	54.45	56.16	76.21	55.83	55.25	55.13
IRWQISC	-	77.35	75.24	41.32	42.40	45.39	77.40	41.63	42.55	45.80
BI	-	3.56	3.38	3.84	4.28	3.72	4.26	5.06	5.11	6.26

Table 2. Aquatic insect specimens collected and the calculated bio-index values in river of Northwest of Iran, Ardabil Province, 2019

Stations	Family									Total	BI	Water Quality
	Ephemeroidea	Notonectidae	Hydropsychidae	Chironomidae	Culicidae	Asellidae	Gammaridae	Glossiphoniidae	Enidae			
1	55	5	4	0	0	0	184	1	3	243	3.56	Excellent
2	38	1	2	0	0	0	85	2	0	128	3.47	Excellent
3	59	0	1	0	0	2	589	1	0	652	3.83	Very good
4	8	0	1	0	0	0	198	20	0	227	4.28	Good
5	67	2	0	0	0	0	404	1	0	474	3.76	Excellent
6	184	0	0	15	0	0	184	0	0	203	4.25	Good
7	0	0	4	55	0	0	152	1	0	212	5.05	Fair
8	2	0	4	69	0	0	172	1	0	248	5.11	Fair
9	9	0	5	72	4	37	71	8	0	206	6.26	Fairly poor
Total	422	8	21	211	4	39	2039	35	3	2593	3.97	Very good

Table 3. Pearson statistical test results and correlation between physicochemical parameters (Northwest of Iran, Ardabil Province, 2019)

Parameter	Turbidity	pH	Temp.	DO	Oxygen (Sat. Per.)	TDS	Phosphate	Colonies	Nitrate	BOD	COD	EC	Total Hardness	Ammonium	Fecal Coliform
Turbidity	1	-0.92	0.09	-0.93	-0.89	0.97	0.96	0.99	0.98	0.98	0.98	0.97	0.99	0.92	0.90
Ph		1	0.25	0.88	0.72	-0.95	-0.84	-0.91	-0.88	-0.88	-0.91	-0.86	-0.93	-0.83	-0.88
Temp.			1	-0.03	-0.26	-0.05	0.20	0.06	0.10	0.13	0.03	0.20	0.06	0.25	-0.04
DO				1	0.92	-0.97	-0.83	-0.89	-0.85	-0.88	-0.88	-0.90	-0.93	-0.83	-0.72
Oxygen (Sat. Per.)					1	-0.87	-0.88	-0.87	-0.84	-0.87	-0.83	-0.92	-0.88	-0.80	-0.65
TDS						1	0.89	0.95	0.92	0.92	0.94	0.93	0.97	0.87	0.84
Phosphate							1	0.98	0.98	0.98	0.95	0.98	0.96	0.94	0.91
Colonies								1	0.99	0.99	0.99	0.98	0.99	0.95	0.94
Nitrate									1	0.99	0.99	0.97	0.97	0.96	0.95
BOD										1	0.98	0.98	0.98	0.96	0.93
COD											1	0.95	0.98	0.95	0.94
EC												1	0.97	0.93	0.87
Total Hardness													1	0.93	0.90
Ammonium														1	0.91
Fecal Coliform															1

Coefficients description: value of -1 meaning a total negative linear correlation, 0 being no correlation, and + 1 meaning a total positive correlation.

Table 4. Relationship between physicochemical parameters and biological index (Northwest of Iran, Ardabil Province, 2019)

Model	Unstandardized Coefficients		Standardized Coefficients	T	Sig.*
	B	Std. Error	Beta**		
(Constant)	40.235	13.127		3.065	.018
pH	-4.837	1.771	-.718	-2.731	.029
(Constant)	3.517	.522		6.739	.000
Turbidity	.104	.054	.591	1.939	.094
(Constant)	12.816	3.283		3.904	.006
DO	-1.588	.617	-.697	-2.575	.037
(Constant)	31.918	14.011		2.278	.057
Temperature	-1.624	.826	-.596	-1.965	.090
(Constant)	9.871	3.151		3.133	.017
%DO	-.084	.048	-.551	-1.748	.124
(Constant)	-4.137	3.522		-1.175	.278
TDS	.027	.011	.676	2.426	.046
(Constant)	3.652	.671		5.445	.001
Phosphate	2.077	1.695	.420	1.225	.260
(Constant)	3.505	.601		5.832	.001
Colonies	.048	.029	.531	1.658	.141
(Constant)	3.588	.660		5.440	.001
Nitrate	.073	.054	.455	1.353	.218
(Constant)	3.634	.594		6.113	.000
BOD	.091	.063	.481	1.451	.190
(Constant)	3.361	.688		4.882	.002
COD	.090	.055	.525	1.632	.147
(Constant)	-.426	3.336		-.128	.902
EC	11.103	7.668	.480	1.448	.191
(Constant)	2.502	1.066		2.348	.051
Total Hardness	.012	.007	.568	1.828	.110
(Constant)	3.507	.998		3.515	.010
Ammonium	3.547	3.822	.331	.928	.384
(Constant)	3.755	.635		5.909	.001
Fecal Coliform	.020	.018	.393	1.132	.295
(Constant)	5.867	1.032		5.682	.001
IRWQI	-.027	.018	-.492	-1.495	.179
(Constant)	7.297	1.829		3.989	.005
NSFWQI	-.047	.029	-.520	-1.611	.151

*p-value < 0.05: significant

**Slope of linear regression equation

Discussion

Beneficial insects play an important role in human life. A group of these insects are considered as biological indicators of water quality, including Ephemeroptera, Plecoptera and Trichoptera. This group of insects indicates the appropriate quality of water for human use (43–46).

Based on the study findings in the previous

section, the activity of fish farms can negatively affect the amount of DO by decreasing the amount of this parameter in fish farms' effluent stations as compared to upper stations. According to the findings of this study, the amount of this parameter gradually reduced from the upper sections to the lower sections of the river showing that the effluent contained a lot of

wastes of fishes, leading, in turn, to an increase in nitrate, phosphate and solid materials concentration levels, water turbidity, a higher chemical decomposition level of organic materials and a stronger need for oxygen (47). In an agreement with the findings of this study, previous research conducted in Kermanshah Province, Khuzestan Province, Yassoj, Iranian Kurdistan, the United States and Turkey have also reported that the amount of DO in the effluent of fish farms is significantly reduced (47–50).

This study revealed that the amount of pH in Stations 3 and 7 was influenced by the effluent of fish farms and was thus significantly reduced. In a study by Khoshakhlagh et al. (51), several reasons were discussed for such a decline in pH including the decomposition of wastes of fishes and food leftovers in the effluent and an increase in organic materials and pollution. The results of this study agree with the findings reported by Boventura et al. (52) and Zarzuela et al. (53) but disagree with the findings of studies by Selong and Helfrich and Dugel (49, 54).

The findings of this study showed a significant increase in the parameters of TDS, electrical conductivity, fecal coliform, nitrate, ammonium, phosphate, BOD and COD in the effluent of fish farms. Millard et al., Hynes et al., Homewood et al., Stephens et al., Pillary et al., Adem et al., Miller and Semens and Sobhani et al. have all reported in their studies that food leftovers and remaining, waste of fishes, metabolic fish products and cleansing pools can lead to such an increase in the above-mentioned parameters (9, 49, 55–60). The findings of this study agree with those previously reported from Khuzestan, Shahr-e Kurd, Yassoj, Mazandaran Province, Golestan Province, Kermanshah Province, Turkey and Minnesota in the United States (45, 47, 49, 50, 61–63). A rise in dissolved solids can also increase the amount of turbidity as also reported by Hosseini et al. (45). Because the physicochemical parameters including turbidity, TDS, phosphate, faecal coliform, nitrate, BOD, COD, electrical conductivity, total hard-

ness, and ammonium are all somehow related to oxygen consumption, so they can lead to a decrease in DO and, as a result, oxygen saturation percentage. Also, some of these compounds, such as nitrate and ammonium, cause changes in the pH of water due to some chemical interactions, especially with the production of acidic compounds, so the reason for the negative correlation between these parameters can be explained.

The findings of this research revealed an increase in the temperature of the effluent of fish farms throughout the year. Khoshakhlagh et al. named several possible reasons for such a boost in the temperature: the repelled heat of fish metabolism, water touching the bed and walls of the pools, absorbing more sunlight because of suspended food leftovers, extra energy produced by pollutant influents and decomposition of organic compounds by microorganisms in the water (51). A study conducted by Axeler et al. also corroborates the findings of this study (64) which revealed an increase in the total amount of hardness in the effluent of fish farms. These results are in agreement with previous research findings conducted in Minnesota in the United States, Kermanshah Province and Chaharmahal and Bakhtiari Province (48, 52, 64). Boyd has stated that using calcic materials produced by limestone in fish farms can lead to an increase in the overall hardness of water (65).

Stations 1, 2 and 6 located in the upper section of fish farms, and hence safeguarded against effluents, can fit into the ‘good quality’ category based on NSFQI and IRWQI_{SC} Indices. The changes resulted from fish farms activities significantly reduced the amount of NSFQI and IRWQI_{SC} Indices in fish farms’ effluent stations; thus, they can be categorized into the ‘moderate quality’ and ‘fairly bad quality’ categories, respectively. Based on the NSFQI index, all the stations located in the lower section of the river belong to the ‘moderate quality’ category whereas based on IRWQI_{SC} index, stations 4 and 8 fall into the

‘fairly bad’ category and stations 5 and 9 fit into the ‘moderate quality’ category. Such findings are in line with the results observed in Golestan Province based on the NSFQI index (63). All the studies conducted on rivers receiving the effluent of fish farms in East Azerbaijan Province, Golestan Province and Gorgan University based on the NSFQI index have proved the effects of fish farms’ activities on the quality of rivers by various degrees (23, 66–69).

According to the results, stations 1 and 2, which were located in the upper section of the first fish farm and were guarded against the pollutants of this farm, hosted a higher number of Amphipoda and Ephemeroptera species, which mostly live in freshwaters; their biological index was also excellent in quality. The reason for the higher numbers of Amphipoda and Ephemeroptera in the effluent station of the first fish farm (Fig. 2) is probably because the water was not polluted enough to negatively affect such arthropods. The relative effect of effluent pollutants from the first fish farm on station 4 was significant in a way that the number of the above-mentioned species drastically reduced in this station; Leach species was also detected and collected in this station. An increase in the number of such species in Station 5 shows that the environment for their thriving is quite felicitous. Located in the upper section of the second fish farm, Stations 6 hosted Amphipoda and Chironomidae species (two species which are less sensitive to pollution). Thus, Stations 3 and 6 fit into the ‘very good quality’ category regarding biological index. The number of Amphipoda species decreased in Station 7 and other later stations, but the number of other less-sensitive species, such as Chironomidae, increased; some species which are frequently observed in polluted waters, such as Diptera and Isopoda, were also observed in Station 9. Therefore, Stations 7 and 8 fall into the ‘good quality’ category and Station 9 into the ‘moderate quality’ category. In general, it seems that because of the short-

covering distance of the study, the negative effects of pollutants resulting from the effluent of fish farms were higher in lower stations (70).

The existence of a positive correlation between fecal coliform and dissolved solids is acceptable because an increase in the amount of fecal coliform could increase the load of organic compounds in water, leading in turn to a rise in the electrical conductivity index. The negative correlation between fecal coliform and DO is also acceptable because when the coliform-containing effluent enters the river, the oxygen level drastically decreases. The positive correlations among dissolved solids, turbidity and electrical conductivity result from their mutual effects on each other because when the number of dissolved solids increases, the amounts of turbidity and electrical conductivity of water also increase. The positive correlations among nitrate, electrical conductivity, phosphate, BOD and COD parameters are a result of deriving from the same source in a way that an increase in the amount of nitrate can raise the amounts of other mentioned parameters (71).

The investigation of the effects of physiochemical parameters and Hilsenhoff biological Index on other parameters of the study by the linear regression analysis revealed that there were significant relationships between these two Indices and pH, DO and dissolved solids parameters. These results agree with the findings of the studies conducted in Nigeria in terms of pH parameter, in Mazandaran Province and Lorestan Province in terms of dissolved solids parameter and finally in Nigeria, Mazandaran Province and Lorestan Province in terms of DO (36–38). This effect was positive regarding dissolved solids parameter (Sig= 0.046, B= 0.027) but negative regarding DO parameter (Sig= 0.037, B= -1.588) and pH parameter (Sig= 0.029, B= -4.837). The quantity and quality of the influent organic materials pouring into water resources (72) are influenced by fish farm activities and the effluent discharge of such farms (52, 71). This can pol-

lute waters and negatively affect the quality and quantity of water resources (66, 72, 73), and effective parameters can bring about drastic changes in the ecosystem, such as losing diversity, losing the whole population of one species, increasing the number of species feeding on organic materials, losing sensitive species and replacing resistant species (72). Finally, such changes can result in an increase or decrease in the biological index as observed in the stations in this study. The relationship between effective parameters and the Hilsenhoff biological Index is represented in Fig. 2. As observed, there were no significant relationships between the Hilsenhoff biological Index and the two qualitative Indices of NSFQI and IRWQI showing that the measured parameters are not similar in the above-mentioned Indices.

To conduct this study, there were some limitations that faced challenges, including economic limitations and problems related to sampling. These problems were solved by conducting composite sampling and managing points and the number of samples so that the resulting data have the maximum compatibility with the real conditions of the studied area. Also, there were problems in the analysis of some parameters, which were solved by using portable devices.

Conclusion

The stations located in the upper section of fish farms can be categorized into the ‘good quality’ category based on the NSFQI and IRWQI_{SC} Indices. The activities of fish farms can negatively impact relevant parameters that affect the NSFQI index and its lower values in the effluent receiving stations and even in lower-section stations, hence putting these stations in the ‘moderate quality’ category. However, in terms of the IRWQI_{SC} index, the effluent stations and the lower ones all belong to ‘fairly bad quality’ and ‘moderate quality’

categories. The main reason for the changes in the above indicators along the river and especially in the downstream stations of the farms is related to receiving effluents from the farms and the presence of relatively high amounts of various pollutants based on the results of the tests. The two water quality indicators generally had almost the same trends in all stations and minor changes are caused by the parameters including BOD, nitrate, phosphate and etc. in their calculation. Therefore, it is necessary to consider the necessary arrangements for different uses of water in downstream of the river and act with caution for drinking and household uses. The linear regression analysis revealed a significant relationship between the Hilsenhoff biological Index and the physicochemical parameters of pH, DO and total dissolved solids. Given that the activity of fish farms and their effluent discharge can negatively change the physical, chemical and biological parameters of receiving waters, it is essential to consider necessary treatments for such effluents so that health and ecological risks are efficiently reduced. For purposes such as drinking, it is recommended to use different water treatment methods, including observing the physical distance of water harvesting for the possibility of self-purification, physicochemical treatment and disinfection, as much as possible depending on the local conditions, so that the quality of water could be acceptable in terms of national standards for various uses as well as surface water. Moreover, it is of high importance to conducting more studies in this field, continuous monitoring and measure Indices relevant to the type of utilization and check them against with the standards.

Acknowledgements

We hereby express our sincere gratitude to the Vice President for Research at Ardabil University of Medical Sciences who approved and financially supported this research project

and also to the experts working at Water and Wastewater Laboratory of School of Health who cooperated in conducting relevant experiments.

Ethical considerations

The study was approved by the Ethical Committee of Ardabil University of Medical Sciences, Iran (Code of ethics: IR. ARUMS. REC.1397.195). This study was financially supported by the Ardabil University of Medical Sciences (Project No. 314)

Conflict of interest statement

The authors declare there is no conflict of interest.

References

- Majnunian H (1998) River conservation, biophysical attributions, habitat values and extraction rules. 1st Ed. Tehran. environmental protection organization. In: EPA, editor. Tehran, EPA. p. 16–22.
- USEPA (1979) National secondary drinking water regulations.
- Afshinnehzad Y (1994) Rivers of Iran: Ministry of Energy, Jamab Consulting Engineers.
- Yusefzadeh A, Shams Khoram Abadi G, Godini H, Hossein Zadeh I, Safari M (2014) Evaluation of water quality of Khoramrood River Khorramabad with water quality index (NSFWQI) and its zoning with Geographic Information System (GIS). *Yafteh*. 15(5): 82–92.
- Behbahaninia A, Salmasi R (2008) Investigation and recognition of physicochemical properties of Jajrood River pollutants. *Hum Environ*. 6: 34–43.
- Nastiti KDM, Sayekti RW, Yuliani E (2023) Water quality and distribution of Selorejo Reservoir using the Pollution Index, Oregon-WQI, and NSF-WQI methods. *J Tek Pengair: J Water Resour Eng*. 14 (1): 64–75.
- Fortes ACC, Barrocas PRG, Kligerman DC (2023) Water quality indices: Construction, potential, and limitations. *Ecol Indic*. 157: 111187.
- Sánchez E, Colmenarejo M, Vicente J, Rubio A, García M, Travieso L, Borja R (2007) Use of the water quality index and dissolved oxygen deficit as simple indicators of watersheds pollution. *Ecol Indic*. 7(2): 315–328.
- Bordalo A, Nilsumranchit W, Chalermwat K (2001) Water quality and uses of the Bangpakong River (Eastern Thailand). *Water Res*. 35(15): 3635–3642.
- Karr JR (1998) Rivers as sentinels: using the biology of rivers to guide landscape management. *River ecology and management: lessons from the Pacific Coastal ecoregion* Springer, New York.
- Siriwardhana KD, Jayaneththi DI, Herath RD, Makumbura RK, Jayasinghe H, Gunathilake MB (2023) A Simplified Equation for Calculating the Water Quality Index (WQI), Kalu River, Sri Lanka. *Sustainability*. 15(15): 12012.
- Boyacioglu H (2006) Surface water quality assessment using factor analysis. *AJOL*. 32(3): 389–393.
- Pesce SF, Wunderlin DA (2000) Use of water quality indices to verify the impact of Córdoba City (Argentina) on Suquia River. *Water Res*. 34(11): 2915–2926.
- Jonnalagadda S, Mhere G (2001) Water quality of the Odzi River in the eastern highlands of Zimbabwe. *Water Res*. 35 (10): 2371–2376.
- Larijani S, Hossein B, Kaviani A, Ziaei AN (2022) Sensitivity assessment of two water quality models NSFWQI and IRWQISC in Haraz River. *Iran J Irrig Drain*. 16(4): 816–830.

16. Asadpour Lomer S, Ebrahimi K, Liaghat A, Alipour F (2024) Assessing Temporal-Spatial Variations and Classifying Water Quality of the Dinachal River, Iran, through Field Data Collection. *Iran J Soil Water Res.*
17. Abbaspour M, Javid A, Habibi E (2014) Determination of Physical and Chemical Parameters of Khersan River Water and Investigation of its Annual Changes. *Int J Environ Sci Tech.* 15(4): 1–11.
18. Farzadkia M, Djahed B, Shahsavani E, Poureshg Y (2015) Spatio-temporal evaluation of Yamchi Dam basin water quality using Canadian water quality index. *Environ Monit Assess.* 187: 1–15.
19. NSF (2012) National Sanitation Foundation. Available at: <http://www.Nsfconsumer.org>.
20. Nikunahad A, Kazembeigi F (2009) Comparison of water quality parameters for choosing the best of the index in the Karkheh Dam. *Iran Water Res J.* 3(4): 69–73.
21. Fabiano D, Altair B, Marcia C, Sonia M, Maria J (2008) Water quality index as a simple indicator of aquaculture effects on aquatic bodies. *Ecol Indic.* 8(5): 476–484.
22. Torabian A, Hashemi S, Khalili R, Ferdusipoor S (2004) Effects of harvesting on water quality in the river downstream of the dam Mamlu using QUAL2E model. *J Environ Stud.* 30(35): 69–76.
23. Hoseinzadeh A, Korsandi H, Rahimi N, Hosizadeh S, Alipour M (2013) Water quality assessment of AydooghMosh River using the NSFQI Quality Index and Liou Pollution Index. *Uromiyeh Med J.* 24(2): 156–162.
24. SawooEP Agency (2014) Standard of Iran Water Quality Index In: Agency EP, editor. Tehran: EPA.
25. Shaker Donyavi A, Cheloyan P, Moridi A, Khalili R (2023) Assessment of Spatio-Temporal Changes in Water Quality of Babilroud River Using IRWQIsc, NSFQI and CCME Water Quality Index. *J Water Irrig Manag.* 13(2): 551–563.
26. Hafezieh M (2001) Aquatic insects as indicators of organic contamination of water. *Iran J Fish Sci.* 10(1): 19–36.
27. Lenat DR (1993) A biotic index for the southeastern United States: derivation and list of tolerance values, with criteria for assigning water-quality ratings. *J North Am Benthol Soc.* 12(3): 279–290.
28. Mokhtari SA, Farzadkia M, Esrafil A, Kalantari RR, Gholami M (2017) Application of dispersive liquid-liquid micro-extraction as a simple assisted clean-up and preconcentration technique for GC/MS determination of selected PAHs extracted from sewage sludge by Soxhlet and ultrasound assisted extraction method. *Desalination Water Treat.* 66(5): 176–183.
29. Hilsenhoff WL (1988) Rapid Field Assessment of Organic Pollution with a Family-Level Biotic Index. *J North Am Benthol Soc.* 7: 65–68.
30. Taowu M, Qinghui H, Hai W, Zijian W, Chunxia W, Shengbiao H (2008) Selection of benthic macroinvertebrate-based multimetrics and preliminary establishment of biocriteria for the bioassessment of the water quality of Taihu Lake, China. *Acta Ecol Sin.* 28(3): 1192–1200.
31. NafisiBehabadi M, Ahmadi M (2001) Identification of invertebrate species of current waters. Khaibar, Tehran.
32. Balayo C (2019) Determining the diversity and distribution of macroinvertebrates in Tende Sub-Bay of Murchison Bay, Lake Victoria (Pearl Aquatics Cage Fish Farm). Undergraduate Dissertation. School of Biosciences Collection. Makerere University. Uganda.
33. Nasri M, Ghoghji A, Podeh HT (2023) Investigation of Zayandeh River water quality based on Macrobenthos as biological indicators. *Int J Aquat Biol.* 11(3): 213–221.
34. Zhang J, Jiang P, Chen K, He S, Wang B, Jin X (2020) Development of biological water quality categories for streams us-

- ing a biotic index of macroinvertebrates in the Yangtze River Delta, China. *Ecol Indic.* 117: 106650.
35. Malvandi H, Moghanizade R, Abdoli A (2021) The use of biological indices and diversity indices to evaluate water quality of rivers in Mashhad, Iran. *Biologia.* 76: 959–971.
 36. Shukrisarvey M, Ahmadi M, Rahmani H, Kamrani E (2015) Evaluation of Water Quality Based on Hilsenhoff Biological Indices, Shannon-Wiener Diversity and Environmental Indices in Tajan River. *J Fish Sci Tech.* 3(4): 43–55.
 37. Heydari N, Yazdian H, Zahrai B, Nematollah JafarzadehhagigiFard (2011) Biological Assessment of Kashkanrood River Based on Diversity and Population Structure of Macrobenthos. The 1st National Conference on Environmental Protection and Planning, 2014 February 27, Hamadan Islamic Azad University, Hamadan, Iran.
 38. Ibiwumi AA, Kabir PKO, Abiodun SA (2013) Impact of Physicochemical Parameters on Benthic Macro-invertebrates Assemblage of Erelu Reservoir in Oyo Town, Nigeria. *Asian J Biol Sci.* 2(12): 328–336.
 39. Mehrjo F, Abdoli A, Hashemi SH, Hosseinabadi F (2020) Development of a multimetric index based on benthic macroinvertebrates for downstream Jajrud in Tehran Province, Iran. *Iran J Fish Sci.* 19(1): 286–296.
 40. Feldman D, Farris J, Moore M, Cooper C (2010) A characterization of benthic macroinvertebrate communities in agricultural drainage ditches of the northeast Arkansas Delta, USA. *Agricultural drainage ditches: Mitigation wetlands for the 21st century Research Signpost, Kerala*, pp. 17–35.
 41. Bui DT, Khosravi K, Tiefenbacher J, Nguyen H, Kazakis N (2020) Improving prediction of water quality indices using novel hybrid machine-learning algorithms. *Sci Total Environ.* 721: 137612.
 42. Rostami MH, Vahidi H, Tayefeh M, Ahmadikallan M (2020) Evaluating water quality of kangir river by means of NSF water quality index and the effect of reducing the load of point pollutants on increasing the index rate using QUAL-2KW model. *J Wetl Ecobiol.* 12(1): 81–98.
 43. Malekei-Ravasan N, Bahrami A, Shayeghi M, Oshaghi MA, Malek M, Mansoorian AB, Vatandoost H (2013) Notes on the Iran Caddisflies and Role of Annulipal-pian Hydropsychid Caddisflies as a Bio-monitoring Agent. *J Arthropod Borne Dis.* 7(1): 71–82.
 44. Abbasi M, Doosti S, Vatandoost H, Hosseini-Vasoukolaei N (2020) Study on the Fauna of Aquatic Insects in Northwestern Iran. *J Arthropod Borne Dis.* 14(1): 1–7.
 45. Shaeghi M, Dehghan H, Pakdad K, Nikpour F, Absavaran A, Sofizadeh A, Akhavan AA, Vatandoost H, Aghai-Afshar A. Faunistic Study of the Aquatic Arthropods in a Tourism Area in Northern Iran. *J Arthropod Borne Dis.* 2017 May 27. 11(2): 286–301.
 46. Shayeghi M, Vatandoost H, Gorouhi A, Sanei-Dehkordi AR, Salim-Abadi Y, Karami M, Jalil-Navaz MR, Akhavan AA, Shiekh Z, Vatandoost S, Arandian MH. Biodiversity of Aquatic Insects of Zayandeh Roud River and Its Branches, Isfahan Province, Iran. *J Arthropod Borne Dis.* 2014 Apr 9. 8(2): 197–203.
 47. Hosseini SH, Sajjadi MM, Kamrani E, Sourinejad I, Ranjbar H (2013) Impact of rainbow trout (*Oncorhynchus mykiss*) farm effluents on water physico-chemical parameters of Ryjab River (Kerman-shah Province). *J Aquat Ecol.* 2(4): 39–29.
 48. Weisi T, Ahmadifard N, Agh N, Kamali M (2019) The effect of Palangan Fish Farming on the macro-invertebrate of the Sirvan River Using Shannon and Hilsinhof

- Biological Indices. *Anim Environ.* 11(3): 345–354.
49. Gasemi S, Mafigolami R (2015) Effects of effluent from Ponds Nourali district fish farming on the changes in the oxygen content of the Gargar River and provides an optimal management plan for fish pools. *Quart J Wetl Ecol.* 28: 75–84.
 50. Fadaeifard F, Raissy M, Faghani M, Majlesi A, NodehFarahani G (2012) Evaluation of physicochemical parameters of waste water from rainbow trout fish farms and their impacts on water quality of Koohrang stream Iran. *Int J Fish Aquac.* 4(8): 170–177.
 51. SobhanArdakani S, Mehrabi Z, Ehteshami M (2014) Effect of Aquaculture Farms Wastewater on Physicochemical Parameters of Kabkian River, 2011–12. *Mazandaran Med J.* 24(113): 140–149.
 52. Pulatsu S, Rad F, Köksal G, Aydın F, Benli AK, Topçu A (2004) The impact of rainbow trout farm effluents on water quality of Karasu stream, Turkey. *Turk J Fish Aquat Sci.* 4(1): 9–15.
 53. Khoshakhlagh M, Kamrani E, Abrahamdarajeh E, Sorinejad E (2015) The effect of Rainbow trout farms effluents on benthic macro-invertebrates of Marber River in Semirom. *J Aquat Ecol.* 5(1): 103–112.
 54. Cairns J, Dickson K, editors (1973) Biological methods for the assessment of water quality: ASTM.
 55. Boaventura R, Pedro AM, Coimbra J, Lencastré E (1997) Trout farm effluents: characterization and impact on the receiving streams. *Environ Pollut.* 95(3): 379–387.
 56. Selong J, Helfrich L (1998) Impacts of trout culture effluent on water quality and biotic communities in Virginia headwater streams. *N Am J Aquac.* 60(4): 247–262.
 57. Hynes HBN (1977) A Key to the Adults and Nymphs of the British Stoneflies (Plecoptera). *Freshwater Biological Assn.*
 58. Miller D, Semmens K (2002) Waste Management in Aquaculture. College of Agriculture, Forestry, and Consumer Sciences West Virginia University.
 59. Boyd CE (2003) Guidelines for aquaculture effluent management at the farm-level. *Aquac.* 226(1–4): 101–112.
 60. Homewood J, Purdie D, Shaw P (2005) Influence of sewage inputs and fish farm effluents on dissolved nitrogen species in a chalk river. *Water Air Soil Pollut.* 4(6): 117–125.
 61. Stephens WW, Farris JL (2004) A bio-monitoring approach to aquaculture effluent characterization in channel catfish fingerling production. *Aquac.* 241(1–4): 319–330.
 62. Maillard VM, Boardman GD, Nyland JE, Kuhn DD (2005) Water quality and sludge characterization at raceway-system trout farms. *Aquac Eng.* 33(4): 271–284.
 63. Rennert B (1994) Water pollution by a land-based trout farm. *J Appl Ichthyol.* 10(4): 373–378.
 64. Naderijelodar M, Esmaelisari A, Ahmadi M, Sefabadi S, Abdoli A (2010) Investigation of Pollution Caused by Rainbow Trout Fish Workshops on Water Quality Parameters of the River of Haraz. *Environ Sci.* 4(2): 21–36
 65. Mazaheri Kohanestani Z, Ghorbani R, Hajimoradloo A, Naeimi A (2014) The effect of trout farm effluents on the water quality parameters of Zaringol Stream (Golestan, Iran) based on NSFQI and WQI indexes. *Environ Resour Res.* 1(2): 191–201.
 66. Axler RP, Tikkanen C, Henneck J, Schuldt J, McDonald M (1997) Characteristics of effluent and sludge from two commercial rainbow trout farms in Minnesota. *Fish culture.* 59(2): 161–172.
 67. Boyd CE (2015) Calcium and magnesium use in aquaculture. *Global Aquaculture Alliance, USA.*

68. Mahdi Sadeghi AB, Nasser Bay, Nafiseh Soflaie, Mohammad Hadi Mehdinejad, Morteza Mallah (2015) The survey of Zarin-Gol River water quality in Golestan Province using NSF-WQI and IRWQISC. *J Health Fld.* 3(3): 27–33.
69. Noroozi N, Sadeghi M, Shahbazi A (2020) Study of changes in water quality index of Ziarat River at 2018 in Gorgan. *J Res Environ Health.* 6(3): 250–259.
70. Nasirahmadi K, Yosefi Z, Tarasoli A (2013) Water quality zoning of the river Haraz according to NSFQI index. *Mazandaran Med J.* 22(92): 64–71.
71. Ramezani A, ahmadimoghadam M, jafari M (2014) Water Qualitative Zoning of Goderkhosh River Based on NSFQI Index and Applying Geographic Information System (GIS). *Iran J Watershed Manag Sci Eng.* 8(25): 1–8.
72. Shayeghi M, MoradiAsl E, Saeidi Z, Mozafari E, Poudat A, Salimi M, Vatan-dooost H (2016) Aquatic Insects Fauna of Meshkin Shahr, Ardabil Province, North-western Iran, 2014. *J Marine Sci Res Dev.* 6(4): 1–3.
73. Ebrahimi E, Fathi P, Ghodrati F, Naderi M, Pirali A (2017) Assessment of Tajan River water quality with the use of biological and quality indicators. *Iran Sci Fish J.* 26: 139–152.