

Original Article

Relationship between Some Environmental and Climatic Factors on Outbreak of Whiteflies, the Human Annoying Insects

Abedin Saghafipour¹; *Alireza Zahraei-Ramazani²; Hassan Vatandoost^{2,3}; Amin Asadollahi²; Reza Fouladi-Fard⁴; Amir Hamta⁵; Ali Hasanwand²

¹Department of Public Health, Faculty of Health, Qom University of Medical Sciences, Qom, Iran

²Department of Medical Entomology and Vector Control, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran

³Department of Environmental Chemical Pollutants and Pesticides, Institute for Environmental Research, Tehran University of Medical Sciences, Tehran, Iran

⁴Research Center for Environmental Pollutants, Qom University of Medical Sciences, Qom, Iran

⁵Clinical Research Development Center (CRDU), Qom University of Medical Sciences, Qom, Iran

(Received 27 Mar 2019; accepted 21 Mar 2020)

Abstract

Background: The reports of numerous outbreaks of whiteflies from different parts of the world have increased its medical importance. The aim of this study was to determine relationship between environmental changes and climatic factors with the outbreak of the whitefly population in Tehran, the capital of Iran.

Methods: This study was carried out in urban areas of Tehran, where the increasing population of whiteflies was reported frequently during 2018. In order to entrap the whiteflies, 20 yellow sticky cards smeared with white refined grease were installed on the trunks of the trees at twice per month as trapping time intervals. The captured flies were transferred and conserved in cans containing 70% alcohol and were counted accurately under a stereomicroscope. To determine the relationship between air quality index, precipitation, air temperature and air humidity as environmental and climatic factors with the abundance of whiteflies, change point analysis and Generalized Estimating Equations (GEE) was used.

Results: The most density of white flies per trap was 256.6 and 155.6 in early October and late September respectively. The number moved closer to zero from November to April. The population of whiteflies was inversely correlated with the level of air quality index ($p=0.99$) and precipitation ($p=0.95$), and it had a direct correlation with the high temperature. Also, the population of whiteflies had a direct correlation with the level of air humidity in the first half of the year

Conclusion: According to these findings, during spring and summer from early May to early October.

Keywords: Change point analysis; Climatic factors; Environmental change; Outbreak; Tehran

Introduction

Whiteflies (Hemiptera: Aleyrodidae) feed on a wide range of hosts in a way that for some species more than 900 plant species have been identified (1, 2). Cucurbits and ornamental plants, agricultural crops, palms, and weeds are the main hosts of this pest, though there are many weeds which are the secondary hosts (3). The life cycle of whiteflies from egg to adult complete one month depending upon environmental temperature. Adult whiteflies may be surviving for one to two months (4). This insect is con-

sidered a health problem and an important medical pest that can also threaten human safety in some cases. Accidental entry of a whitefly into the human respiratory tract can cause inflammation and infection in the upper respiratory tract leading to the emergence of opportunistic fungal and bacterial infections (5). The population of this insect has increased in many parts of the world which has caused many problems for humans, especially in urban areas (6). According to experts, repeated and uncontrolled use of various formulations and concentrations

*Corresponding author: Dr Alireza Zahraei-Ramazani, E-mail: azahraei@tums.ac.ir

of pesticides can have many adverse effects, such as the resistance of pests to pesticides, the emergence of new pests, and the eradication of natural enemies (parasitoids and predators). Whiteflies are among the pests that have been evolved by continuous use of chemicals and the lack of proper management of pesticides (7). In addition to direct physical and biological harm for human, these insects cause a sharp decline in the production of agricultural products. Also, these flies cause the growth and development of saprophytic fungi on their honeydew which reduces the quality and nutritional value, as well as the consistency and shape of the products. Moreover, they physically damage non-productive plants. Nowadays, whiteflies, due to their increased resistance to various types of pesticides (8), cause more damage to a large number of crops and ornamental plants. Male and female winged insects feed on the leaf juice of plants. This leads to yellow spots on the leaves that directly damage the host plant and make it seem short and sick. Insect secretions on plant leaves can cause fungal growth (9). So far, about 1556 species of whiteflies have been identified in different parts of the world (10). Whiteflies in Iran were identified for the first time during the faunistic research and the identification of flies in Fars Province in 1995 (11). After that 14, 18 and 24 species of whiteflies have been reported in Isfahan (9), Gilan (12) and Golestan Provinces respectively (13). Subsequently, in 2000, morphological and biological studies were conducted on common species in Esfahan, which revealed that whiteflies in Esfahan were of the European race (9). Such flies are greenhouse pests, but, unfortunately, the lack of proper management of chemicals and pesticides has caused resistance to some of the pesticides and adverse environmental conditions in these insects. As a result, these flies have been able to adapt themselves to greenhouse conditions and easily grow and reproduce. Some experts also believe that these flies have been released into the open air through whitefly infestations, and since the ecosystem cycle and the population of predator in-

sects have not been balanced in the environment, this has led to a widespread outbreak of whiteflies in the open air (14). Moreover, since these insects are expected to reproduce in places more similar to greenhouse in terms of climate and food resources, it is necessary to identify the ecological factors in the reproduction of the insects, and take measures to control them outdoors (15). Over the past few years, the outbreak of whiteflies in Iran, especially in the residential areas of Tehran, has caused different allergies to humans. Therefore, the present ecological study was conducted with the aim of determining the relationship between some environmental and climatic factors on outbreak of whiteflies, the human annoying insects Tehran; District 6 in 2018.

Materials and Methods

Study area

The current study was carried out in the urban area of Tehran; the capital of Iran (District 6) suffers from a severe air quality index, where the abundance of whiteflies has been reported frequently during 2018 (Fig. 1).

The city of Tehran is located between the mountainous region and the plain. Tehran's climate is generally described as mild in spring and fall, hot and dry in summer, and cold and wet in winter. Based on the 2016 census, Tehran has a population of approximately 10 million. Vegetation coverage in Tehran is including natural and hand-planted forests (16). The air pollution indicators including Air Quality Index (AQI) with monthly activity of whiteflies were received from the Iran Meteorological Organization (17).

Study design

To entrap whiteflies, 20 yellow sticky cards (fly traps) of 30×50cm (18), depending on the diameter of the tree, were installed on the trees each time at twice per month as trapping time intervals from Early April to late March 2018. Totally 480 yellow sticky cards have been ap-

plied for catching adult whiteflies. In addition, we replaced the old traps with new one in each visit.

The chemicals on the traps were odorless and smeared with purified white grease purchased wholesale from reputable stores. The traps were installed on the trees of Maple, Acacia, European ash, Sycamore, and Berries for the purpose of catching whiteflies. In this regard, to examine the effect of colorful traps on whiteflies, three different colors of yellow, blue and green for attracting and capturing of them were used in a selected greenhouse. In this study, colored cards of 10×22cm in two heights of 1.3 and 2.3m were installed on tomato plants. After two weeks, the numbers of trapped insects on colored cards were counted.

Whiteflies were isolated and counted in two ways: traps (sticky cards) were immersed in warm water for 2 minutes (min) until the grease on the surface containing the flies was softened, and the flies were released in hot water. Then, the insects were removed from the hot water using a brush and were transferred and conserved in cans containing 70% alcohol. Then, at appropriate times, the whiteflies in the canned glass were released into the appropriate plates and carefully counted under a stereomicroscope. In another method, the traps containing whiteflies were peeled off the trees, and given the grid pattern on the surface of traps, the number of flies in every grid was randomly counted and the total number of whiteflies was estimated on the surface of the traps. The yellow and blue colored traps used in this study were supplied by Russel IPM Company; and the green traps were made of Tangle foot adhesive supplied by Kerman Chemistry Company (applied on green cards with brush).

Statistical analysis

Change analysis is organized to answer two questions 1) whether there are any change points or not 2) if so in which times change point(s) occurs. We deal with Hypothesis test and estimation in first and second question, respectively.

Generally, suppose X_1, X_2, \dots, X_n to be sequence of independent random variables with probability distribution function F_1, F_2, \dots, F_n respectively. Change point analysis intended to test following null hypothesis $H_0: F_1 = F_2 = \dots = F_n$ versus alternative hypothesis $H_1: F_1 = \dots = F_{k_1} \neq F_{k_1+1} = \dots = F_{k_2} \neq F_{k_2+1} = \dots = F_{k_q} \neq F_{k_q+1} = \dots = F_n$

Where $1 < K_1 < K_2 < \dots < K_q < n$ and q is unknown parameter that shows number of change points and K_1, K_2, \dots, K_q are the change points and have to be estimated. Although in majority of studies which point analysis has been applied probability distribution function supposed to be normal (19), in current study with the respect to response variable, it is let be Poisson. So our problem to detect change point (s) refer to X_1, X_2, \dots, X_c that are consequence of Poisson random variable with parameters $\lambda_i, i = 1, 2, c$ respectively. The aim was to test following null hypothesis $H_0: \lambda_1 = \dots = \lambda_c$ versus alternative hypothesis

$$H_1: \lambda_1 = \dots = \lambda_k = \lambda \neq \lambda_{k+1} \dots = \lambda_c = \lambda'$$

that shows there are k unknown change points. Likelihood function based on null hypothesis is:

$$L_0(\lambda) = \frac{e^{-c\lambda} \lambda^{\sum_{i=1}^c x_i}}{\prod_{i=1}^c x_i!}$$

so likelihood estimation based on that is:

$$\hat{\lambda} = \frac{\sum_{i=1}^c x_i}{c}$$

and under alternative hypothesis likelihood function is obtained

$$L_1(\lambda, \lambda') = \frac{e^{-k\lambda} \lambda^{\sum_{i=1}^k x_i} \cdot e^{-(c-k)\lambda'} \lambda'^{\sum_{i=k+1}^c x_i}}{\prod_{i=1}^k x_i! \cdot \prod_{i=k+1}^c x_i!}$$

by letting

$$M_k = \sum_{i=1}^k x_i, \quad M = \sum_{i=1}^c x_i, \\ M_k' = M - M_k = \sum_{i=k+1}^c x_i$$

the likelihood estimation function of λ and

$$\lambda' \text{ are given by } \hat{\lambda} = \frac{M_k}{c}, \hat{\lambda}' = \frac{M_k'}{c-k}.$$

SIC¹ under null hypothesis is found as

$$SIC(c) = -2 \log L_0(\hat{\lambda}) + \log c$$

and corresponding alternative hypothesis determined from

¹ Schwarz information criterion

$$SIC(k) = -2\log L_1(\hat{\lambda}, \hat{\lambda}') + 2\log c$$

So, according to the information criterion principle H_0 was rejected if $SIC(c) > \min_{1 \leq k \leq c-1} SIC(k)$. In order to estimating change point(s) $SIC(\hat{k}) > \min_{1 \leq k \leq c-1} SIC(k)$ was used (19, 20). After determination of change point (s) in the next stage for considering correlation between data, which were collected over time, marginal longitudinal model was applied. Link function because of being count response variable, log Poisson fulfilled. Coefficients were estimated by Generalized Estimating Equations (GEE). This method provides predictive model for the response variable by explanatory variables and it takes into account possible correlation between repeated measures of depend variable of a subject. In this study since data are collected over time it is likely that repeated measure on an individuals are correlated. When GEE method is used in order to analyze longitudinal data, correlation structure is formulated and possible correlations between measurements over time is incorporated (21).

Suppose that Y_{ij} is a count and we are interested to investigate changes in expected count to the covariates. Since counts are often modeled as poisson random variable, using poisson variance function and log link function, marginal model for Y_{ij} would be illustrated as follow:

The mean of Y_{ij} (μ_{ij}) is related to the independent variables by log link function:

$$\log(\mu_{ij}) = X'_{ij}\beta$$

Also the variance of each Y_{ij} depends on μ_{ij} :

$$Var(Y_{ij}) = \phi \mu_{ij}$$

In addition to these, an unstructured pairwise correlation pattern is assumed for the within-subject association among the repeated responses vector:

$$corr(Y_{ij}, Y_{ik}) = \alpha_{jk}$$

The vector of parameters α shows the pairwise correlation among responses.

The marginal model specified above is a log-

linear regression model, with an extra-poisson variance assumption (22).

So, to determine the relationship between air quality index, precipitation, air temperature and air humidity as environmental and climatic factors with the abundance of whiteflies this model was applied. The goodness of fit of model was evaluated by QIC.

Results

The density of whiteflies per trap in different seasons were calculated during the 12-month period of sampling in district No. 6 of Tehran in 2018. The most density of white flies per trap was 256.6 and 155.6 in early October and late September respectively due to low temperature and rainfall and high humidity (Fig. 2). The density of whiteflies per trap in other times of year was shown in Table 1. The number moved closer to zero from November to April. The population of whiteflies was inversely correlated with the level of air quality index ($p= 0.99$) and precipitation ($p= 0.95$), and it had a direct correlation with the high temperature. Also, the population of whiteflies had a direct correlation with the level of air humidity in the first half of the year, and it was inversely correlated in the final months of the year.

In the current study, up to the 13th whiteflies trapping (early October), using the marginal model, the estimates were calculated by the GEE method, and considering the Poisson link function, the following results were obtained:

A) For a unit of increase in air quality index (AQI), the population of whiteflies decreased by 0.98 (decreases).

B) For a unit of increase in temperature, the population of whiteflies increased by 1.18 (increases).

C) For a unit of increase in humidity, the population of whiteflies increased by 1.04 (increases).

D) For a unit of increase in precipitation, the population of whiteflies decreased by 0.99 (de-

creases) (Table 2). In this method, all four variables were significant.

In addition, the following results were obtained after the 13th measurement (early October):

A) For a unit of increase in air quality index (AQI), the population of whiteflies decreased by 0.98 (decreases).

B) For a unit of increase in temperature, the population of whiteflies increased by 2.37 (increases).

C) For a unit of increase in humidity, the population of whiteflies decreased by 0.92 (decreases).

D) For a unit of increase in precipitation, the population of whiteflies decreased by 0.44 (decreases) (Table 3). In this method, none of the four variables were significant. In both GEE models, QIC showed models were well-fitted.

Table 1. Comparison of Air Quality Index (AQI) with monthly activity of whiteflies, Tehran, Iran in 2018

Variable Month	Density of whitefly per trap	Air quality index (AQI)
Early April	00.00	71.52
Late April	00.00	71.68
Early May	22.55	75.53
Late May	24.45	79.56
Early June	24.70	82.93
Late June	24.75	94.31
Early July	24.55	93.67
Late July	25.45	87.81
Early August	34.00	89.47
Late August	50.00	90.75
Early September	127.8	102.73
Late September	155.6	100.68
Early October	256.6	88.00
Late October	99.25	84.47
Early November	00.00	93.06
Late November	00.00	128.12
Early December	00.00	136.15
Late December	00.00	138.14
Early January	00.00	142.28
Late January	00.00	145.85
Early February	00.00	121.45
Late February	00.00	112.15
Early March	00.00	110.59
Late March	00.00	108.14

Table 2. Generalized estimating equation (GEE) models predicting number of flies before 13th measurement, Tehran, Iran, 2018

Variable	Coefficient	S.E.	P- value
Air pollution	0.98	0.001	<0.001
Temperature	1.18	0.004	<0.001
Humidity	1.04	0.003	<0.001
Rain fall	0.99	0.000	<0.001
Constant	12.98	2.65	<0.001

Table 3. Generalized estimating equation (GEE) models predicting number of flies after 13th whiteflies trapping, Tehran, Iran in 2018

Variable	Coefficient	S.E.	P- value
Air pollution	0.98	7.40	0.99
Temperature	2.37	3.31	0.53
Humidity	0.92	<0.0001	-
Rain fall	0.44	6.35	0.95
Constant	1	556.34	1.00




Fig. 1. Geographic location of the study area in District 6, Municipality of Tehran, Iran




Fig. 2. Average monthly rainfall, temperature and humidity, Tehran, Iran in 2018




Fig. 3. Change point of whiteflies activity, Tehran, Iran, 2018

(N= number of collected whiteflies; Index= times of trapping)

Discussion

The results of this study showed that the population of the whiteflies was inversely correlated with the level of air quality index and precipitation, and it had a direct correlation with high temperature. Also, the population of whiteflies had a direct correlation with the level of air humidity in the first half of the year, and it was inversely correlated in the final months of the year.

Modern transportation and rapid trading of plants, cuttings, branches and other parts of plants, which often contain eggs, larvae and whitefly nymph, have led to the transfer of these pests into new places. However, there may be few whiteflies that enter a new place independently and without encountering the natural enemies with which they are evolving. Appropriate level of humidity and temperature is another factor for the developing population of whiteflies. The whiteflies were commonly able to complete their life cycle including egg, larvae, nymph and adult in the temperature of 15–35 °C while survival was usually reduced at temperatures <20 °C or >30 °C (23, 24). The rate of survival of the whiteflies at unfavorable low and

high temperatures were also affected by host plants (25). For instance, according to life span of mulberry trees in Tehran City, white flies strongly infest these trees (26) and when the leaves fall, the whiteflies population also decreases.

Importing of seedlings and cuttings of flowers, as well as the types of wood susceptible to the growth of whiteflies are one of the most important factors in the spread of whiteflies in different places. Whiteflies unable to fly long distances (27). Also, lack of precipitation is another factor in the survival of whiteflies. The high level of pollutants containing CO₂ in the air of Tehran, due to defective fuel consumption of worn out vehicles and other airborne pollution, is another important factor in the survival of whiteflies in urban areas since these flies live in conditions similar to greenhouse. It should be mentioned that, in fact, the high level of pollution and weather conditions are important factors in the reproduction of whiteflies in Tehran (28). The abundance of whiteflies in Tehran can cause health problems including itching, red and sore eyes, runny nose, allergies, and problems in the respiratory system of individuals, especially asthmatics (29). Children and people with poor immune system, the elderly populations and pregnant women are more susceptible to these problems. Also, the presence of whiteflies in food and beverages, besides causing fear, is concerning in terms of contamination with pathogenic microorganisms. Therefore, this study was conducted in different months of the year to consider the effects of temperature, precipitation, humidity and environmental contaminants. The materials on the wings and bodies of the whiteflies can act as the pollen and cause allergic reactions in individuals (30). Rain and low humidity (below 60%) and low or high temperatures disrupt development of the insects. Whiteflies stay in one place because they cannot migrate long distances. The longest reported displacement of this insect is 7km (31, 32). Whiteflies can also cause problems for the eyes and respiratory tracts (31). The reasons for eradica-

tion of the natural enemies of whiteflies include the destruction of their reproductive habitats, uncontrolled use of authorized and unauthorized pesticides, and the lack of using specialized biological controlling methods. These factors influence the growth of whiteflies in metropolitan cities like Tehran. Reducing the use of smoky and pollutant vehicles and monitoring the factories that produce vehicles with incomplete combustion (33, 34) is one of the major factors in making the environment unstable for whiteflies. The use of different least-hazardous pesticides to reduce the resistance of whiteflies, as well as the use of systemic toxins for non-productive trees (35), and reducing the use of insecticides as the only methods of controlling whiteflies, and ultimately the use of growth regulators and integrated pest management (36) will help control these pests in such cities as Tehran.

Conclusion

It should be mentioned that, in fact, the high level of pollution in different times of year and weather conditions are important factors in the reproduction of whiteflies in Tehran. According to these findings, during spring and summer from early May to early October that temperature and humidity for development of white flies are supplied in Tehran City, personal protection against these pests was recommended by Tehran residents.

Acknowledgements

This study was part of a postgraduate thesis in Medical Entomology and Vector Control, supported financially by Tehran University of Medical Sciences (Grant number: 29826). Ethical clearance was earned from the Institutional Ethics Committee of Tehran University of Medical Sciences. Authors thankful for the support of the Department of Medical Entomology and Vector Control sincerely. The authors declare that there is no conflict of interest.

References

1. Doukas D, Payne CC (2007) Greenhouse whitefly (Homoptera: Aleyrodidae) dispersal under different UV-light environments. *J Econ Entomol.* 100: 389–397.
2. Sadeh D, Nitzan N, Shachter A, Chaimovitch D, Dudai N, Ghanim M (2017) Whitefly attraction to rosemary (*Rosmarinus officinalis L.*) is associated with volatile composition and quantity. *PLoS One.* 12: e0177483.
3. De Barro PJ, Liu SS, Boykin LM, Dinsdale AB (2011) *Bemisia tabaci*: A statement of species status. *Annu Rev Entomol.* 56: 1–19.
4. Bogran CE, Heinz KM (2002) Whiteflies. Texas Agricultural Extension Service. p.7
5. Tracy JM (2011) Insect allergy. *Mt Sinai J Med.* 78: 773–783.
6. Polston JE, Anderson PK (1997) The emergence of whitefly-transmitted geminiviruses in tomato in the western hemisphere. *Plant Dis.* 81: 1358–1369.
7. Naveen NC, Chaubey R, Kumar D, Rebijith KB, Rajagopal R, Subrahmanyam B, Subramanian S (2017) Insecticide resistance status in the whitefly, *Bemisia tabaci* genetic groups Asia-I, Asia-II-1 and Asia-II-7 on the Indian subcontinent. *Sci Rep.* 18: 40634.
8. VanDoorn A, de Vries M, Kant MR, Schuurink RC (2015) Whiteflies glycosylate salicylic acid and secrete the conjugate via their honeydew. *J Chem Ecol.* 41(1): 52–58.
9. Fekrat L, Shishehbor P (2007) Some biological features of cotton whitefly, *bemisia tabaci* (Homoptera: Aleyrodidae) on various host plants. *Pakistan J Biol Sci.* 10: 3180–3184.
10. Hernández-Suárez E, Martin JH, Gill RJ, Bedford ID, Malumphy CP, Reyes Betancort JA, Carnero A (2012) The aleyrodidae (Hemiptera: Sternorrhyncha) of the Canary Islands with special reference to *Aleyrodes*, *Siphoninus*, and the challenges of puparial morphology in *Bemisia*. *Zootaxa.* 3212: 1–76.

11. Alemansoor H, Fallahzadeh, M (2004) Bio ecology of ash whitefly, *Siphoninus phillyreae* (Halliday) (Hom: Aleyrodidae) in the Fars Province, Iran. *Pajouhesh and Sazandegi*. 62: 64–70.
12. Shahbazvar N, Sahragard A, Manzari SH, Hosseini R, Hajizadeh J (2010) A faunal study of whiteflies (Hemiptera: Aleyrodidae) and their parasitoids in Guilan Province, Iran. *Entomofauna*. 17: 269–284.
13. Ghahari H, Abd-Rabou S, Ostovan H, Samin N (2007) Whiteflies (Homoptera: Aleyrodidae) and their host plants in Golestan Province, Iran. *Plant and Ecosystem*. 12: 17–28.
14. Ghahari H, Hatami B (2000) Morphological and biological studies of greenhouse whitefly *Trialeurodes vaporariorum* westwood (Homoptera: Aleyrodidae) in Isfahan. *J Water Soil Sci*. 4: 141–154.
15. Antignus Y, Nestel D, Cohen S, Lapidot M (2001) Ultraviolet-deficient greenhouse environment affects whitefly attraction and flight-behavior. *Environ Entomol*. 30: 394–399.
16. Heshmati GA (2007) Vegetation characteristics of four ecological zones of Iran. *Int J Plant Prod*. 2: 215–224.
17. Iran Meteorological Organization. Iran (2019) current weather. Available at: <http://www.irimo.ir/index.php?newlang=eng>.
18. Lu Y, Bei Y, Zhang J (2012) Are yellow sticky traps an effective method for control of sweetpotato whitefly, *Bemisia tabaci*, in the greenhouse or field? *J Insect Sci*. 12: 113.
19. Chen J, Gupta AK (2011) Parametric statistical change point analysis: with applications to genetics, medicine, and finance: Springer Science and Business Media. p. 120.
20. Chen J, Gupta AK, Pan J (2006) Information Criterion and Change Point Problem for Regular Models. *Indian J Statistics*. 68: 252–282.
21. Cui J, Qian G (2007) Selection of working correlation structure and best model in GEE analyses of longitudinal data. *Commun Stat-Simul Comput* 36: 987–996.
22. Fitzmaurice GM, Laird NM, Ware JH (2011) *Applied Longitudinal Analysis*, 2nd Edition. p. 752.
23. Bonato O, Lurette A, Vidal C, Fargues J (2007) Modelling temperature-dependent bionomics of *Bemisia tabaci* (Q-biotype). *Physiol. Entomol*. 32: 50–55.
24. Muniz M, Nombela G (2001) Differential variation in development of the B- and Q-biotypes of *Bemisia tabaci* (Homoptera: Aleyrodidae) on sweet pepper at constant temperatures. *Environ. Entomol*. 30: 720–727.
25. Tsueda H, Tsuchida K (2011) Reproductive differences between Q and B whiteflies, *Bemisia tabaci*, on three host plants and negative interactions in mixed cohorts. *Entomol Exp Appl*. 141: 197–207.
26. David V, Ragupathy E (2004) Whiteflies (homoptera: Aleyrodidae) of Mulberry, *Morus alba* L., in India. *Pestology*. 28(10): 24–33.
27. Mound, LA, Halsey SH (1978) *Whitefly of the World*. British Museum (Natural History)/ John Wiley and Sons, Chichester, p. 340.
28. Alomar O, Goula M, Albajes R (2001) Colonization of tomato fields by predatory mirid bugs (Hemiptera: Heteroptera) in northern Spain. *Agriculture, Ecosystems and Environment*. 89: 105–115.
29. Grier TJ, LeFevre DM, Duncan EA, Esch RE, Coyne TC (2012) Allergen stabilities and compatibilities in mixtures of high-protease fungal and insect extracts. *Ann Allergy Asthma Immunol*. 108: 439–447.
30. Traidl-Hoffmann C, Kasche A, Menzel A, Jakob T, Thiel M, Ring J, Behrendt H (2003) Impact of pollen on human health: more than allergen carriers? *Int Arch Allergy Immunol*. 131: 1–13.
31. Cohen S, Melamed-Madjar V (1978) Prevention by soil mulching of spread of tomato yellow leaf curl virus transmitted by

- Bemisia tabaci* (Gennadius) (Hemiptera: Aleyrodidae) in Israel. Bulletin Entomol Res. 68: 465–470.
32. Cohen S, Berlinger M (1986) Transmission and cultural control of whitefly-borne viruses. Agric Ecosystems Environ. 17: 89–97.
 33. Burnett T (1949) The effect of temperature on an insect host-parasite population. Ecology. 30: 113–134.
 34. Crowder DW, Ellsworth PC, Tabashnik BE, Carrière Y (2008) Effects of operational and environmental factors on evolution of resistance to pyriproxyfen in the sweetpotato whitefly (Hemiptera: Aleyrodidae). Environ Entomol. 37: 1514–1524.
 35. Vosman B (2016) Different mechanisms of whitefly resistance in cabbage and its wild relatives. plant and animal genome XXIV Conference, Plant and Animal Genome.
 36. Chandler D, Bailey AS, Tatchell GM, Davidson G, Greaves J, Grant WP (2011) The development, regulation and use of biopesticides for integrated pest management. Philos Trans R Soc Lond B Biol Sci. 366: 1987–1998.