



Original Article

Journal of Food Safety and Hygiene

Journal homepage: <http://jfsht.tums.ac.ir>

Antibiotic susceptibility pattern of *Staphylococcus aureus* isolated from fresh lettuce (*Lactuca sativa*) and food safety knowledge of vegetable vendors in Birnin Kebbi, Nigeria

Shamsudeen Muhammad Muhammad*, Adamu Almustapha Aliero

Department of Microbiology, Faculty of Life Science, Kebbi State University of Science and Technology, Aliero, Nigeria.

ARTICLE INFO

Article history:

Received 19 Jun. 2023

Received in revised form

23 Aug. 2023

Accepted 02 Sep. 2023

Keywords:

Lettuce (*Lactuca sativa*);*Staphylococcus aureus*;

Antibiotic susceptibility;

Food safety

ABSTRACT

In recent years, the consumption of fresh products has increased due to the multiple contributions of nutrients and functional properties. This study aims to determine the susceptibility pattern of *Staphylococcus aureus* in fresh lettuce and assess the food safety practice of vegetable vendors in the Birnin Kebbi metropolis. A total of 28 fresh lettuce samples were collected from various locations (Tudun Wada, Rafin Atiku, Badariya, and Bayan Kara) in the Birnin Kebbi metropolis, chosen randomly from vegetable vendors. A structured questionnaire was utilized to assess food safety practices. *S. aureus* was isolated and identified using cultural and biochemical characteristics and antibiotic sensitivity was conducted by the Kirby-Baur disk diffusion method. Only 7.14% of the vegetable vendors reported using gloves while handling vegetables. 17.86% reported washing their hands properly after handling waste or garbage. Only 7.14% reported wearing an apron while working. 50% reported properly cleaning the vegetable storage area before storing new products. 14.29% reported washing vegetables after purchasing them from the market, while None of the respondents reported covering their products while selling them. Out of a total of 28 samples tested, 21(75%) were found to be positive for *S. aureus*. Antibiotic susceptibility pattern of *S. aureus* showed resistance in Ampicillin (100%), Amoxicillin (28.57%), Tetracycline (17.86%), Rifampicin (39.29%), Chloramphenicol (14.29%), Neomycin (53.57%), Methicillin (89.29%), Cefpodoxime (35.71%) respectively. Furthermore, none of the *S. aureus* were resistant to Gentamycin. The high occurrence of *S. aureus* found in fresh lettuce suggests contamination could have occurred before or after harvesting. It is recommended that cleaning and sanitation practices be introduced to enhance the safety and quality of this vegetable.

Citation Muhammad ShM, Aliero AA. Antibiotic susceptibility pattern of *Staphylococcus aureus* isolated from fresh lettuce (*Lactuca sativa*) and food safety knowledge of vegetable vendors in Birnin Kebbi, Nigeria. J food safe & hyg 2023; 9(3): 185-196

1. Introduction

Fruits and vegetables are considered to be crucial elements of a healthy diet due to their low-fat and low-energy density.

They are also relatively high in essential vitamins, minerals, and other beneficial bioactive compounds. Additionally, they are a valuable source of dietary fiber (1). Leafy greens like lettuce, spinach, and kale are commonly eaten raw or lightly cooked, making them susceptible to potential health risks from harmful

*Corresponding author. Tel.: +2348166923586

E-mail address: deenshams2000@gmail.com



Copyright © 2023 Tehran University of Medical Sciences. Published by Tehran University of Medical Sciences.

This work is licensed under a Creative Commons Attribution-NonCommercial 4.0 International license (<https://creativecommons.org/licenses/by-nc/4.0/>).

Non-commercial uses of the work are permitted, provided the original work is properly cited.

bacteria or pathogens on their leaves. Additionally, being grown close to the ground increases the chances of contamination from soil and other environmental sources, further emphasizing the need for proper handling and washing before consumption (2).

Lettuce is an annual leafy vegetable belonging to the Asteraceae family, commonly consumed raw in the form of salad. Indeed, lettuce is susceptible to contamination due to various factors, including inadequate sanitation practices, poor drainage systems, and unhygienic planting, harvesting, packaging, transportation, and storage methods that are sometimes employed in its production and distribution (3).

Additionally, lettuce can become contaminated with pathogenic microorganisms at various stages of the food production process, from the farm to the market. One common source of contamination is the use of cattle manure as fertilizer or the proximity of vegetable fields or water sources to cattle farms. These sources have been linked to instances of vegetable contamination in the past (4).

Not adequately washing and peeling food items can serve as a breeding ground for different microorganisms, potentially resulting in foodborne illnesses. These illnesses can be caused by different types of pathogens that enter the human body through the oral route, such as by eating contaminated food or drinking contaminated water (5).

While raw fruits and vegetables are known for their numerous health benefits, research has indicated that they rank as the second most common source of foodborne illness outbreaks. These outbreaks not only present a public health risk but also lead to substantial

economic losses for the food industry (6). Both in developed and developing regions, foodborne pathogens are a major cause of infectious disease outbreaks, and incidents of foodborne illness related to the consumption of fresh produce have seen a significant increase in recent years (7).

Fresh vegetables can become contaminated by microbes at many points along the farm-to-fork supply chain. Fresh produce cultivation, harvest, preparation, washing, transportation to stores, distribution chains, and even the final stage in the consumer's kitchen can all result in contamination. Both contaminated soil and water might result in contamination during agriculture. Food contamination may also be influenced by the environment in which these foods are handled, such as food contact surfaces or processing equipment (8).

Staphylococcus aureus is a common bacterium found in 30% of healthy individuals and can cause infections ranging from mild to severe (9). It has various virulent genes that encode toxins and enzymes. Humans are the primary reservoir, and food contamination can occur through direct or indirect contact. In recent years, there has been a notable surge in bacterial resistance to antibiotics (10). This rise in antimicrobial drug resistance among foodborne pathogens is primarily linked to the overuse of antibiotics in livestock, aquaculture, and agricultural practices for both preventive and therapeutic reasons (11). It is crucial to note that antibiotic-resistant bacteria and antimicrobial resistance genes can transfer between animal and human populations through various routes, including direct contact with animals or their environment and indirect contact through contaminated food sources (12). Staphylococcal enterotoxins, which are the cause

of staphylococcal food poisoning, can develop and be produced by *S. aureus*. In outbreaks of foodborne illness, pathogenic *Staphylococcus* strains' enterotoxins are frequently involved. Food contamination with an enterotoxigenic strain of *Staphylococcus* spp. is the primary precondition for staphylococcal food poisoning (13). Temperature, pH, and water activity requirements must be met for enterotoxins to be created. Upon being produced by the strains, staphylococcal enterotoxins are resistant to a variety of environmental conditions, such as freezing, drying, heat treatment, and low pH conditions, which easily destroy the enterotoxin-producing strains themselves (14).

S. aureus is a bacterial species known for its ability to develop resistance to antibiotics. The β -lactam antibiotics, which include penicillin, methicillin, flucloxacillin, dicloxacillin, nafcillin, oxacillin, and cloxacillin, are typically effective in treating infections caused by *S. aureus*. However, some strains of *S. aureus* have acquired the *mecA* gene, leading to methicillin-resistant *S. aureus* (MRSA), which is resistant to all penicillins, cephalosporins, and carbapenems. The widespread distribution of MRSA worldwide and its ability to acquire resistance to multiple antibiotics have raised concerns about treatment options (15).

Nigeria is a significant producer of fruits and vegetables, contributing to the livelihoods of both rural and urban communities. Leafy vegetables are commonly used for cooking to improve the taste of the soup and for nutritional purposes, medicinal properties and are used to manage ailments such as diarrhea, stomach issues, coughing, and malaria (16). Typically, these vegetables are cooked in soups or stews and

served with a carbohydrate-rich staple food (17). Regrettably, in Nigeria, fruits and vegetables are often unclean and not properly sanitized before being sold in the market, leading to unsafe and substandard produce. There has not been a comprehensive investigation on the issue of antibiotic resistance in fresh lettuce in Birnin Kebbi. Additionally, there is a lack of data regarding comprehensive the prevalence of antibiotic-resistant *Staphylococcus aureus* and food safety knowledge level among street vendors that sell fresh produce in the Birnin Kebbi metropolis. Understanding the prevalence and antibiotic resistance patterns of *Staphylococcus aureus* in fresh lettuce and the level of knowledge among street traders is essential to ensure the safety of the food supply and to prevent the spread of infections. Thus, this research aims to establish the susceptibility of *Staphylococcus aureus* in fresh lettuce to commonly available antibiotics and evaluate the food safety knowledge of vegetable vendors in the Birnin Kebbi metropolis.

2. Materials and Methods

2.1. Collection of fresh lettuce

A total of 28 fresh lettuce samples were collected from various locations (Tudun Wada, Rafin Atiku, Badariya, and Bayan Kara) in the Birnin Kebbi metropolis, chosen at random from vendors who display their vegetables openly for customers to see and purchase. Lettuce samples were obtained from each vendor using the same approach as a regular customer, and then individually placed in sterile plastic bags. These samples were subsequently transported in a cool box at low temperatures to the Microbiology Laboratory of

Kebbi State University of Science and Technology Aliero for further analysis.

2.2. Food safety practice data collection

To assess the demographic characteristics and food safety practices of vegetable vendors, a structured questionnaire was utilized. Trained research assistants and the researchers themselves conducted interviews with the participants, using the structured questionnaire which included questions related to the demographic information of the respondents adapted from previous works (18,19). Moreover, questions regarding food safety practices were adapted from previously published studies (20) and were presented on a five-point scale, with a score of 1 representing strong disagreement and a score of 5 representing strong agreement.

2.3. Samples preparation

To prepare each fresh lettuce sample for analysis, the initial step involved slicing or cutting it into small pieces using a sterilized tray and knife while wearing latex gloves for sanitary purposes. Subsequently, 25 g of each sample were weighed and combined with 225 mL of sterile normal saline solution. The mixture was homogenized using a stomacher device and then incubated at 37°C for a duration of 24 h.

2.4. Isolation and identification of *S. aureus*

Following homogenization and incubation, one milliliter of each sample was extracted and inoculated onto a sterile Mannitol Salt Agar (MSA) plate, which was subsequently streaked and incubated at 37°C for 24 h. The typical colonies obtained from each MSA agar plate were further sub-cultured on nutrient agar to facilitate identification.

From the total of 28 samples, one presumptive culture of *Staphylococcus aureus* was randomly chosen and identified using culture characteristics on Mannitol Salt Agar, gram-staining, and biochemical reactions following the guidelines outlined in Bergey's Manual of Systematic Bacteriology (21).

2.5. Antibiotic sensitivity testing

The confirmed *S. aureus* isolates underwent antibiotic susceptibility testing using the Kirby Bauer disk diffusion technique, following the guidelines provided by the Clinical and Laboratory Standard Institute (CLSI). Eight different antibiotics were utilized in this process, namely ampicillin (10 µg), amoxicillin (30 µg), tetracycline (30 µg), gentamycin (10 µg), rifampicin (5 µg), chloramphenicol (30 µg), neomycin (10 µg), methicillin (5 µg), and cefpodoxime (10 µg). The inhibition zone diameters were measured and compared to predetermined breakpoints outlined in the CLSI protocol (22). Based on the results, the antibiotic susceptibility of the isolates was interpreted as sensitive, intermediate, or resistant.

3. Results

3.1. Prevalence of *S. aureus* in fresh lettuce samples

This study collected fresh lettuce samples (n=28) from street vegetable vendors in Birnin kebbi. Table 1 presents the number of positive samples for *S. aureus* along with their prevalence percentage. Out of a total of 28 samples tested, 21(75%) were found to be positive for *S. aureus*, while the remaining 7(25%) were negative.

Table 1. Prevalence of *S. aureus* (n=28) in fresh lettuce samples

Organism	Number (%) positive	Number (%) negative
<i>Staphylococcus aureus</i>	21(75)	7(25)

Table 2. Demographic characteristics of vegetable vendors

Parameter	Frequency	Percentage
Gender		
Male	28	100
Female	0	0
Age		
<18	4	14.23
19-40	20	71.43
41-60	4	14.23
>60	0	0
Education		
Primary	18	64.29
Secondary	5	17.86
Higher	0	0
No formal education	5	17.86
Number of years in business		
0-5	5	17.86
6-10	15	53.57
>10	8	28.57
Marital status		
Single	20	71.43
Married	8	28.57
Divorced	0	0
Food safety training		
Yes	0	0
No	28	100

3.2. Demographic characteristics of vegetable vendors

The socio-demographic profile of respondents is presented in Table 2. Out of 28 respondents, all 100% were males. The majority of the participants are aged between 19-40 (71.43%), followed by those aged below 18 (14.23%) and those aged 41-60 (14.23%). In terms of education, 64.29% of the participants have completed primary education, while 17.86% have had no formal education, and only 17.86% have completed secondary education. Regarding the number of years in business, 53.57% of the participants have been in business for 6-10 years, while 28.57% have been in business for more than 10 years, and 17.86% have been in business for 0-5 years. In terms of marital status, 71.43% of the participants are single, while 28.57% are married. Finally, none of the participants received food safety training.

3.3. Food safety practices of vegetable vendors

Table 3 represents the responses (in percentages) to various food safety questions. All 28 respondents (100%) reported eating or drinking at their workplace. Only 7.14% reported using gloves while handling vegetables, while 71.23% reported not using gloves. 17.86% reported washing their hands properly after handling waste or garbage, while 82.12% reported not doing so. All 28 respondents (100%) reported washing their hands after using the toilet. Only 7.14% reported wearing an apron while working, while 92.86% reported not wearing one. 50% reported properly cleaning the vegetable storage area before storing new products, while the other 50% did not.

All 28 respondents (100%) reported using clean polyethylene bags for packaging. 14.29% reported washing vegetables after purchasing them from the market, while 85.71% reported not doing so. None of the respondents reported covering their products while selling them.

3.4. Results of antibiotic susceptibility pattern for *S. aureus* isolates from fresh lettuce

The antibiotic-resistant pattern of *S. aureus* from lettuce samples is shown in Table 4. All 28 isolates (100%) were resistant to ampicillin. The majority of isolates (28.57%) were resistant to amoxicillin. 17.86% of the isolates were resistant to tetracycline. None of the isolates were resistant to gentamycin. A significant proportion of isolates (39.29%) were resistant to rifampicin. 14.29% of the isolates were resistant to chloramphenicol. More than half of the isolates (53.57%) were resistant to neomycin. The vast majority of isolates (89.29%) were resistant to methicillin. 35.71% of the isolates were resistant to cefpodoxime. Overall, the resistance pattern of the *S. aureus* isolates indicates that there is significant resistance to several antibiotics, particularly ampicillin, methicillin, and neomycin.

Table 3. Food safety practices of vegetable vendors

Questions	Response (n)%	
	YES	NO
Do you eat or drink at the workplace?	28(100)	0(0)
Do you use gloves while handling vegetables?	2(7.14)	20(71.23)
Do you wash your hands properly after handling waste/garbage?	5(17.86)	23(82.12)
Do you wash your hand after using the toilets?	28(100)	0(0)
Do you wear an apron the whole working?	2(7.14)	26(92.86)
Do you properly clean the vegetable storage area before storing new products?	14(50)	14(50)
Do you use clean polyethylene bags for packaging?	28(100)	0(0)
Do you wash the vegetables after purchase from the market?	4(14.29)	24(85.71)
Do you cover your product while selling?	0(0)	28(100)

Table 4. Antibiotic susceptibility pattern for *S. aureus* (n=28) isolates from fresh lettuce

Antibiotics	Concentration (ug)	Zones of inhibition (mm)					
		Number (%) of resistant		Number (%) of intermediate		Number (%) of sensitive	
Ampicillin	(10 µg),	28(100)		0(0)		0(0)	
Amoxicillin	(30 µg),	8(28.57)		6(21.43)		14(50)	
Tetracycline	(30 µg),	5(17.86)		2(7.14)		21(75)	
Gentamycin	(10 µg),	0(0)		5(17.86)		23(82.14)	
Rifampicin	(5 µg),	11(39.29)		1(3.57)		16(57.14)	
Chloramphenicol	(30 µg)	4(14.29)		3(10.71)		21(75)	
Neomycin	(10 µg),	15(53.57)		0(0)		13(46.43)	
Methicillin	(5 ug)	25(89.29)		2(7.14)		1(3.57)	
Cefpodoxime	(10 µg).	10(35.71)		4(14.29)		14(50)	




Figure 1. Antibiotic susceptibility test for *S. aureus* (n=28) isolates from fresh lettuce

4. Discussion

This study presents novel findings regarding the occurrence of antibiotic-resistant *Staphylococcus aureus* in lettuce, as well as the food safety measures implemented by vegetable vendors in the city of Birnin Kebbi, Nigeria. Furthermore, it establishes a foundation of knowledge that can serve as a benchmark for future studies in this area. Fresh produce can become contaminated at different points along the entire production chain, from its cultivation on the farm to its consumption at the table (23). Various factors, including human contact, harvesting equipment, transportation containers, insects, dust, rinse water, ice, vehicles for transport, and processing equipment, can all play a role in postharvest contamination (24).

In this study, only 50% reported properly cleaning the vegetable storage area before storing new products, 14.29% reported washing vegetables after purchasing

them from the market and none of the respondents reported covering their products while selling them. This could be contributing factor to the contamination of the vegetable. According to previous research, an inadequate understanding of food safety can result in food handlers adopting unsatisfactory hygiene practices (25). Additional research has indicated that simply providing food safety and hygiene knowledge does not guarantee the adoption of safe food behavior or practices (26).

The prevalence of *Staphylococcus aureus* contamination reported in this study is comparatively higher than rates observed in other regions. For instance, a study conducted in Tehran found that 45 out of 350 (12.85%) vegetable and salad samples tested positive for *S. aureus* (27). In a study conducted by (28), investigating antibiotic-resistant *S. aureus* in leafy greens and clinical sources, it was found that 31 (28.18%) of the retail

vegetables tested positive for *S. aureus*. Among the vegetables, lettuce had the highest prevalence, with 13 out of 15 samples (86.7%) being positive for *S. aureus*. Another study in the Bekaa Valley of Lebanon revealed that lettuce samples had a significantly higher microbial load, including coliforms, *E. coli*, and *S. aureus*, compared to parsley samples (29). Similarly, a study conducted in China to investigate the prevalence and characterization of *S. aureus* in retail vegetables found that 24 (5.73%) samples were positive for *S. aureus*, with lettuce having the highest prevalence at 13 out of 84 samples (15.48%) (30). *Staphylococcus aureus* has the potential to contaminate vegetables when there is contact with unwashed hands during the selection process. Its presence in food raises public health concerns due to its ability to produce enterotoxins, posing a risk of food poisoning (31,32). Recently, there has been a significant alarm in both public and clinical sectors regarding the emergence and spread of antibiotic-resistant strains of *S. aureus*, which has implications for managing staphylococcal infections (33,34). Multiple studies have documented the presence of antibiotic-resistant *S. aureus* strains in various food samples from different countries (28,35,36).

In this particular study, it was found that all of the isolates (100%) demonstrated resistance to ampicillin, while 89.29% exhibited resistance to penicillin, (53.57%) were resistant to neomycin and the vast majority of isolates (89.29%) were resistant to methicillin. MRSA, methicillin-resistant *Staphylococcus aureus*, is notorious for its resistance not only to methicillin and oxacillin but also to various other antibiotics. This bacterium has emerged as a significant cause of infections, posing a major global health threat due to its ability to produce

numerous toxins. In recent years, MRSA has been increasingly detected in the community, outside the confines of healthcare facilities. This includes its presence in food and food-producing animals worldwide (37). The presence of MRSA in food raises further awareness about potential routes of transmission and the need for monitoring and controlling antibiotic resistance in both clinical and food environments.

In this particular study, it was found that all of the isolates (100%) demonstrated resistance to ampicillin, while 89.29% exhibited resistance to penicillin. These resistance rates were in agreement with those reported by (38) in their research on leaf vegetables in Korea, where a resistance rate of 96.3% was observed. The resistance patterns to other antimicrobials such as tetracycline, erythromycin, gentamicin, and ciprofloxacin were found to be comparable to the findings of (30). In their study, it was reported that 49.4% of *S. aureus* isolates from food products displayed resistance to tetracycline, 24.1% to erythromycin, and 13.8% to gentamicin. Over time, there have been changes in antibiotics and the resistance patterns exhibited by foodborne pathogens. These patterns can vary among isolates from the same or different sources. Several factors contribute to these differences, including geographical locations, the specific bacterial species involved, the type of animal production systems employed, the extent of antibiotic usage, the techniques used for sampling, and the duration of the sampling period. These diverse factors collectively influence the observed variations and similarities in antibiotic resistance patterns among foodborne pathogens (39,40).

5. Conclusion

The findings of this study demonstrate the presence of resistant *Staphylococcus aureus* isolates in lettuce samples, indicating inadequate sanitation and improper handling practices during harvesting and marketing. To mitigate contamination risks, farmers need to adopt the best agricultural and handling practices, while food vendors, processors, and consumers should adhere to good hygiene practices. These results also provide valuable information to clinicians for effective treatment of patients. Therefore, it is crucial for the government to closely monitor both pre-and post-harvest activities of vegetable producers and sellers to minimize disease risks. Additionally, regular surveillance of antibiotic susceptibility is necessary to monitor changes in resistance patterns.

Conflict of interest

The authors declare that there is no conflict of interest.

Acknowledgment

This study was supported by Kebbi State University of Science and Technology, Nigeria.

References

1. Brookie KL, Best GI, Conner TS. Intake of raw fruits and vegetables is associated with better mental health than intake of processed fruits and vegetables. *Front Psychol* 2018; 9: 487.
2. Mohamed MA, Siddig EE, Elaagip AH, et al. Parasitic contamination of fresh vegetables sold at central markets in Khartoum state, Sudan. *Ann Clin Microbiol Antimicrob* 2016; 15: 1–7.
3. Bekele F, Tefera T, Biresaw G, et al. Parasitic contamination of raw vegetables and fruits collected from selected local markets in Arba Minch town, Southern Ethiopia. *Infect Dis Poverty* 2017; 6: 1–7.
4. Castro-Ibáñez I, Gil MI, Tudela JA, et al. Microbial safety considerations of flooding in primary production of leafy greens: a case study. *Food Res Int* 2015; 68: 62–69.
5. Amoah P, Drechsel P, Abaidoo RC, et al. Effectiveness of common and improved sanitary washing methods in selected cities of West Africa for the reduction of coliform bacteria and helminth eggs on vegetables. *Trop Med Int Heal* 2007; 12: 40–50.
6. Hussain MA, Dawson CO. Economic impact of food safety outbreaks on food businesses. *Foods* 2013; 2: 585–89.
7. Stine SW, Song I, Choi CY, et al. Application of pesticide sprays to fresh produce: a risk assessment for hepatitis A and Salmonella. *Food Environ Virol* 2011; 3: 86–91.
8. Machado-Moreira B, Richards K, Brennan F, et al. Microbial contamination of fresh produce: what, where, and how? *Compr Rev Food Sci Food Saf* 2019; 18: 1727–50.
9. Taylor TA, Unakal CG. *Staphylococcus aureus*. *Stat Pearls* 2022; 2: 8-15.
10. Stanton IC, Bethel A, Leonard AFC, et al. What is the research evidence for antibiotic resistance exposure and transmission to humans from the environment? A systematic map protocol. *Environ Evid* 2020; 9: 1–8.
11. Al Amin M, Hoque MN, Siddiki AZ, et al. Antimicrobial resistance situation in animal health of Bangladesh. *Vet World* 2020; 13: 2713.
12. Faour-Klingbeil D, Murtada M, Kuri V, et al. Understanding the routes of contamination of ready-to-eat vegetables in the Middle East. *Food Control* 2016; 62: 125–33.
13. Grisoldi L, Karama LM, Armani A, et al. *Staphylococcus aureus* enterotoxin in food of animal

- origin and staphylococcal food poisoning risk assessment from farm to table. *Italian J Anim Sci* 2021; 20: 677–90.
14. Hennekinne JA, De Buyser ML, Dragacci S. *Staphylococcus aureus* and its food poisoning toxins: characterization and outbreak investigation. *FEMS Microbiol Rev* 2012; 36: 815–36.
 15. Andreoletti O, Buhka H, Buncic S, et al. Assessment of the Public Health significance of methicillin-resistant *Staphylococcus aureus* (MRSA) in animals and foods scientific opinion of the panel on biological hazards. *EFSA J* 2009; 993: 1–73.
 16. Bakari UM, Usman M. Marketing of some selected vegetables: In Yola-North and South local government areas of Adamawa state, Nigeria. *Int J Eng Sci* 2013; 2: 11.
 17. Lawal OM, Idowu-Adebayo F, Enujiugha VN. Nutritional assessment of Nigerian ethnic vegetable soups (Marugbo, Tete, and Ila). *J Nutr Food Lipid Sci* 2018; 1: 32–39.
 18. Mama M, Alemu G. Prevalence, antimicrobial susceptibility patterns and associated risk factors of *Shigella* and *Salmonella* among food handlers in Arba Minch University, South Ethiopia. *BMC Infect Dis* 2016; 16: 1–7.
 19. Tessema AG, Gelaye KA, Chercos DH. Factors affecting food handling practices among food handlers of Dangila town food and drink establishments, North West Ethiopia. *BMC Pub Health* 2014; 14: 1–5.
 20. Akabanda F, Hlortsi EH, Owusu-Kwarteng J. Food safety knowledge, attitudes and practices of institutional food-handlers in Ghana. *BMC Pub Health* 2017; 17: 1–9.
 21. Holt JG, Krieg NR, Sneath PHA, et al. 1994. *Bergey's Manual of determinate bacteriology*.
 22. Clinical and Laboratory Standard Institute (CLSI). 2021. Performance standards for antimicrobial susceptibility testing, 30th ed. West Valley. 345-411 p.
 23. Ailes EC, Israel K. Microbial concentrations on fresh produce are affected by postharvest processing, importation and season. *J Food Prot* 2008; 71: 2389–97.
 24. Beuchat LR. Ecological factors influencing survival and growth of human pathogens on raw fruits and vegetables. *Microbes Infect* 2002; 4: 413–23.
 25. Giritlioglu I, Batman O, Tetik N. The knowledge and practice of food safety and hygiene of cookery students in Turkey. *Food Control* 2011; 22: 838–42.
 26. Al-Shabib NA, Mosilhey SH, Husain FM. Cross-sectional study on food safety knowledge, attitude and practices of male food handlers employed in restaurants of King Saud University, Saudi Arabia. *Food Control* 2016; 59: 212–17.
 27. Mir SA, Shah MA, Mir MM, et al. Microbiological contamination of ready-to-eat vegetable salads in developing countries and potential solutions in the supply chain to control microbial pathogens. *Food Control* 2018; 85: 235–44.
 28. Yigrem C, Eribo B, Sangre K. Incidence of antibiotic resistant *Staphylococcus aureus* in leafy greens and clinical sources. *J Microbiol Res* 2022; 12: 13–23.
 29. Halablal M, Sheet I, Holail H. Microbiological quality of raw vegetables grown in Bekaa Valley Lebanon. *Americ J Food Technol* 2011; 6: 129–39.
 30. Wu S, Huang J, Qingping W, et al. Prevalence and characterization of *Staphylococcus aureus* isolated from retail vegetables in China. *Front Microbiol* 2018; 9: 1–10.
 31. Eni AO, Oluwawemitan IA, Oranusi SU. Microbial quality of fruits and vegetables sold in Sango Ota, Nigeria. *Afric J Food Sci* 2010; 4: 291–96.
 32. Mossel DAA, Corry JEL, Struijk CB, et al. 1995. *Essentials of the microbiology of foods: a textbook for advanced studies*. 1st ed. John Wiley & Sons. 123 p.
 33. Shankar PR. The evolving threat of antimicrobial resistance. Options for action. *Australas Med J* 2012; 5:

- 243.
34. Thomas S, Espeisse O. Antimicrobial resistance. *New Engl J Med* 2020; 17: 330–33.
 35. Wang W, Zoolagnia B, Tao J, et al. Enterotoxigenicity and antimicrobial resistance of *Staphylococcus aureus* isolated from retail food in China. *Front Microbiol* 2017; 8: 2256.
 36. Degaga B, Sebsibe I, Belete T, et al. Microbial quality and safety of raw vegetables of Fiche Town, Oromia, Ethiopia. *J Environ Public Health* 2022; 10: 11-25.
 37. Lee AS, De Lencastre H, Garau J, et al. Methicillin-resistant *Staphylococcus aureus*. *Nat Rev Dis Prim* 2018; 4: 1–23.
 38. Hong J, Kim Y, Kim J, et al. Genetic diversity and antibiotic resistance patterns of *Staphylococcus aureus* isolated from leaf vegetables in Korea. *J Food Sci* 2015; 80: M1526–M1531.
 39. Yelin I, Kishony R. Antibiotic resistance. *Cell* 2018; 172: 1136.
 40. Larsson DGJ, Flach CF. Antibiotic resistance in the environment. *Nat Rev Microbiol* 2022; 20: 257–69.