



Detection of Mandibular Incisive Canal Using CBCT and its Anatomical Relationships in Different Nationalities: A Review Study

Paniz Ranji¹, Hamed Mortazavi², Yaser Safi¹, Solmaz Valizadeh¹, Amirali Shirian³ and Katayoun Ghaffari^{1*}

1. Department of Oral and Maxillofacial Radiology, School of Dentistry, Shahid Beheshti University of Medical Sciences, Tehran, Iran

2. Department of Oral Medicine, School of Dentistry, Shahid Beheshti University of Medical Sciences, Tehran, Iran

3. Department of Prosthodontics, Dental branch, Islamic Azad University of Medical Sciences, Tehran, Iran

* Corresponding author

Katayoun Ghaffari, DDS

Department of Oral and Maxillofacial Radiology, School of Dentistry, Shahid Beheshti University of Medical Sciences, Tehran, Iran

Tel: +98 9122955178

Email: Katayoon.ghaffari@gmail.com

Received: 28 Aug 2025

Accepted: 25 Sept 2025

Citation to this article

Ranji P, Mortazavi H, Safi Y, Valizadeh S, Shirian AA, Ghaffari K. Detection of Mandibular Incisive Canal Using CBCT and its Anatomical Relationships in Different Nationalities: A Review Study. *J Iran Med Council.* 2026;9(3):600-11.

Abstract

Background: The Mandibular Incisive Canal (MIC) is the mesial extension of the inferior alveolar nerve, frequently encountered during surgical procedures such as implant placement and bone graft harvesting. Thorough radiographic evaluation, particularly using three-dimensional imaging modalities like Cone-Beam Computed Tomography (CBCT), is crucial to prevent nerve injury and related complications. This review summarizes studies that investigated MIC detection using CBCT.

Methods: Following the PRISMA guidelines, an electronic search was performed in PubMed/MEDLINE and Scopus, identifying 45 studies, of which 35 met the inclusion criteria. An electronic search was conducted in August 2023 using PubMed/MEDLINE and Scopus with MeSH terms and keywords: “mandibular incisive canal,” “incisive canal,” “computed tomography, cone beam,” “CBCT,” “inferior alveolar nerve,” and “mental foramen.” Relevant studies published in English up to August 28, 2023, were included.

Results: MIC was detected in 87.11% of the cases examined by CBCT. The canal was more prevalent in females (F:M = 1.12:1), with a mean patient age of 45.11 years (range 10-88). The average MIC length and width were 11.30 mm and 1.72 mm, respectively. The mean distances from the MIC to the buccal cortex, lingual cortex, inferior mandibular border, and alveolar crest were 3.71, 4.98, 9.37, and 15.76 mm, respectively. The longest and widest MICs were observed in the Chinese population, whereas the smallest dimensions were found in the Brazilian population. MIC prevalence was highest among Malays (100%) and lowest among Indians (86.22%).

Conclusion: CBCT is a valuable tool for detecting the MIC. Careful assessment of the MIC and interforaminal region is recommended to prevent nerve injury and associated complications such as sensory disturbances and hemorrhage.

Keywords: Brazil, China, Imaging, Mandibular nerve, Mental foramen, Malaysia, Spiral Cone-Beam Computed Tomography

Introduction

The Mandibular Incisive Canal (MIC) is the mesial extension of the Inferior Alveolar Canal (IAC), housing the incisive nerve and blood vessels responsible for innervating the lower anterior teeth. The MIC is typically less corticated and smaller in diameter compared to the IAC (1-3). Some researchers propose that the incisive nerve passes through intramedullary spaces rather than a defined bony canal, making it difficult to detect using conventional imaging (4). In certain cases, the MIC is not radiographically evident due to its plexiform nature (5).

Olivier first described the anterior course of the inferior alveolar nerve within the IAC until its division at the mental foramen, where the mental nerve exits and the incisive nerve continues mesially within the MIC, parallel to the anterior teeth roots (6,7).

The interforaminal area is commonly used for procedures like implant placement and bone graft harvesting (8,9). Although traditionally considered a safe zone devoid of major anatomical structures, complications such as hemorrhage and neurosensory disturbances have been reported (10,11). Implant placement in this region may risk MIC perforation, leading to epineurium edema, sensory disturbances, paresthesia, and potential osseointegration failure (12,13).

To avoid such complications, careful evaluation of the MIC and anterior loop is essential (14,15).

The anterior loop describes the mandibular canal's curvature before it opens at the mental foramen (16). While conventional radiography is widely used, its limitations in detecting the MIC and anterior loop are well-known. Cross-sectional imaging, particularly CBCT, allows precise identification of these structures (17,18). Although ultrasound may provide information on the mental foramen, it lacks the diagnostic accuracy of CBCT (19,20).

This review investigates the MIC's anatomical characteristics and its clinical significance, emphasizing variations across populations to aid in preventing surgical complications.

Materials and Methods

An electronic search was conducted in August 2023 using PubMed/MEDLINE and Scopus with MeSH terms and keywords: “mandibular incisive canal,” “incisive canal,” “computed tomography, cone beam,” “CBCT,” “inferior alveolar nerve,” and “mental foramen.” Relevant studies published in English up to August 28, 2023, were included. Manual cross-referencing supplemented the search. After screening 361 studies, 249 were shortlisted based on titles and abstracts. Studies focusing solely on panoramic imaging or cadaveric assessments were excluded (1,7,21). After full-text review, 35 studies met the inclusion criteria (Figure 1).




Figure 1. Flowchart showing the step-by-step identification of the studies via databases.

Data collected included

- Study authors and year
- Sample size
- Patient demographics (sex and age)
- MIC prevalence
- CBCT device, software, and slice thickness used
- Patient ethnicity
- MIC mean length and width
- Distances from MIC to anatomical landmarks

Statistical analysis used means for continuous variables and percentages for categorical data. Quality assessment was independently conducted by two reviewers, with a third resolving disagreements.

Results**Study selection**

Thirty-five studies assessing the MIC *via* CBCT (published between 2009 and 2023) were included. Study characteristics are detailed in tables 1 and 2.

Demographics

MIC was observed in both sexes, with a higher prevalence among females (F:M=1.12:1). The mean age was 45.11 years (range 10–88) and sample sizes ranged from 50 to 1008.

CBCT examination details

Twelve CBCT units (*e.g.*, *i-CAT*, *NewTom*) and 14 software platforms were employed, with slice thicknesses varying from 0.01 mm to 1 mm (mean 0.5 mm). Imaging parameters like *kVp* (60–120) and *mA* (1–36) varied across studies.

Ethnic variations

MIC prevalence differed across ethnicities, ranging from 83.1% (Americans) to 100% (Malays). Chinese subjects had the longest (12.96 mm) and widest (2.5 mm) canals, while Brazilian populations had the smallest measurements.

Table 1. Reported articles of MIC presence and its characteristics in various ethnicities in the literature up until August 2023

Paper	Year	Patients	Age (yrs)	Incisive canal presence in BCT	CBCT and Applied <i>kVp</i>	Software	Slice thickness	Race	CBCT recommendation	Canal location	Mean length of MIC
Makris <i>et al</i> 22	2009	100 M=48 F=52	10-80 Mean =54.9	83.5%	NewTom -	Newtom software	-	-	✓	Buccal	15.13
Pires <i>et al</i> 23	2009	89 F=51 M=38	-	83.1%	i-CAT 120 <i>kVp</i>	-	-	American	-	-	7±3.8
Parnia <i>et al</i> 24	2012	96 M>F	20-77 M=46.60	83%	Planmeca Promax -	Romexis	1 mm	-	-	-	-
Al-Ani <i>et al</i> 25	2012	60	18-80	100%	i-CAT 120 <i>kVp</i>	SimPlant	-	Malay Indian Chinese	✓	Buccal	-
Dalili <i>et al</i> 14	2012	84 M=41 F=43	13-77 Mean =49	92.3% M=F R=L	NewTom	-	1 mm	Iranian	-	Buccal	-
Orhan <i>et al</i> 12	2013	356 F=190 M=166	25-83 Mean =47.6	91%	NewTom 120 <i>kVp</i>	NNT	0.3 mm	Turkish	-	Buccal	12.4 M>F
Yovchev <i>et al</i> 26	2013	140 F=70 M=70	18-79 Mean =46.6	92.9% Unilateral =24.2 Bilateral =68.6	ILUMA	Kodak dental imaging	-	-	-	-	-
Rosa <i>et al</i> 27	2013	352 M=129 F=197	-	97%	i-CAT	i-CAT Vision	0.5 mm	Brazilian	✓	lingual	9.11 ±3.0

Contd. table 1.

Sahman <i>et al</i> 28	2014	243 F=131 M=113	-	94.4%	NewTom.	NNT	-	Turkish	✓	-	-
Raitz <i>et al</i> 29	2014	150 M=75 F=75	-	90% R>L	i-CAT	-	0.25 mm	-	✓	-	-
Pereira- Maciel 30	2015	100 F=63 M=37	20-80 Mean =50.2	100%	i-CAT 120 kVp	iCAT software	-	Brazilian	✓	Buccal	9.8 ±3.8 L=R M=F
Ramesh <i>et al</i> 31	2015	120	30-50	71.66% Unilateral =23.33 Bilateral =48.33	Planmeca 90 kVp	Planmeca romexis	-	Indian	-	-	10.17 ±4.68
Panjnoush <i>et al</i> 10	2015	200 F=110 M=90	48.78± 13.65	97.5%	Planmeca	Planmeca romexis	0.1 mm	Iranian	✓	Buccal	R=10.4 8±4.53 L=10.40 ±4.52 M= 10.44
De brito <i>et al</i> 32	2016	90 M=56, F=34	15-76 Mean =26.3	24.4 % M=F R>L	i-CAT	i-CAT vsion	0.25 mm	-	✓	-	-
Kong <i>et al</i> 33	2016	50 M=25 F=25	18-42 Mean =29.82 ±7.0	100%	Newtom	-	-	Han chinese	-	Buccal	L=17.84 R=17.73 M=17.78
Kabak <i>et al</i> 16	2016	100 M=53 F=47	15-72 Mean= 38.3± 14	92%	Galileos GAX5	-	-	Belarusian	✓	-	-
Sakhdari <i>et al</i> 15	2016	200 M=100 F=100	21-68 Mean =47	87.5%	NewTom	NNT	1 mm	Iranian	✓	-	-
Haghanifar <i>et al</i> 34	2017	207 M=97 F=110	45.7± 13.83		Cranex 3D	On demand	1 mm	Iranian	✓	-	-
Isik <i>et al</i> 17	2017	430 M=185 F=245	15-86 Mean =47.14	89.1%	J.Morita	I-Dixel	-	Turkish	✓	-	-
Sener <i>et al</i> 35	2017	70		74.3%	Kodak 9000 3D	Kodak	-	Turkish	-	-	-
Gomes <i>et al</i> 8	2018	100 M=50 F=50	27-73	78% F>M	i-CAT	Dental slice	0.01 mm	-	✓	Buccal	19.9 ±0.4 M>F
Lim <i>et al</i> 1	2018	100 M=47 F=53	18-80	-	KaVo	eXam vision	-	Chinese malay indian	✓	Buccal at start, lingual at end	11.31 ±2.65 Malays> chinese> indians
Valizadeh <i>et al</i> 19	2018	256 M=123 F=133	>18	100%	NewTom	-	-	-	-	-	7.15
Xie <i>et al</i> 36	2019	1008 F=487 M=521	>18 M=41.1 ±14.81	87% ?	Newtom VG 110 kVp	NNT viewer	-	Chinese	-	-	9.46

Contd. table 1.

Zhang <i>et al</i> 11	2019	535 F=308 M=227	17-88 Mean =36	92.8% M>F	Newtom	NNT	0.5 mm	Chinese	-	-	13.4 ±3.3
Malusare <i>et al</i> 9	2019	150	-	87%	Kodak CS	CS software	-	Indian	✓	-	R=13.4 L=12.4 M=12.9
Gilis <i>et al</i> 37	2019	50	20-83 Mean =48	99%	Planmeca promax	Romexis	0.2 mm	-	✓	-	6.65 ±4.1
Ghonche <i>et al</i> 38	2019	103 M=39 F=64	20-80 Mean =53.18 ±14.4	90% Bilateral =68% Unilateral =22%	Newtom	NNT	-	Iranian	✓	Buccal (at the initiation point)	7.3 mm R>L
Ayesha <i>et al</i> 39	2019	80 M=F	20-60 (18- 80)?	43.89% L>R	Scanora 3D	On Demand3D scanora3D	-	-	-	-	-
Ferreira barbosa <i>et al</i> 40	2020	847 Bilateral =493 Unilateral =153	18-69	76.3% F>M Mainly 4 th to 6 th decades	CS9000 3D gendex CB500 i-CAT next i-CAT classic	Carestream 3D	-	-	✓	-	7.7 mm M>F L=R
Ramaswamy <i>et al</i> 4	2020	100 F=54 M=46	20-70	-	Carestream	-	-	Indian	✓	Buccal	F=13.43 ±3.03 M=13.93 ±4.18 Mean =13.68
Alshamrani <i>et al</i> 41	2021	93	18-50	96.8% L>R	Planmeca promax	Romexis	-	Saudi arabian	✓	-	-
Martins <i>et al</i> 42	2022	50 F=33 M=17	14-74 Mean =38?	100% ?	i-CAT 120 kVp	Blue Sky plan	-	North brazilians		-	-
Nikkerdae <i>et al</i> 3	2022	378 M=185 F=193	14-75 Mean =47.08 ±12.6	97.62% R=L	Newtom	NNT	0.5 mm	Iranian	✓	Buccal	R=12 ±3.29 L=12.21 ±3.38 M=12.1
Nasher <i>et al</i> 5	2023	180 M=74 F=106	16-80	95.6% F>M R=L	Vatech	Ez3D	0.5 mm	Yemeni	✓	Buccal	12.74 ±4.4

Table 1. Reported articles of MIC presence and its characteristics in various ethnicities in the literature up until August 2023

Width of MIC (diameter)	Distance A*	Distance B	Distance C	Distance D	Distance E	Distance F	Distance G	Distance H	Distance I	Distance J	Distance K	Distance L
	M =3.59 ±1.03	M =5.65 ±1.7	M =11.15 ±1.76	-	M=4.8 ±1.7	M=5.6 ±1.6	M=9.4 ±1.7	-	-	-	11.5	-
	2.2 ±1.1		10.4 ±2.3	-	3.3 ±1.7	5.2±2	10.1 ±2.5	-	2.4	-	10.2 ±2.4	-

Contd. table 1.

1.47 ±0.50	-	-	-	-	-	-	-	-	3.48 ±1.17	4.46 ±1.40	8.72 ±1.43	-
-	-	-	-	-	-	-	-	-	3.15 ±1.28	4.78 ±2	9.86 ±2.51	-
-	-	-	-	-	-	-	-	-	-	-	-	-
-	-	-	-	-	-	-	-	-	-	-	10.5	-
-	-	-	-	-	-	-	-	-	-	-	-	-
1.48 ±0.66	-	-	-	-	-	-	-	-	-	-	-	-
R=1.9 ±0.45 L=1.9 ±0.41 M=1.92	-	-	-	-	-	-	-	-	-	-	-	-
-	-	-	-	-	-	-	-	-	-	-	-	-
-	-	-	-	-	-	-	-	-	4.62 ±1.41	6.25 ±2.03	7.06 ±2.95	10.25 ±2.27
2.57 ±0.40	1.48 ±0.22	4.46 ±0.26	9.42 ±0.77	-	4.47 ±0.35	5.56 ±0.69	9.09 ±0.65	-	-	-	-	-
-	-	-	-	-	-	-	-	-	-	-	-	-
-	-	-	-	-	-	-	-	-	-	-	-	-
-	M =2.97 ±0.83	M=5.25 ±1.52	M=9.92 ±1.64	M=17.86 ±2.60	M=4.21 ±1.22	M=5.96 ±1.66	M=11.50 ±3.92	M=17.72 ±4.17	-	-	-	-
-	-	-	-	-	-	-	-	-	-	-	-	-
2.04 ±1.1	-	-	-	-	-	-	-	-	-	-	-	-
R=1.12 ±0.31 L=1.06 ±0.28 M=1.09	-	-	-	-	-	-	-	-	-	-	-	-
-	-	-	-	-	-	-	-	-	-	-	-	-
-	-	-	-	-	-	-	-	-	-	-	-	-
1.4 ±0.1 M=F	-	-	-	-	-	-	-	-	4	6.2	10	-
-	M =3.76 ±1.11	M=6.63 ±1.51	-	-	M=5.89 ±1.70	M=4.72 ±1.61	-	-	5.06	5.47	-	-
-	-	-	-	-	-	-	-	-	-	-	-	-
-	-	-	-	-	-	-	-	-	-	-	-	-
2.5 ±0.5	-	-	-	-	-	-	-	-	3.7 ±0.9	5.1 ±1.6	8.9 ±1.7	19.5 ±3.8
-	-	-	-	-	-	-	-	-	-	-	-	-
-	2.16 ±1.21	5.56 ±1.98	10.33 ±1.98	16.14 ±5.76	-	-	-	-	-	-	-	-
R=1.19 ± 0.48 L=1.29 ±0.93 M=1.24	-	-	-	-	-	-	-	-	-	-	-	-

Contd. table 1.

-	-	-	-	-	-	-	-	-	4.33 ±1.47	4.34 ±1.53	9.41 ±1.83	-
-	M =2.6 ±1.27	M=5.13 ±1.7	M=9.32 ±1.92	M=16.37 ±5.90	M=3.96 ±1.43	M=4.61 ±1.65	M=8.76 ±2.07	M=18.60 ±3.51	-	-	-	16.63 ±7.98
F=1.74 ±0.40									F=3.56 ±1.01	F=4.01 ±1.13	F=6.94 ±1.65	
M=1.94 ±0.58	-	-	-	-	-	-	-	-	M=4.01 ±1.13	M=4.33 ±1.51	M=8.53 ±1.55	-
Mean =1.84									Mean =3.78	Mean =4.17	Mean =7.7	
-	-	-	-	-	-	-	-	-	4.88	5.54	9.94	-
1.29 ?	3.15 ±1.27	5.41 ±1.80	8.95 ±2.41	-	-	-	-	-	-	-	-	-
R=1.89 ±0.46									R=2.08 ±0.94	R=4.18 ±1.52	R= 9.68 ±1.95	
L=1.94 ±0.45	-	-	-	-	-	-	-	-	L=2.15 ±1.02	L=4.14 ±1.54	L= 9.37 ±1.65	-
M=1.91									M=2.11	M=4.16	M=9.52	
1.66 ±0.44	-	-	-	-	-	-	-	-	3.12 ±1.09	4.35 ±1.5	8.61 ±1.9 mm	16.77 ±4.06

*Distance A: Distance of the initial portion of the Mandibular Incisive Canal (MIC) to the buccal table.

Distance B: Distance of the initial portion of the Mandibular Incisive Canal (MIC) to the lingual table.

Distance C: Distance of the initial portion of the Mandibular Incisive Canal (MIC) to the inferior mandibular border.

Distance D: Distance of the initial portion of the Mandibular Incisive Canal (MIC) to the alveolar canal.

Distance E: Distance of the final portion of the Mandibular Incisive Canal (MIC) to the buccal table.

Distance F: Distance of the final portion of the Mandibular Incisive Canal (MIC) to the lingual table.

Distance G: Distance of the final portion of the Mandibular Incisive Canal (MIC) to the inferior mandibular border.

Distance H: Distance of the final portion of the Mandibular Incisive Canal (MIC) to the alveolar canal.

Distance I: Distance of the Mandibular Incisive Canal (MIC) to the buccal table.

Distance J: Distance of the Mandibular Incisive Canal (MIC) to the lingual table.

Distance K: Distance of the Mandibular Incisive Canal (MIC) to the inferior mandibular border.

Distance L: Distance of the Mandibular Incisive Canal (MIC) to the alveolar canal.

Table 2. Demographic characteristics and variables of the articles

Variables	Outcome
Sex	F>M F=53% (3064) M=47% (2692) F/M:1.12
Age	10-88 years Mean:45.11
MIC presence in CBCT	87.11% (24.4-100)
Mean length of MIC (mm)	11.30(6.65-19.9)
Mean width of MIC (mm)	1.72(1.09-2.50)
Distance of the MIC to buccal cortex (mm)	3.71(2.11-5.06)
	Initial portion 2.73 Final portion 4.43
Distance of the MIC to lingual cortex (mm)	4.98(4.16-6.25)
	Initial portion 5.34 Final portion 5.27
Distance of the MIC to inferior mandibular border (mm)	9.37(7.06-11.5)
	Initial portion 9.92 Final portion 9.77
Distance of the MIC to alveolar crest (mm)	15.76(10.2-19.5)
	Initial portion 16.79 Final portion 18.16

Mandibular Incisive Canal (MIC).

Table 3. Ethnic characteristics and variables of the studies

	America	Belarus	Brazil	Chinese	Indian	Iranian	Malaysia	Saudi arabia	Turkish	Yemen
MIC presence (%)	83.1	92	99 (97-100)	93.26 (87-100)	86.22 (71.66-100)	92.98 (87.5-97.62)	100	96.80	87.2 (74.3-94.4)	95.6
MIC length (mm)	7	-	9.45	12.96	12.01	9.94	11.31	-	12.40	12.74
MIC width (mm)	-	-	1.38	2.50	2.20	1.57	-	-	1.92	1.66
MIC to buccal (mm)	2.40	-	4.62	3.97	3.99	2.10	4.10	4.88	-	3.12
MIC to lingual (mm)	-	-	6.25	5.11	4.80	4.16	5.12	5.54	-	4.35
MIC to inferior alveolar (mm)	10.20	-	7.06	9.38	8.78	9.52	9.86	9.94	10.5	8.61
MIC to alveolar (mm)	-	-	10.25	19.5	-	-	-	-	-	16.70

Mandibular Incisive Canal (MIC).

MIC prevalence

The average MIC prevalence was 87.11% (range 24.4-100%). CBCT was consistently recommended for preoperative assessment. Buccal canal positioning was most common, with a single study reporting a lingual position.

Anatomical variations

The average MIC length was 11.30 mm and the diameter 1.72 mm. Mean distances from the canal to anatomical landmarks were:

- Buccal cortex: 3.71 mm
- Lingual cortex: 4.98 mm
- Inferior mandibular border: 9.37 mm
- Alveolar crest: 15.76 mm

Measurements varied between the initial and terminal portions of the MIC.

Discussion

The MIC is a critical anatomical structure extending from the IAC, housing neurovascular elements supplying the anterior mandible. First documented in 1928 (6), its presence was later corroborated by anatomical and radiographic studies (1,43). CBCT

has significantly improved MIC visualization (8). Anatomical assessments on cadavers have detected MIC in up to 90% of the cases (5,7).

Typically, the MIC courses inferiorly and lingually from the mental foramen, become indistinct near the central incisors as it merges into a neurovascular plexus (29,40). This plexus facilitates midline cross-innervation. The incisive nerve is now considered a prominent structure rather than a mere anatomical variation (22).

Detection rates vary due to ethnic differences, imaging methods, and examiner expertise (32). Prevalence ranged from 24.4 to 100% (19), with Malays showing the highest and Indians the lowest detection rates (2,4,9,25,31). Visibility is influenced by factors like canal cortication, age-related bone changes, and skeletal development (40,44).

While some studies reported no sex-based difference in MIC prevalence (14,15,22,26,29,39), others noted a higher prevalence in females or males, particularly among Chinese cohorts (36). Laterality also varied, with some studies showing right-side dominance (29,32), others left-side (39,41), and some reporting bilateral symmetry (3,5,14). Bilateral presentation

was generally more common (26,31,38).

Most studies identified a buccal canal position, although a Brazilian study noted lingual placement (27). Implant placement should account for MIC location to prevent complications (3,33). Ethnic, age, and sex differences underline the need for preoperative CBCT (40).

Accurate measurements of MIC dimensions and positions are vital for safe surgical planning. Direct trauma may cause sensory disturbances; indirect trauma (*e.g.*, hematoma) may exert pressure on the nerve (1). Implants near the MIC can cause neurosensory damage, edema, and osseointegration failure. Arterial injury within the MIC can result in severe bleeding during surgery (12).

A MIC diameter exceeding 2 mm may impair implant stability and increase complication risks (14,23). The average MIC diameter in studies was 1.72 mm, with larger diameters noted in Chinese and Indian populations.

Panoramic radiography underestimates MIC detection due to inherent limitations like distortion (46). CBCT, with superior cross-sectional imaging, provides reliable preoperative assessment (33).

High-resolution MRI has been proposed for detailed analysis but remains impractical due to cost and availability (33). CBCT remains the recommended tool for identifying the MIC.

Conclusion

Given its high prevalence, clinicians should recognize the MIC's anatomical significance to prevent nerve injury, sensory disturbances, and intraoperative bleeding. With an 87.11% detection rate, CBCT is invaluable for mapping the MIC, anterior loop, and mental foramen. The variability among populations underscores the necessity of individualized CBCT assessment before interforaminal surgical interventions. Further research is essential to better understand anatomical variations and optimize patient outcomes.

Funding

This study received no specific financial support.

Conflict of Interest

There was no conflict of interest in this manuscript.

References

1. Mardinger O, Chaushu G, Arensburg B, Taicher S, Kaffe I. Anatomic and radiologic course of the mandibular incisive canal. *Surg Radiol Anat* 2000;22:157–161.
2. Lim JTS, Kang WJ, Bapat RA, Kanneppady SK, Pandurangappa R. Evaluation of mandibular incisive canal using cone beam computed tomography in Malaysians. *J Maxillofac Oral Surg* 2019;18:596–603.
3. Nikkerdar N, Golshah A, Mahmoodivesali R, Falah-Kooshki S. Assessment of implant-related anatomical landmarks in the mandibular interforaminal region in an Iranian population using cone-beam computed tomography. *Contemp Clin Dent* 2022;13:125–34.
4. Ramaswamy P, Kiran C, Raju BM, Swathi M, Sharanya G. Evaluation of mandibular incisive canal and its relationship to adjacent anatomical landmarks using cone-beam computed tomography. *J Oral Maxillofac Radiol* 2020;8:23–9.
5. Nasher A, Juain A, Da'er S, Farhan A, Alareqi N. The Prevalence of mandibular incisive canal among a sample of Yemeni adults obtained from cone-beam computed tomography. *Saudi J Oral Dent Res* 2023;8:50–8.
6. Olivier E. The inferior dental canal and its nerve in the adult. *Br Dent J* 1928;49:356–8.
7. Mraiwa N, Jacobs R, Moerman P, Lambrichts I, van Steenberghe D, Quirynen M. Presence and course of the incisive canal in the human mandibular interforaminal region: two-dimensional imaging versus anatomical observations. *Surg Radiol Anat* 2003;25:416–23.
8. Gomes LT, de Almeida Barros Mourão CF, Braga CL, de Almeida LFD, de Mello-Machado RC, Calasans-Maia

- MD. Anatomic evaluation of the incisive canal with cone beam computed tomography and its relevance to surgical procedures in the mental region: a retrospective study in a Brazilian population. *Oral Maxillofac Surg* 2018;22:379-84.
9. Malusare P, Navalkar A, Das D, Patil B. Assessing the dimensions of mandibular incisive canal and its relationship to adjacent anatomical landmarks using cone beam computed tomography in Indian sub-population – a retrospective study. *J Indian Acad Oral Med Radiol* 2019;31:100.
 10. Panjnoush M, Rabiee ZS, Kheirandish Y. Assessment of location and anatomical characteristics of mental foramen, anterior loop and mandibular incisive canal using cone beam computed tomography. *J Dent* 2016;13:126–32.
 11. Zhang YQ, Yan XB, Zhang LQ, Xie XY, Liu DG, Zhang ZY. Prevalence and morphology of mandibular incisive canal: comparison among healthy, periodontitis and edentulous mandibles in a population of the Beijing area using cone-beam computed tomography. *Chin J Dent Res* 2019;22:241–9.
 12. Orhan K, Icen M, Aksoy S, Ozan O, Berberoglu A. Cone-beam CT evaluation of morphology, location, and course of mandibular incisive canal: considerations for implant treatment. *Oral Radiol* 30:64–75.
 13. Fuentes R, Arias A, Bucchi C, Saravia D, Dias F. Prevalence and morphometric characteristics of the mandibular incisive canal through panoramic radiographs in a Chilean population. *Int J Morphol* 2017;35:931–7.
 14. Kajan ZD, Salari A. Presence and course of the mandibular incisive canal and presence of the anterior loop in cone beam computed tomography images of an Iranian population. *Oral Radiol* 2012;28:55–61.
 15. Sakhdari S, Hafezi L, Esmaili M. Prevalence of the inferior alveolar nerve's anterior loop and mandibular incisive canal by use of Cone Beam Computed Tomography (CBCT) in an Iranian population. *J Res Dent Maxillofac Sci* 2016;1:14–21.
 16. Kabak SL, Zhuravleva NV, Melnichenko YM, Savrasova NA. Study of the mandibular incisive canal anatomy using cone beam computed tomography. *Surg Radiol Anat* 2017;39:647–55.
 17. Işık BK, Taşşöker M, Menziletoğlu D, Şener S, Esen A. Assessment of the mandibular incisive canal by panoramic radiograph and cone-beam computed tomography. *Selcuk Dent J* 2017;4:17-22.
 18. de Oliveira-Santos C, Souza PHC, de Azambuja Berti-Couto S, Stinkens L, Moyaert K, Rubira-Bullen IRF, et al. Assessment of variations of the mandibular canal through cone beam computed tomography. *Clin. Oral Investig* 2012;16:387–93.
 19. Valizadeh S, Alesheikh H, Iranparvar Alamdari M. Cone-beam computed tomographic assessment of anatomical variations of the interforaminal region in a selected Iranian population. *J Dent Sch* 2021;38:153–57.
 20. Çağlayan F, Sümbüllü MA, Akgül HM. Is ultrasonography sufficient for evaluation of mental foramen? *Dentomaxillofac Radiol* 2019;48:20180252.
 21. Uchida Y, Noguchi N, Goto M, Yamashita Y, Hanihara T, Takamori H, et al. Measurement of anterior loop length for the mandibular canal and diameter of the mandibular incisive canal to avoid nerve damage when installing endosseous implants in the interforaminal region: a second attempt introducing cone beam computed tomography. *J Oral Maxillofac Surg* 2009;67:744–50.
 22. Makris N, Stamatakis H, Syriopoulos K, Tsiklakis K, van der Stelt PF. Evaluation of the visibility and the course of the mandibular incisive canal and the lingual foramen using cone-beam computed tomography. *Clin Oral Implants Res* 2010;21:766-71.
 23. Pires CA, Bissada NF, Becker JJ, Kanawati A, Landers MA. Mandibular incisive canal: cone beam computed tomography. *Clin Implant Dent Relat Res* 2012;14:67–73.
 24. Parnia F, Moslehifard E, Hafezeqoran A, Mahboub F, Mojaver-Kahnamoui H. Characteristics of anatomical landmarks in the mandibular interforaminal region: a cone-beam computed tomography study. *Med Oral Patol Oral Cir Bucal* 2012;17:e420-5.
 25. Al-Ani O, Nambiar P, Ha KO, Ngeow WC. Safe zone for bone harvesting from the interforaminal region of the

mandible. *Clin Oral Implants Res* 2013;24 Suppl A100:115-21.

26. Yovchev D, Deliverska E, Indjova J, Zhelyazkova M. Mandibular incisive canal: a cone beam computed tomography study. *Biotechnol Biotechnol Equip* 2013;27:3848–51.

27. Rosa MB, Sotto-Maior BS, Machado Vde C, Francischone CE. Retrospective study of the anterior loop of the inferior alveolar nerve and the incisive canal using cone beam computed tomography. *Int J Oral Maxillofac Implants* 2013;28:388-92.

28. Sahman H, Sekerci AE, Sisman Y, Payveren M. Assessment of the visibility and characteristics of the mandibular incisive canal: cone beam computed tomography versus panoramic radiography. *Int J Oral Maxillofac Implants* 2014;29:71-8.

29. Raitz R, Shimura E, Chilvarquer I, Fenyo-Pereira M. Assessment of the mandibular incisive canal by panoramic radiograph and cone-beam computed tomography. *Int J Dent* 2014;2014:187085.

30. Pereira-Maciel P, Tavares-de-Sousa E, Oliveira-Sales MA. The mandibular incisive canal and its anatomical relationships: a cone beam computed tomography study. *Med Oral Patol Oral Cir Bucal* 2015 Nov 1;20:e723-8.

31. Ramesh AS, Rijesh K, Sharma A, Prakash R, Kumar A, Karthik. The prevalence of mandibular incisive nerve canal and to evaluate its average location and dimension in Indian population. *J Pharm Bioallied Sci* 2015;7(Suppl 2): S594-6.

32. de Brito ACR, Nejaim Y, de Freitas DQ, de Oliveira Santos C. Panoramic radiographs underestimate extensions of the anterior loop and mandibular incisive canal. *Imaging Sci Dent* 2016;46:159-65.

33. Kong N, Hui M, Miao F, Yuan H, Du Y, Chen N. Mandibular incisive canal in Han Chinese using cone beam computed tomography. *Int J Oral Maxillofac Surg* 2016;45:1142-6.

34. Haghanifar S, Moudi E, Bijani A. Mandibular canal and its incisive branch: a CBCT study. *World Fam Med J* 2017;15:133–40.

35. Sener E, Onem E, Akar GC, Govsa F, Ozer MA, Pinar Y, et al. Anatomical landmarks of mandibular interforaminal region related to dental implant placement with 3D CBCT: comparison between edentulous and dental mandibles. *Surg Radiol Anat* 2018;40:615-23.

36. Xie L, Li T, Chen J, Yin D, Wang W, Xie Z. Cone-beam CT assessment of implant-related anatomy landmarks of the anterior mandible in a Chinese population. *Surg Radiol Anat* 2019;41:927-34.

37. Gilis S, Dhaene B, Dequanter D, Loeb I. Mandibular incisive canal and lingual foramina characterization by cone-beam computed tomography. *Morphologie* 2019;103:48-53.

38. Ghoncheh Z, Moghaddam Zadeh B, Shaeri S. Prevalence and position of mandibular incisive canal, anterior loop of the mandibular canal and lingual foramen using cone beam computed tomography. *J Craniomaxillofac Res* 2020;6:143-50.

39. Ayesha RT, Pachipulusu B, Govindaraju P. Assessment of prevalence and position of mandibular incisive canal: a cone beam computed tomography study. *Tzu Chi Med J* 2019;32:205-10.

40. Barbosa DA, Kurita LM, Pimenta AV, Teixeira RC, Silva PG, Ribeiro TR, et al. Mandibular incisive canal-related prevalence, morphometric parameters, and implant placement implications: a multicenter study of 847 CBCT scans. *Med Oral Patol Oral Cir Bucal* 2020;25:e337-e345.

41. Alshamrani AS, Tokhtah RA, Al-Omar A. Cone-beam computed tomography evaluation of prevalence and location of mandibular incisive canal in patients attending King Saud University dental hospital. *J Orthod Sci* 2021;10:21.

42. Martins VB, Oenning AC, Guimarães LC, Junqueira JL, Franco A. CBCT study on the prevalence, morphology and position of the mandibular incisive canal in a North-Brazilian population. *J Clin Exp Dent* 2022;14:e534-e540.

43. Bavitz JB, Harn SD, Hansen CA, Lang M. An anatomical study of mental neurovascular bundle-implant relationships. *Int J Oral Maxillofac Implants* 1993;8:563-7.

44. Jalili MR, Esmaeelinejad M, Bayat M, Aghdasi MM. Appearance of anatomical structures of mandible on panoramic radiographs in Iranian population. *Acta Odontol Scand* 2012;70:384-9.
45. Gala V, Tirpude V, Shah D, Doshi A, Fernandes G. A retrospective cone beam computed tomography (CBCT) study of the assessment of the length of the anterior loop of the inferior alveolar nerve. *Int J Sci Res Sci Technol* 2018;4:828-32.
46. Jacobs R, Mraiwa N, Van Steenberghe D, Sanderink G, Quirynen M. Appearance of the mandibular incisive canal on panoramic radiographs. *Surg Radiol Anat* 2004 Aug;26:329-33.