

Research Article



The Effect of Sensorimotor Synchronization on Gait Spatiotemporal Parameters in Women with Multiple Sclerosis

Mozhgan Memarmoghadam^{1*}, Mansoureh Shahraki², Behzad Mohammadi Orangi³, Luca Correale⁴, Leonardo A. Peyré-Tartaruga⁵

1. Department of Motor Behavior and Sport Biomechanics, Faculty of Sport Sciences, University of Mazandaran, Mazandaran, Iran.
2. Department of Sport Sciences, Faculty of Literature and Humanities, University of Zabol, Zabol, Iran.
3. Department of Sport Science, Faculty of Humanities, Damghan University, Damghan, Iran.
4. Department of Public Health, School of Experimental Medicine and Forensic Sciences, University of Pavia, Pavia, Italy.
5. Exercise Research Laboratory, Universidade Federal do Rio Grande do Sul, Porto Alegre, Brazil.



Citation Memarmoghadam M, Shahraki M, Mohammadi Orangi B, Correale L, Peyré-Tartaruga LA. The Effect of Sensorimotor Synchronization on Gait Spatiotemporal Parameters in Women with Multiple Sclerosis. *Journal of Modern Rehabilitation*. 2025; 19(1):71-79. <http://dx.doi.org/10.18502/jmr.v19i1.17511>

doi <http://dx.doi.org/10.18502/jmr.v19i1.17511>

Article info:

Received: 08 Jun 2024

Accepted: 23 Sep 2024

Available Online: 01 Jan 2025

ABSTRACT

Introduction: Although gait rehabilitation based on sensorimotor synchronization (auditory and visual) is stimulating for patients with neurological disorders and older people, there is little evidence in patients with multiple sclerosis (MS). Therefore, this study investigates the effect of synchronization with rhythmic visual stimulus on gait spatiotemporal parameters, bilateral symmetry, and locomotor rehabilitation index in women with MS.

Materials and Methods: Spatiotemporal parameters, bilateral symmetry and the locomotor rehabilitation index were obtained before and after 6 weeks (three times per week, 30 min each session) of locomotor training. We compared these findings between two groups of 10 patients, each with an expanded disability status scale (EDSS) of 3-6, who performed the gait training with or without rhythmic visual stimulus.

Results: Time×group interaction effect indicated greater significant improvements in the group with rhythmic visual stimulus in self-selected walking speed (SSWS) (P=0.041), stride frequency (P=0.009), stance time (P=0.021), and locomotor rehabilitation index (P=0.036). Stride length, double stance, and swing time were improved in the group with rhythmic visual stimulus, but this change was not significant. Also, bilateral symmetry did not change significantly in the experimental group. Therefore, rhythmic visual stimulation can help improve functional mobility and locomotor rehabilitation index in patients with MS, especially due to the improvements in the temporal parameters of gait.

Conclusion: Therefore, synchronizing gait with a rhythmic visual stimulus can be an effective therapeutic strategy to improve gait and main temporal parameters in patients with MS.

Keywords:

Multiple sclerosis;
Sensorimotor; Synchronization;
Gait

* Corresponding Author:

Mozhgan Memarmoghadam, PhD.

Address: Department of Motor Behavior and Sport Biomechanics, Faculty of Sport Sciences, University of Mazandaran, Mazandaran, Iran.

Tel: +98 (913) 3272189

E-mail: mmemarmoghaddam@yahoo.com



Copyright © 2025 Tehran University of Medical Sciences. Published by Tehran University of Medical Sciences
This work is licensed under a Creative Commons Attribution-NonCommercial 4.0 International license (<https://creativecommons.org/licenses/by-nc/4.0/>).
Noncommercial uses of the work are permitted, provided the original work is properly cited.

Introduction

Multiple sclerosis (MS) is a demyelinating disease of the central nervous system, leading to a wide range of disabilities and disorders in the transmission of neural messages throughout the brain and spinal cord [1, 2]. Gait disorder is one of the most common problems reported in these patients [3]. Liparoti et al. reported that patients with MS show longer stance time, double stance and swing time than healthy people [4]. Bilateral coordination also worsens, and reduced gait speed is a decisive factor in this impairment [5]. This condition indicates impaired gait mechanics in these patients [5], reflected in changes in gait spatiotemporal parameters [6]. Therefore, it is crucial to develop treatment strategies to improve gait biomechanics parameters in these patients.

Sensory disorders play a major role in hampering motor control and coordination in patients with MS [7]. Therefore, treatment strategies based on providing additional sensory cueing to support movement performance, such as sensorimotor synchronization exercises, can be a suitable method to overcome motor control and coordination disorders in patients with MS [8]. Sensorimotor synchronization consists of generating a sequence of motor responses at a frequency similar to a sensory stimulus, ie, producing a motor response that coincides with a sensory stimulus [9]. Motor and cognitive rehabilitation (such as sensorimotor synchronization exercises) can enhance the functional and structural plasticity of the brain in patients with MS [10]. In patients with MS, the evidence shows that plasticity limits the clinical effects of injury by creating patterns of brain activity different from healthy volunteers and improvement in motor function following exercises is accompanied by reorganizing those altered patterns [11]. Also, exercising a practical movement under time constraints can be effective for practicing timing and treating patients' coordination problems [12]. So, it seems that exercises based on sensorimotor synchronization can help improve these patients' gait mechanics.

A limited number of studies have investigated the effect of synchronization with different sensory stimuli in patients with MS [13-15]. In most studies, the effect of synchronization with rhythmic auditory stimulus on the gait performance of these patients has been investigated, which indicates the positive effects of this type of stimulus on improving the gait mechanics of patients with MS [13, 14]. However, as far as we know, only one study investigated the effect of synchronization with rhythmic

visual stimulus on gait performance in patients with MS [15]. The study findings showed that practicing gait with a rhythmic visual stimulus improved stride length, stride time, double support time, gait cadence and speed. However, this difference was not statistically significant between the experimental and the control groups [15]. Further studies have also reported a positive effect of this type of intervention for people suffering from other neurological diseases, such as Parkinson disease [16]. Research design problems in these studies include short intervention time and no control group. Despite these limitations, the improvement of gait spatiotemporal parameters of these patients shows that gait training with rhythmic sensory stimuli can be a promising rehabilitation strategy in patients with MS. Researchers state that more investigation, stronger evidence, and higher quality are required to ensure the extent of the positive effects of this intervention on the gait rehabilitation of patients with MS [13-15].

The present study is the first study that, while controlling the limitations of previous studies, investigates the effect of gait synchronization with the rhythmic visual stimulus on gait spatiotemporal parameters, motor rehabilitation index, and bilateral symmetry in patients with MS.

We hypothesized that gait training synchronized with rhythmic visual stimuli would improve gait spatiotemporal parameters, including self-selected walking speed (SSWS), stride frequency, stride length, stance time and swing, double stance, and locomotor rehabilitation index, to a greater extent compared to gait training alone. Also, we hypothesized that bilateral symmetry would improve significantly by adding the synchronization method to gait training due to the inherent stimulus of controlling the stepping and additional gains in the gait speed already observed in the previous study.

Materials and Methods

Twenty women with MS (relapsing-remitting MS) members of the Zahedan MS Association, Zahedan City, Iran, were selected using available and targeted sampling methods [17]. The criteria for participating in the present research included 1) Being 18 years old or more, 2) Presenting an extended disability status scale of 3 to 6, 3) Walking ability of at least 100 feet without physical assistance, 4) Lack of treatment and recurrence and exacerbation of the disease in the past 30 days, 5) Lack of cardiovascular and rheumatic diseases, 6) lack of severe pain in the lower joints, 7) Lack of participation in any regular exercise for the past three months, 8) Normal or

corrected vision with glasses [15]. The neurologist determined the degree of disability of the patients with an expanded disability status scale (EDSS) questionnaire for MS patients. All participants completed a written informed consent form to participate in the study. They were divided by the simple random method into an experimental group with 10 participants (gait training with rhythmic visual stimulus) and a control group with 10 participants (simple gait). Eight qualisys motion analysis cameras and the Qualisys Track Manager software, version 2014, were used to measure gait biomechanics. To measure the parameters of gait stability, right and left leg stance time and swing time, after extracting the gait cycle and toe-off event, the time interval between heel contact (HC) to toe-off has been considered stance time and the time interval between toe off to HC was considered swing time [18]. After measuring the temporal-spatial parameters of gait, the symmetry index was used to evaluate the bilateral symmetry [5]. Also, the locomotor rehabilitation index was determined through the ratio of SSWS to optimal walking speed (OWS) [19].

Exercises were performed for 6 weeks, three sessions per week, and 30 minutes per session. The exercises consisted of gait synchronization with a rhythmic visual stimulus. The stimulus was created with the design of a device including a Musedo (MT-100) metronome, an optical LED connected to the metronome and a piece of wire 10 cm in front of the right inner edge of a hat. At the beginning of the first session of the experimental group exercise program, the visual stimulus frequency was set 10% higher than each participant's preferred walking cadence [15], determined during a 10 m walking test. In this test, a 10 m route was covered at a preferred speed, the number of steps and the time of covering the mentioned distance were recorded and the number of steps in one minute was calculated as the preferred walking cadence of each participant [20]. After setting up the device for each participant, they were asked to cover a distance of 6 m after adapting their steps to the desired stimulus frequency, rotating 180 degrees and returning to the route's beginning [16]. In each exercise session, the participant could split the 30 minute time into smaller intervals if walking for 30 minutes consecutively was difficult. At the end of each week, each participant's preferred walking cadence was measured using a 10 m walking test at the preferred speed [20]. Then, based on the preferred walking cadence plus 10%, the visual stimulus frequency was set for the following week's exercises. The exercises were performed individually, and an instructor walked two steps behind the examinees to support and prevent the participants from falling. The participants in the control group performed only simple gait (Figure 1

shows the consort chart of the study). Three participants, including two in the experimental group and one in the control group, were excluded from the initial statistical sample due to receiving corticosteroids and not attending regularly in the exercise sessions. It should be noted that the pre-test and post-test were conducted by people blind to the grouping. (Table 1 shows the general characteristics of the group participants)

Statistical analysis

Firstly, we analyzed the data qualitatively for the normal distribution using the Shapiro-Wilk test and the homogeneity of the variances, which was confirmed by Levene's test. To evaluate the effect of different interventions, a mixed model ANOVA was used with interventions (gait with and without rhythmic visual stimulation) and time (pre- and post-training) as fixed factors and subjects as random factors (Jasp). Data were reported as mean and standard deviation, with an alpha of $P=0.05$. Finally, Hedges's g and 95% confidence interval (CI) were used to determine the magnitude of the changes, bringing a more practical approach to the obtained results. The effect size (ES) was classified as trivial (<0.19), small (0.20–0.49), medium (0.50–0.79) and large (0.80–1.29) [21].

Results

Time \times group interaction effects indicated greater significant improvements in the group with rhythmic visual stimulus in SSWS ($P=0.041$), stride frequency ($P=0.009$), stance time ($P=0.021$) and locomotor rehabilitation index ($P=0.036$). Stride length, double stance, and swing time improved in the group with rhythmic visual stimulus, but this change was not significant. Also, bilateral symmetry did not change significantly in the experimental group (Table 2).

Violin plots showing the distribution of the gait spatio-temporal parameters and locomotor rehabilitation index according to the time conditions (pre-test and post-test) for the control group (just gait, pink) and the synchronization group (gait synchronization with rhythmic visual stimulus, green). Moreover, the two-sided violin plots indicate the data distribution for each variable (Figure 2).

The ES was 0.15 with a [95% CI, -0.2%, 0.57%] between pre-test and post-test conditions. On the other hand, in the rhythmic training with unequalized total training volume (RTUV) condition, the ES was 0.51 (i.e. medium) with [95% CI, 0.09%, 0.97%]. For cross-sectional area (CSA) values, the ES between the differ-




Figure 1. The consort chart of the study

JMR

ent frequencies of the rhythmic training with equalized total training volume (RTEV) condition was -0.02 (i.e. insignificant) with a [95% CI, -0.43% , 0.40%], while in the RTUV condition, the ES was 0.63 (i.e. medium) with [95% CI, 0.2% , 1.10%] (Figure 2).

Discussion

The present study compared the changes in gait spatiotemporal parameters, locomotor rehabilitation index, and bilateral symmetry after 6 weeks between gait training executed with and without synchronization with rhythmic visual stimuli in a group of patients with MS.

Our main findings showed that the self-selected walk speed, stride frequency, stance time, and locomotor rehabilitation index significantly improved after synchronizing gait with the rhythmic visual stimulus. However, no

significant change was observed despite improved stride length, double stance, and swing time. These findings confirm our first hypothesis. Also, bilateral symmetry did not significantly improve, which rejected our second hypothesis.

Our findings confirm the previous results from studies on spatiotemporal parameters of gait with shorter training periods [15]. Shahraki et al. (2019) showed that gait training with the rhythmic visual stimulus improves stride length, stride time, double support time, cadence, and speed. However, this difference between the experimental and control groups was not statistically significant [15]. The results of the present study, which had a longer training period than the previous study [15], suggest that the specific improvements due to the inclusion of rhythmic visual stimulus as a mode to synchronize the stepping gait were related to temporal parameters more

Table 1. General characteristics of the participants in the groups (n=20)

Variables	Mean±SD		P
	Control Group	Experimental Group	
Participants (male/female)	10(0/10)	10(0/10)	-
Age (y)	40.9±9.2	38.1±12.1	0.149
Height (cm)	1.63±0.06	1.60±9.70	0.195
Body mass (kg)	60.9±11.5	66.1±7.2	0.753
BMI (kg/m ²)	25.6±3.4	26.5±3.5	0.585
EDSS	4±0.8	4.2±0.9	0.684
Lower Limb Length (m)	0.88±0.03	0.87±0.05	0.368

BMI: Body mass index; EDSS: Expanded disability status scale.

JMR

The results for the independent t-test comparing control and experimental groups.

Table 2. Mixed model ANOVA test results for gait spatiotemporal parameters, bilateral symmetry and locomotor rehabilitation index in groups

Variables	Mean±SD				P		
	Control Group		Experimental Group		Group	Time	Group×Time
	Pre-test	Post-test	Pre-test	Post-test			
Self-selected walk speed (m/s)	0.55±0.23	0.6±0.29	0.43±0.19	0.75±0.32	0.885	0.006	0.041*
Stride length (m)	0.75±0.26	0.80±0.29	0.71±0.26	0.95±0.30	0.598	0.009	0.074
Stride frequency (Hz)	0.77±0.06	0.75±0.13	0.60±0.12	0.77±0.14	0.086	0.037	0.009*
Stance time (s)	0.99±0.22	0.95±0.20	1.19±0.23	0.86±0.21	0.485	0.004	0.021*
Swing time (s)	0.40±0.05	0.41±0.06 [‡]	0.49±0.10	0.47±0.23 [§]	0.137	0.945	0.604
Locomotor rehabilitation index (%)	51.6±17.4	55.1±20.8	28.3±12.8	50.4±21.9	0.068	0.006	0.036*
Relative stance time (%)	70.3±3.6	68.8±3.5	69.3±4.1	64.5±7.0	0.092	0.045	0.292
Double stance (s)	0.30±0.1	0.29±0.13	0.35±0.1	0.23±0.1	0.960	0.028	0.060
SL symmetry (%)	32.3±17.7	29.8±17.3	4.88±4.27	3.80±2.71	<0.001	0.495	0.780
SF symmetry	8.75±11.1	8.78±8.02	5.28±6.73	6.45±6.35	0.371	0.758	0.768
ST symmetry	13.5±12.2	9.6±10.4	7.1±4.13	10.4±11.9	0.490	0.889	0.131
SW symmetry	14.1±11.4	32.5±46.3	16.1±9.15	12.5±7.35	0.295	0.314	0.138
DC symmetry	27.1±17.9	29.7±26.9	21.3±16.7	25.7±17.2	0.518	0.491	0.854

Abbreviations: SL: Stride length; SF: Stride frequency; ST: Stance; SW: Swing.

JMR

Superscript symbols indicate statistically significant differences (P<0.01 and P<0.05).




Figure 2. Violin plots

JMR

Note: The distribution of the gait spatiotemporal parameters and locomotor rehabilitation index according to the time conditions (pre-test and post-test). The control group was pink and the experimental group was green.

than spatial parameters. These specific adaptations positively impacted the functional mobility observed here, such as gait speed. In addition to the observed biomechanical improvements, the bilateral coordination was probably enhanced due to higher SSWS done by the synchronized group. Previous evidence has demonstrated that the bilateral accuracy (temporal symmetry between right-left stepping) and bilateral consistency (variability of the temporal symmetry) increase at a faster speed than self-selected in patients with MS [22]. Furthermore, the improvements found in the temporal variables of gait are particularly interesting as they impact cardinal points of movement disorders in MS [23]. Indeed, stride frequency and stance time impairments are highly related to gait speed and disease staging (EDSS). Although reductions in temporal gait parameters due to MS could be part of adopting a more cautious gait strategy, these impairments result in a higher risk of falls [24].

Temporal-specific adaptations in response to an intervention based on temporal gait control raise the question of whether training with visual-spatial stimuli (e.g. visual cues) could bring superior adaptations in spatial variables such as stride length. Future studies could test these possible specific adaptations on mobility gains. This study provides pertinent information for the organization and application of interventions for patients with MS. Individuals with high levels of bilateral asymmetry and low bilateral coordination may likely have important benefits from the rhythmic visual synchronization used in the present study.

Abnormalities in postural control during gait in patients with MS are related to slow movement toward stability limits and delayed responses to perturbations and displacements during gait [25]. The improved responses in the present study may be related to the mitigation of this slowness of movement during gait.

After the intervention, the improved locomotor rehabilitation index indicates that subjects perform their routine gait activities with greater economy and improved pendulum mechanics [19, 26]. The locomotor rehabilitation index, which expresses the relationship between the conceptually, SSWS, is defined as the speed at which people choose to walk in their daily lives because it is the most comfortable speed and OWS is defined as the speed at which the contribution of mechanical parameters is optimized and the metabolic energy expenditure is minimized, is a complementary analysis of spatiotemporal parameters in gait assessment. According to this index, walking is functional and healthy when the SSWS and OWS coincide. Therefore, the locomotor rehabilitation index can be considered a vital performance marker with clinical relevance [19]. This improved metabolic economy may increase physical activity and the quality of movements other than walking (e.g. load lifting, instrument handling) due to the lower degree of fatigue [27]. Since gait is the most commonly reported exercise among adults, walking faster than usual paces may signal additional gains in primary health-related outcomes in the follow-up [27].

Patients with MS adopt a conservative strategy to maintain stability by increasing the duration of the walking cycle [4]. Besides, gait disorder in patients with MS indicates motor control disorder in these patients, which is inversely related to gait speed [4]. Hence, as shown in this study, it seems that improvements in spatiotemporal parameters, especially temporal parameters of gait, following sensory-motor synchronization training with a rhythmic visual stimulus may indicate increased motor control and improve the stability of gait in patients with MS. As previously mentioned, treatment strategies based on providing additional sensory cueing to support movement performance can be a good way to overcome motor control and coordination disorders in patients with MS [8]. Sensorimotor synchronization activates several motor areas of the brain, including the prefrontal cortex, supplementary motor area, premotor cortex, right lower parietal lobe, posterior superior temporal gyrus, anterior insula, cerebellum, and putamen [28]. Activation of brain motor area improves muscle activation and leads to better motor control [29].

The EDSS was used to homogenize the study groups, allowing them to assess gait disorders in these patients. On the other hand, the walking training route in this study was 6 m; that is, the participants in the study returned to the beginning of the route after a distance of 6 m and walked along this route during the entire training period [16]. However, in this protocol, the participants seemed to restart the synchronization process in each turn instead of walking slowly and evenly in a straight line for a long time. Similarly, in daily life, people rarely walk in a straight line and often stop, change direction, and start again, and these protocols are more seen in people's normal lives.

No studies have been performed to evaluate the effect of synchronizing gait with a rhythmic visual stimulus on brain plasticity in patients with MS [30]. The effect of synchronization with the rhythmic visual stimulus on improving spatiotemporal parameters of gait in these patients may result from this rehabilitation intervention on reorganizing or restoring the altered brain patterns. Also, these changes may be the consequence of activity-dependent myelination. Future studies are suggested for clinicians to investigate this issue. Also, clinicians can recommend the above protocol as an effective, cheap, and usable rehabilitation method as a complementary treatment along with drug treatment for patients and medical and rehabilitation centers.

Here, we investigated spatiotemporal parameters, bilateral symmetry, and locomotor rehabilitation index recov-

ery after 6 weeks of locomotor training, comparing these findings between two groups of 10 patients, each performing the gait training with or without rhythmic visual stimulus. We demonstrated that rhythmic visual stimulus improved functional mobility and locomotor rehabilitation index in patients with MS. These improvements were possible because of the rhythmic visual stimulus employed in the present study, constituting an effective method to qualify gait training methods. In addition, we showed that the ability to walk was additionally improved, precisely due to temporal adjustments of gait provided by the rhythmic visual stimulation. Here, we showed that patients with MS sparing improved about $\approx 74\%$ of their initial functional mobility, adding rhythmic visual stimulation—a proportion much higher than just gait.

Conclusion

Rhythmic visual Stimulus can help improve functional mobility and locomotor rehabilitation index in patients with MS, especially due to the improvements in the temporal parameters of gait.

Study limitations

This study also had some limitations. There was no control group with no exercise. It was not possible to control the eating patterns, daily activity patterns, and rest of the participants. Also, it was impossible to match the participants' social, economic, cultural, and health status, mental health, and motivation levels.

Ethical Considerations

Compliance with ethical guidelines

This study was approved by the Ethics Committee of the [University of Mazandaran](#) (Code: IR.UMZ.REC.1399.012). All procedures performed in this study were in accordance with the ethical standards of the institutional and or national research committee and with the 1964 Helsinki Declaration and its later amendments or comparable ethical standards.

Funding

This research did not receive any grant from funding agencies in the public, commercial, or non-profit sectors.

Authors' contributions

Data collection and writing the original draft: Mozghan Memarmoghdam and Mansoureh Shahraki; Review, editing and supervision: All authors.

Conflict of interest

The authors declared no conflict of interest.

Acknowledgments

The authors hereby thank and appreciate all the researchers worldwide who have done research in the field of rehabilitation to improve the physical activity of patients with MS. In addition, the authors also thank all those who participated in this study.

References

- [1] Compston A, Coles A. Multiple sclerosis. *Lancet*. 2008; 372(9648):1502-17. [DOI:10.1016/S0140-6736(08)61620-7] [PMID]
- [2] Baigi SFM, Sarbaz M, Sobhani-Rad D, Mousavi AS, Dahmardeh F, Kimiafar K. Investigating the effects of telerehabilitation on improving the physical activity of individuals with multiple sclerosis: A systematic review of randomized controlled trial. *Journal of Modern Rehabilitation*. 2024; 18(1):1-14. [DOI:10.18502/jmr.v18i1.14724]
- [3] Minden SL, Frankel D, Hadden L, Perloff J, Srinath KP, Hoaglin DC. The Sonya Slifka longitudinal multiple sclerosis study: Methods and sample characteristics. *Multiple Sclerosis Journal*. 2006; 12(1):24-38. [DOI:10.1191/135248506ms12620a] [PMID]
- [4] Liparoti M, Della Corte M, Rucco R, Sorrentino P, Sparaco M, Capuano R, et al. Gait abnormalities in minimally disabled people with Multiple Sclerosis: A 3D-motion analysis study. *Multiple Sclerosis and Related Disorders*. 2019; 29:100-7. [DOI:10.1016/j.msard.2019.01.028] [PMID]
- [5] Pau M, Leban B, Deidda M, Putzolu F, Porta M, Coghe G, et al. Kinematic analysis of lower limb joint asymmetry during gait in people with multiple sclerosis. *Symmetry*. 2021; 13(4):598. [DOI:10.3390/sym13040598]
- [6] Rahimibarghani S, Emami-Razavi SZ, Moghadasi AN, Azadvari M, Fard MS, Rahimi-Dehgolan S. Quantitative changes in gait parameters after cycling among multiple sclerosis patients with ataxia: A pilot study. *Journal of Modern Rehabilitation*. 2022; 16(4). [DOI:10.18502/jmr.v16i4.10763]
- [7] Rolak LA. Multiple sclerosis: It's not the disease you thought it was. *Clinical Medicine & Research*. 2003; 1(1):57-60. [DOI:10.3121/cmr.1.1.57] [PMID]
- [8] Ghai S, Ghai I. Effects of rhythmic auditory cueing in gait rehabilitation for multiple sclerosis: A mini systematic review and meta-analysis. *Frontiers in Neurology*. 2018; 9:386. [DOI:10.3389/fneur.2018.00386] [PMID]
- [9] Molinari M, Leggio MG, Thaut MH. The cerebellum and neural networks for rhythmic sensorimotor synchronization in the human brain. *The Cerebellum*. 2007; 6(1):18-23. [DOI:10.1080/14734220601142886] [PMID]
- [10] Prosperini L, Piattella MC, Gianni C, Pantano P. Functional and structural brain plasticity enhanced by motor and cognitive rehabilitation in multiple sclerosis. *Neural Plasticity*. 2015; 2015:481574. [DOI:10.1155/2015/481574] [PMID]
- [11] Tomassini V, Matthews PM, Thompson AJ, Fuglò D, Geurts JJ, Johansen-Berg H, et al. Neuroplasticity and functional recovery in multiple sclerosis. *Nature Reviews Neurology*. 2012; 8(11):635-46 [DOI:10.1038/nrneuro.2012.179] [PMID]
- [12] Shumway-Cook A, Woollacott M. *Motor control: Translating research into clinical practice*. Pennsylvania: Lippincott Williams & Wilkins; 2007. [Link]
- [13] Shahraki M, Sohrabi M, Taheri Torbati HR, Nikkiah K, NaeimiKia M. Effect of rhythmic auditory stimulation on gait kinematic parameters of patients with multiple sclerosis. *Journal of Medicine and Life*. 2017; 10(1):33-7. [PMID]
- [14] Memar Moghaddam M, Shahraki M. [The effect of rhythmic auditory stimulation during gait training on kinematic parameters of gait in patients with multiple sclerosis (Persian)]. *Motor Behavior*. 2018; 10(33):149-64. [DOI:10.22089/mbj.2018.5582.1649]
- [15] Shahraki M, Sohrabi M, Torbati H, Nikkiah K, NaeimiKia M. [The effect of rhythmic visual stimulation on kinematic parameters of gait in patients with multiple sclerosis (Persian)]. *Motor Behavior*. 2019; 11(35):35-48. [DOI:10.22089/mbj.2018.1434]
- [16] Rochester L, Baker K, Hetherington V, Jones D, Willems AM, Kwakkel G, et al. Evidence for motor learning in Parkinson's disease: Acquisition, automaticity and retention of cued gait performance after training with external rhythmical cues. *Brain Research*. 2010; 1319:103-11. [DOI:10.1016/j.brainres.2010.01.001] [PMID]
- [17] Dusti F, Frughi Pur M, Sohrabi M, Taheri HR, Saedi M, Teymoori S. [Effect of instructions internal and far and near external focus of attention on balance of multiple sclerosis patients (Persian)]. *Medical Journal of Mashhad University of Medical Sciences*. 2013; 56(1):15-20. [DOI:10.22038/mjms.2013.453]
- [18] Zeni JA Jr, Richards JG, Higginson JS. Two simple methods for determining gait events during treadmill and overground walking using kinematic data. *Gait & Posture*. 2008; 27(4):710-4. [DOI:10.1016/j.gaitpost.2007.07.007] [PMID]
- [19] Barboza TV, Weizemann C, de Carvalho AR. Causal relationship between spatiotemporal parameters of walking and the locomotor rehabilitation index in healthy people. *Gait & Posture*. 2021; 90:320-5. [DOI:10.1016/j.gaitpost.2021.09.191] [PMID]
- [20] Rochester L, Nieuwboer A, Baker K, Hetherington V, Willems AM, Chavret F, et al. The attentional cost of external rhythmical cues and their impact on gait in Parkinson's disease: effect of cue modality and task complexity. *Journal of Neural Transmission*. 2007; 114(10):1243-8. [DOI:10.1007/s00702-007-0756-y] [PMID]
- [21] Cohen S, Williamson GM. Perceived stress in a probability sample of the United States. In: Spacapan S, Oskamp S, editors. *The social psychology of health*. California: Sage; 1988. [Link]

- [22] Correale L, Montomoli C, Bergamaschi R, Ivaniski-Mello A, Peyré-Tartaruga LA, Buzzachera CF. Bilateral coordination of gait at self-selected and fast speed in patients with multiple sclerosis: A case-control study. *Multiple Sclerosis and Related Disorders*. 2022; 65:104027. [DOI:10.1016/j.msard.2022.104027] [PMID]
- [23] Comber L, Galvin R, Coote S. Gait deficits in people with multiple sclerosis: A systematic review and meta-analysis. *Gait & Posture*. 2017; 51:25-35. [DOI:10.1016/j.gaitpost.2016.09.026] [PMID]
- [24] Chee JN, Ye B, Gregor S, Berbrayer D, Mihailidis A, Patterson KK. Influence of multiple sclerosis on spatiotemporal gait parameters: A systematic review and meta-regression. *Archives of Physical Medicine and Rehabilitation*. 2021; 102(9):1801-15. [DOI:10.1016/j.apmr.2020.12.013] [PMID]
- [25] Cameron MH, Nilsagard Y. Balance, gait, and falls in multiple sclerosis. *Handbook of Clinical Neurology*. 2018; 159:237-50. [DOI:10.1016/B978-0-444-63916-5.00015-X] [PMID]
- [26] Peyré-Tartaruga LA, Coertjens M. Locomotion as a powerful model to study integrative physiology: Efficiency, economy, and power relationship. *Frontiers in Physiology*. 2018; 9:1789. [DOI:10.3389/fphys.2018.01789] [PMID]
- [27] Murtagh EM, Mair JL, Aguiar E, Tudor-Locke C, Murphy MH. Outdoor walking speeds of apparently healthy adults: A systematic review and meta-analysis. *Sports Medicine*. 2021; 51(1):125-41. [DOI:10.1007/s40279-020-01351-3] [PMID]
- [28] Araneda R, Renier L, Ebner-Karestinos D, Dricot L, De Volder AG. Hearing, feeling or seeing a beat recruits a supramodal network in the auditory dorsal stream. *European Journal of Neuroscience*. 2017; 45(11):1439-50. [DOI:10.1111/ejn.13349] [PMID]
- [29] Thaut MH, McIntosh GC, Prassas SG, Rice RR. Effect of rhythmic auditory cuing on temporal stride parameters and EMG. Patterns in hemiparetic gait of stroke patients. *Journal of Neurologic Rehabilitation*. 1993; 7(1):9-16. [DOI:10.1177/136140969300700103]
- [30] Wittwer JE, Webster KE, Hill K. Reproducibility of gait variability measures in people with Alzheimer's disease. *Gait & Posture*. 2013; 38(3):507-10. [DOI:10.1016/j.gaitpost.2013.01.021] [PMID]