

Research Article



The Relationship between the Depression and Anxiety Stress Survey Questionnaire, Salivary Cortisol and Heart Rate Variability

Fateme Sangtarash¹ , Haniyeh Choobsaz¹ , Milad Zarrin¹ , Sousan Salari² , Ehsan Mokari Manshadi³ , Ali Akbar Esmaili⁴, Sayyed Hossein Mozaffari⁵ , Boshra Hatef⁵

1. Department of Physical Therapy, School of Rehabilitation, Tehran University of Medical Sciences, Tehran, Iran.

2. Department of Clinical Psychology, Shahed University, Tehran, Iran.

3. Life Style Institute, Behavioral Science Research Center, Baqiyatallah University of Medical Sciences, Tehran, Iran.

4. Cognitive and Behavioral Research Center, Baqiyatallah University of Medical Sciences, Tehran, Iran.

5. Neuroscience Research Center, Baqiyatallah University of Medical Sciences, Tehran, Iran.

Use your device to scan
and read the article online



Citation Sangtarash F, Choobsaz H, Zarrin M, Salari S, Mokari Manshadi E, Esmaili AA, et al. The Relationship between the Depression and Anxiety Stress Survey Questionnaire, Salivary Cortisol and Heart Rate Variability. Journal of Modern Rehabilitation. 2025; 19(1):80-89. <http://dx.doi.org/10.18502/jmr.v19i1.17512>

<http://dx.doi.org/10.18502/jmr.v19i1.17512>

Article info:

Received: 02 Jul 2024

Accepted: 23 Sep 2024

Available Online: 01 Jan 2025

ABSTRACT

Introduction: Cortisol and heart rate variability (HRV) represent the activity of physiological stress axes. The depression and anxiety stress survey (DASS-42) is widely used to assess stress. This study examines the correlations between the DASS-42 questionnaire and stress markers, salivary cortisol levels and HRV.

Materials and Methods: A total of 195 healthy volunteers (145 males and 50 females) participated in this study. At first, the DASS-42 survey form was completed. The salivary cortisol samples were collected, and the electrocardiograms were recorded. Differences in cortisol levels between baseline and post-trier social stress tests were recorded as changes in cortisol after stress (CCAS). Measurements were made to establish whether the overall DASS and its subscales (stress, anxiety and depression) were correlated with baseline cortisol, CCAS, and HRV indices.

Results: The anxiety-DASS subscale score correlated negatively with the CCAS score in women ($r=-0.429$, $P=0.002$). The DASS score was significantly correlated with heart rate ($r=0.25$, $P=0.007$) and SD2 of the Poincare plot ($r=-0.272$, $P=0.004$) in men. In contrast, women showed significant correlations between total DASS scores with very low-frequency power ($r=-0.40$, $P=0.005$) and detrended fluctuation analysis ($r=-0.30$, $P=0.034$). The anxiety-DASS subscale correlated with HRV in both genders.

Conclusion: The anxiety-DASS subscale represents the sympathetic-adrenal medulla (SAM) activity. Clinicians can estimate the activity of this stress axis by using the anxiety subscale of the DASS questionnaire. Meanwhile, gender differences should be noted when assessing stress.

Keywords:

Anxiety; Salivary cortisol;
Heart rate; Nonlinear analysis

*** Corresponding Author:**

Boshra Hatef, Assistant Professor.

Address: Neuroscience Research Center, Baqiyatallah University of Medical Sciences, Tehran, Iran.

Tel: +98 (212) 2382158

E-mail: boshrahatef@bmsu.ac.ir



Copyright © 2025 Tehran University of Medical Sciences. Published by Tehran University of Medical Sciences
This work is licensed under a Creative Commons Attribution-NonCommercial 4.0 International license (<https://creativecommons.org/licenses/by-nc/4.0/>).
Noncommercial uses of the work are permitted, provided the original work is properly cited.

Introduction

Most health researchers agree that stress is critical for human health and aging [1]. However, stress is associated with an increased risk of various diseases, such as cardiovascular disease, hypertension, and even infectious diseases. The measurement of stress, which lacks consistency and thoroughness, is inherently complex [2].

Stress simultaneously activates two axes: The hypothalamus-pituitary-adrenal (HPA) and the sympathetic-adrenal medulla (SAM). The activation of the HPA leads to increased cortisol in body fluid and the SAM activation affects heart function [1, 3]. The correlation between salivary cortisol and plasma cortisol was greater than 0.9 [4]. Therefore, many studies in basic and clinical sciences use salivary cortisol as a biological marker to examine the activity of the physiological nervous or brain stress system.

Another promising stress marker used is heart rate variability (HRV), which is the variation in the length of the heartbeat interval. Time-domain measurements of HRV include assessing the quality of the Mean \pm SD of the RR interval. Frequency-domain measurements of HRV, including the power of the high-frequency (PHF) and the low-frequency (PLF) components, were calculated. In addition to the linear indicators mentioned, nonlinear features include SD1 and SD2 of the Poincaré plot and $\alpha 1$ the detrended fluctuation analysis (DFA) [3, 5].

The depression, anxiety, and stress (DASS-42) questionnaire has been used to assess stress state instead of the physiological responses as stress markers (cortisol and HRV). Parkitny et al. developed DASS-42. It has high reliability and validity (the Cronbach α of 0.96 to 0.97 for DASS-depression, 0.84 to 0.92 for DASS-anxiety, and 0.90 to 0.95 for DASS-stress). The questionnaire consists of 42 items divided into three sections and reports the levels of DASS-42 [6].

Given the variety of methods employed to assess stress, it is possible to establish a link between the results of self-reports and physiological responses to stress. Physiological responses are a potential pathway linking stress to some morbidities or diseases [7]. If this relationship is confirmed, a standard questionnaire could estimate an individual's health status and indicate possible future health risks, potentially saving time and money. Several studies have reported this association [1, 7]. Some of these studies were conducted in children [8] or adoles-

cents [9] with limited sample sizes, and only one study used the DASS-42 questionnaire [10]. The study participants were in specific physiologic states, such as after curettage or an earthquake [9, 11].

The importance of using the DASS questionnaire as an efficient tool for stress assessment will be examined. The first hypothesis of this study is the relationship between DASS scores and physiological biomarkers of stress (salivary cortisol and HRV) in healthy subjects, as HRV and cortisol have been shown to reflect subjects' stress reports. Secondly, HRV and cortisol may be in good agreement.

Materials and Methods

Study participants

The study design is cross-sectional. This study was conducted among healthy volunteers in Tehran City, Iran, from March 2021 until September 2023. After surveying 500 volunteers, 195 were considered eligible and referred to the Neuroscience Research Center Laboratory of [Baqiyatallah University of Medical Sciences](#). They comprised 145 males and 50 females between the ages of 25 and 50.

The inclusion criteria comprised physical and mental health, no smoking habits, no history of craniocervical surgery or mental illness, no history of systemic diseases (such as diabetes or cardiovascular disease) (based on self-report and clinical psychologist comment) and no participation in a regular exercise program in the past month.

Study procedure

After spending 5 minutes in the testing room, the participants completed the DASS-42 questionnaire, and then 5 mL of cortisol saliva between 9-15 O'clock and 2 minutes of ECG was collected from the participants. Testing was performed while all variables at baseline were recorded. Only saliva samples were collected twice (before and after the trier social stress test [TSST]). The TSST protocol consisted of an interview and a math task, which included 5 minutes of speech preparation before entering the testing room, a 2-minute speech performance for two managers and an 8-min math portion. The participants were asked to subtract the number 13 from 1022 successively. The examiner verbally reported the answers aloud and asked them to start in 1022. If the participant made a mistake, the examiner prompted, "You are wrong. Please start over from 1022" [3].

Study outcomes

Depression and anxiety stress-42 questionnaire

The DASS-42 questionnaire was completed in the quiet environment of an exam room. The participants were carefully instructed to complete the questionnaire according to their living conditions. The form was divided into three sections and consisted of 14 questions reporting levels of DASS-42 based on scores obtained on a 5-point scale [10, 12].

Salivary cortisol level

A small saliva sample (0.5 mL) was collected and frozen at -80°C . After thawing, a human saliva cortisol enzyme-linked immunosorbent assay (ELISA) kit from IBL, Germany, was used. The ELISA was performed according to the kit instructions [3].

Electrocardiography (ECG) recording

ECG recording was performed using equipment from Rib Intelligence Technology, Iran. Three electrodes were used in this study. The first electrode was placed on the midline to the left of the heart. The second electrode was placed in the right lower abdomen, and the third electrode was placed on the left edge of the sternum, below the heart position. HRV can be affected by many factors, including circadian rhythms, body position, level of physical activity before testing, medication intake, speech impairment, and deep breathing.

For this reason, special care was taken to ensure that all users were under the same conditions. ECGs were recorded at the same time of day. During the study, the subjects were in a sitting position without deep breathing or talking. ECG recordings were performed for 2 minutes using standard methods [3, 13]. Johnell developed HRV software, version 2010 in MATLAB software, version 2010 [8]. It was used for HRV analysis. At first, the HRV was extracted from the ECG signal by appropriate filtering and peaks were identified during the analysis process. Since the ECG signal is bandpass filtered between 5 and 20 Hz, the peak R can be estimated by setting the amplitude threshold value and the minimum time interval between consecutive peaks. In the time domain, the Mean \pm SD of the RR were extracted as linear features from the RR series. In the frequency domain, high frequency (HF) power (0.15-0.5 Hz), low frequency (LF) power (0.05-0.15 Hz), ultra-LF power (0-0.04) and the ratio of LF/HF were analyzed [5]. The nonlinear features extracted in the time domain are SD1

and SD2 from Poincaré plots and $\alpha 1$ from DFA and in the frequency domain, spectral entropy (SE) [3, 14].

Statistical analysis

After the data were collected, they were analyzed using the SPSS software, version 25. Since the data were normally distributed based on the Kolmogorov–Smirnov test results, the Pearson correlation coefficient was used to determine the relationships among the variables. Curve estimation was used to determine the nonlinear correlation between cortisol and the frequency domain of HRV (percentage of VLF and HF). The significance level for all the tests was set at $P < 0.05$.

Results

A total of 195 males and females participated in the study. Table 1 shows the Mean \pm SD and sex comparisons of the baseline data. There were no significant differences in the DASS depression score, DASS anxiety score, or heart rate between men and women. In contrast, the DASS-total and DASS-stress scores were significantly greater in men, and the baseline cortisol level was significantly greater in women (Table 1).

Relationships of the DASS-42 total score and its subscale scores with salivary cortisol

There was no significant relationship between the DASS total score and baseline cortisol or changes in cortisol after stress (CCAS) scores in men and women. There was no relationship between the three DASS subscales and salivary cortisol in men; however, in women, only the DASS-anxiety subscale showed a significant negative correlation with the CCAS score ($r = -0.429$, $P = 0.002$) (Table 2).

Relationships between the DASS-42-total score and its subscale scores and HRV

For men, the DASS total score showed a statistically significant positive correlation with heart rate ($r = 0.250$ and $P = 0.007$) and a statistically significant negative correlation with Poincaré plot-SD2 ($r = -0.272$ and $P = 0.004$). The depression subscale score did not significantly correlate with HRV. The anxiety subscale showed a significant positive correlation with HR ($r = 0.282$, $P = 0.002$) and a significant negative correlation with the mean RR ($r = -0.25$, $P = 0.006$) and Poincaré plot-SD2 ($r = -0.291$, $P = 0.002$). The stress subscale showed a significant negative correlation with Poincaré plot-SD2 ($r = -0.218$, $P = 0.002$) and a positive correlation with SE ($r = 0.214$,

Table 1. Comparing Mean±SD of DASS scores and baseline cortisol levels between the two genders

Variables	Mean±SD		P
	Men	Women	
DASS	33.37±23.02	25.46±20.97	0.02*
DASS-depression	8.90±8.62	7.86±8.09	0.44
DASS-anxiety	8.12±6.66	7.06±6.21	0.30
DASS-stress	13.62±9.39	10.42±8.73	0.03*
Baseline-cortisol	6.74±4.02	13.60±6.61	0.00*
Heart rate	78.73±11.58	80.66±11.10	0.31

DASS: Depression and anxiety stress survey.

JMR

*Significant difference based on the t-test.

P=0.027) (Table 3). There was no significant correlation between DASS and its subscales with sample entropy and DFA-α1 of HRV.

In women, the total DASS score positively correlated with the PVLf (r=0.405, P=0.005) and DFA-α1 (r=0.309, P=0.034). Depression subscale scores were significantly positively correlated with the PVLf (r=0.422, P=0.003)

Table 2. DASS scores relationship with CCAS

Gender	Variables	Statistics	DASS	DASS-Depression	DASS-Anxiety	DASS-stress
Men	Cortisol	R	-0.040	-0.106	-0.007	-0.147
		P	0.636	0.210	0.933	0.081
	CCAS	R	-0.095	-0.117	-0.120	-0.129
		P	0.292	0.192	0.183	0.151
Women	Cortisol	R	0.115	0.158	0.128	0.142
		P	0.427	0.272	0.368	0.326
	CCAS	R	-0.194	-0.041	-0.429*&	-0.211
		P	0.178	0.780	0.002*&	0.141

DASS: Depression and anxiety stress survey; CCAS: Changes in cortisol after stress.

JMR

&Significant; *Significant based on the Pearson correlation.

Table 3. DASS scores relationship With HRV in men

Variables		Mean RR	SD	HR	PVLf	PLF	PHF	LF/HF	Poincare plot -SD1	Poincare plot -SD2	SE
DASS	R	-0.230	-0.058	0.250*&	0.106	-0.80	-0.038	-0.015	-0.065	-0.272*&	0.177
	P	0.014	0.537	0.007*&	0.261	0.398	0.686	0.877	0.490	0.004*&	0.068
DASS-depression	R	-0.118	0.028	0.143	0.035	0.083	0.036	-0.087	0.024	-0.094	0.026
	P	0.210	0.768	0.128	0.711	0.384	0.704	0.369	0.797	0.329	0.789
DASS-anxiety	R	-0.255*&	-0.110	0.282*&	0.106	-0.50	-0.064	0.019	-0.104	-0.291*&	0.130
	P	0.006*&	0.244	0.002*&	0.262	0.596	0.498	0.845	0.272	0.002*&	0.183
DASS-stress	R	-0.156	-0.065	0.167	0.048	-0.120	0.055	-0.103	-0.016	-0.218*&	0.214*&
	P	0.096	0.493	0.076	0.615	0.205	0.562	0.287	0.862	0.022*&	0.027*&

JMR

Abbreviations: DASS: Depression and anxiety stress survey; HR: Heart rate; PVLf: Very low-frequency power; PLF: Low-frequency power; PHF: High-frequency power; LF/HF: Low frequency/high frequency; SE: Spectral entropy.

&Significant; *Significant based on the Pearson correlation.

Table 4. DASS scores relationship with HRV in women

Variables		Mean±SD	HR	PVLF	PLF	PHF	LFHF	Poincare plot -SD1	Poincare plot -SD2	DFA -α1
DASS	R	-0.105±0.189	0.095	0.405* ^{&}	0.069	-0.281	0.142	-0.284	-0.080	0.309* ^{&}
	P	0.483±0.202	0.526	0.005 ^{&}	0.647	0.059	0.345	0.053	0.594	0.034 ^{&}
DASS-depression	R	-0.107±0.197	0.085	0.422* ^{&}	0.111	-0.324* ^{&}	0.202*	-0.282	-0.107	0.388* ^{&}
	P	0.471±0.181	0.564	0.003 ^{&}	0.459	0.026* ^{&}	0.174*	0.052	0.469	0.019 ^{&}
DASS-anxiety	R	-0.102±0.193	0.122	0.366* ^{&}	0.126	-0.281	0.120	-0.252	-0.115	0.291* ^{&}
	P	0.487±0.183	0.403	0.020 ^{&}	0.393	0.053	0.417	0.080	0.430	0.043 ^{&}
DASS-stress	R	-0.059±0.109	0.049	0.390* ^{&}	0.011	-0.230	0.115	-0.229	0.019	0.242
	P	0.691±0.461	0.739	0.007 ^{&}	0.941	0.120	0.441	0.117	0.900	0.097

JMR

Abbreviations: DASS: Depression, and anxiety stress survey; SD: Standard deviation; HR: Heart rate; PVLF: Very low-frequency power; PLF: Low-frequency power; PHF: High-frequency power; LF/HF: Low frequency/high frequency; Sample-ENT: Sample entropy; DFA: Detrended fluctuation analysis; Spect-ENT: Spectral entropy.

*Significant; [&]Significant Pearson correlation

and DFA-α1 (r=0.338, P=0.019) and significantly negatively correlated with the PHF (r=-0.324, P=0.003). Anxiety significantly correlated with PVLF (r=0.336, P=0.020) and DFA-α1 (r=0.291, P=0.043). The stress component positively correlated with only the PVLF (r=0.390, P=0.07) (Table 4). There was no significant correlation between DASS and its subscales with sample entropy and SE of HRV.

Relationship between salivary cortisol and HRV

In men, the baseline cortisol only showed a significant negative correlation with the SD of RR (r=-0.189, P=0.048). In women, the baseline cortisol was significantly related to the PVLF (r=0.309, P=0.031) and DFA-α1 (r=0.296, P=0.037) (Table 5). An inverted regression was observed between salivary cortisol and both the PVLF (r=0.361, P=0.00003) and PHF (r=0.233, P=0.003) in all subjects (both sexes) (Figures 1 and 2).

Discussion

This study determined the possible relationships between self-reported measures of stress, the DASS-42 questionnaire, and physiological biomarkers. Our results showed no association between total scores on the DASS or its subscale and baseline cortisol or CCAS levels. However, the DASS-anxiety subscale score was inversely related to CCAS in women. Consequently, two HRV characteristics (HR and Poincare plot-SD2 in men) and (PVLF and DFA-α1 in women) correlated with the total score on the DASS. In both men and women, the DASS-anxiety subscale score was correlated with the HRV score.

Comparing the DASS-42 score with the baseline cortisol concentration and CCAS score

The present study analyzed baseline data from women and men and revealed that baseline cortisol levels in women were approximately twice as high as in men.


Table 5. Relation between salivary cortisol and HRV

Gender	Variables	Statistics	Mean	SD RR	HR	PVLF	PLF	PHF	Sample-ENT	DFA -α1	SE
Men	Cortisol	R	-0.028	-0.186* ^{&}	0.052	0.120	-0.011	-0.113	-0.049	-0.134	0.018
		P	0.769	0.048* ^{&}	0.584	0.202	0.905	0.234	0.640	0.162	0.858
Women	Cortisol	R	-0.097	0.022	0.130	0.309* ^{&}	0.132	-0.280	-0.141 ^{&}	0.296	-0.089
		P	0.502	0.881	0.369	0.031 ^{&}	0.367	0.051	0.0412* ^{&}	0.037	0.700

JMR


Abbreviations: HR: Heart rate; PVLF: Very low-frequency power; PLF: Low-frequency power; PHF: High-frequency power; LF/HF: Low frequency/high frequency; Sample-ENT: Sample entropy; DFA: Detrended fluctuation analysis; SE: Spectral entropy.

*Significant.



JMR

Figure 1. The inverted nonlinear regression between the amount of salivary cortisol and the very low-frequency power in all subjects (n=195; $r=0.361$, $P=0.00003$)



JMR

Figure 2. The inverted nonlinear regression between the amount of salivary cortisol and high-frequency power in all subjects (n=195; $r=0.233$, $P=0.003$)

This result is consistent with previous studies showing that women's baseline cortisol levels are higher than men's [16, 17]. Overall, women had significantly higher morning cortisol levels than men, and women under 50 years of age had higher morning and Δ -cortisol (morning and evening cortisol) levels than matched men [16]. In contrast, in a study conducted among different sexes, serum cortisol levels were greater in depressed men than in depressed women [18]. The results of this study differ in that the participants were healthy, and cortisol was determined in saliva rather than in serum. Salivary cortisol sampling is not stressful or inconvenient for participants and several samplings are possible.

In addition, absolute cortisol and CCAS values were unrelated to the total DASS or DASS subscales, except for the anxiety-DASS subscale in women, which negatively correlated with the CCAS. This study's results are consistent with previous research [19, 20], in which they showed that absolute cortisol levels do not correlate with emotional self-reports of distress anxiety. In contrast, Sungako et al. reported a significant relationship between DASS scores and salivary cortisol levels in women after curettage [10]. These results may be related to the characteristics of the participants examined after curettage, where hormonal fluctuations are high. On the other hand, some studies have suggested a circadian rhythm of cortisol that the lowest cortisol levels occur around midnight and that the peak of cortisol production occurs in the early morning. They determined the cortisol arousal response and estimated the diurnal cortisol levels by sampling several times during the day [20, 21]. It was impossible to evaluate these items in this study. Meanwhile, the current results were observed in healthy young people and might differ from those in other age groups.

Our study showed that the DASS scores of men were significantly greater than women. On the other hand, among employees of engineering colleges in Nigeria, no significant differences were found in DASS-42 scores [8] between males and females. In Iran, no study has compared the performance of the DASS-42 and its subset between males and females. The subjective questionnaire can explain this result; men in Iran face more work-related stresses, and in most cases, men are the only working members of the family [22].

DASS-42 data versus HRV

Theoretically, HRV may be affected by behavioral factors, such as social and conduct disorders [23]. According to our results, some features of HRV are related to DASS-42.

HRV characteristics, such as HR and Pioncare-SD2, showed the strongest correlation with the DASS-42 questionnaire in men. Nonlinear HRV characteristics, such as Pioncare-SD2, indicate the heart's dynamic state. The main advantage of studying nonlinear characteristics is that it provides complete and natural information about biological systems [24]. No previous study using nonlinear features, such as the Poincaré-SD2 score, determined their correlation with self-report questionnaire scores.

In addition to the Pioncare-SD2 and HR subscales, the DASS-anxiety subscale was correlated with the mean RR. In contrast, some studies did not observe significant changes in HRV after acute stress in men [3, 15, 16]. In men, the time domain features of the HRV were correlated with those of the DASS-42.

In contrast, in women, the frequency domain of the HRV was related to the DASS-42 score and its subset. Several frequency HRV parameters have been used to test which of these parameters is most strongly associated with stress, and they represent high frequency, which can reflect vagal parasympathetic innervation. LF is a dual innervation of sympathetic activity and vagal innervation (PA+SA), and the low-frequency/high-frequency ratio has been assumed to represent the sympathy-vagal balance [24]. Moreover, anxiety is related to both low PA and high SA in adults and children [25]. As expected, our research revealed a direct relationship between PLF and DFA- α 1 and an inverse relationship with PHF. This result aligns with that of Mohammadi et al. who reported changes in the frequency of HRV after stress in women [3].

The relationships between the DASS-42 and HRV showed that only two subscales, the DASS-anxiety and DASS-stress subscales, were related to HRV in men, but all three subscales of the DASS-42 were related to HRV in women. Underlying mechanisms could be the sex differences that have been demonstrated in the used HRV parameters [26] in psychological functioning and development of genders since childhood [27] and in handling stressful situations [27].

For both genders, the DASS-anxiety subscale is related to HRV. The DASS-anxiety subscale questions focus more on physiological changes following stress. This section examines the state of breathing, the amount of sweating, the irregular heart rhythm, and the feeling of weakness. This subset seems to have concentrated more on subjective stress perception and coping with it [28]. Then, it may be suggested that the DASS questionnaire evaluates the SAM axis of stress rather than the HPA axis.

Cortisol versus HRV

HRV and cortisol have been shown to reflect physiological changes after stress. We investigated their relationship. For the first time, we also consider differences between the sexes.

A low HRV (a greater sympathetic over parasympathetic dominance) is related to a greater cortisol awakening response [29]. In another study, cortisol waking response was associated with low high frequency and LF, but there was no cross-sectional change in LF/high frequency [30]. Nevertheless, no significant relationship between cortisol and HRV has been reported [31, 32].

In this study, baseline cortisol showed a negative correlation with the SD of the RR in men and a positive correlation with the PVLf in women. A previous study showed that baseline cortisol has a negative correlation with the standard deviation (SD) of the RR and a positive correlation with LF without mentioning differences between men and women [3].

The relationship between cortisol and the HRV index shows the real interaction of the HPA and SAM axes in the heart. When the cortisol concentration increases to the desired level, the VLF and HF indexes decrease. Thus, at very low and very high suboptimal cortisol levels, the parasympathetic nervous system activity index, i.e. HF, will be above the optimal state, and the sympathetic nervous system activity index, i.e. VLP, will be below the optimal state based on cortisol levels.

Despite this relationship between HRV and cortisol, their responses to acute stress may differ. For example, it has been reported that there is a dissociation in the response of both nervous systems to repeated exposure to stress and that after several repeated exposures (habituation), the stress response of cortisol decreases while the HRV response remains high such as some disorders, multiple sclerosis, and stroke [13, 14, 33]. In addition, salivary cortisol levels continued to increase under stress conditions but rapidly decreased under control conditions. Shaverdi et al. demonstrated that SDs of Poincaré plots of HRV increased, and the salivary cortisol changes tended to be in the optimal range after Muslim praying. This means the changes in HRV and cortisol are the same but do not have a linear correlation [34, 35].

Conclusion

The present study showed that only DASS-anxiety was associated with CCAS in women. The time domain of HRV was related to the DASS score in men and the frequency domain of HRV in women. This study showed that the DASS-anxiety subscale reflects SAM activity in both men and women. Therefore, clinicians can make appropriate decisions regarding this stress axis by evaluating the DASS questionnaire, particularly the anxiety subscale. In addition, cortisol is associated with several characteristics of HRV.

Study limitations

One of the limitations of our study was that the number of male and female participants was not the same. The other limitation was the cortisol awakening response and diurnal decline, and their relationships with DASS and HRV were not investigated.

Ethical Considerations

Compliance with ethical guidelines

This study was approved by the Ethics Committee of the [Baqiyatallah University of Medical Sciences](#) (Code: IR. BMSU. REC.1400.051).

Funding

This research did not receive any grant from funding agencies in the public, commercial, or non-profit sectors.

Authors' contributions

Conceptualization and study design: Boshra Hatef, Sayyed Hossein Mozaffari, Aliakbar Esmacili and Ehsan Mokari Manshadi; Data collection: Boshra Hatef, Sou-san Salari, Fateme Sangtarash, Haniyeh Choobsaz and Milad Zarrin; Data analysis: Fateme Sangtarash, Sayyed Hossein Mozaffari, Haniyeh Choobsaz, Fateme Sangtarash and Boshra Hatef; Drafting the manuscript, review and final approval: All authors.

Conflict of interest

The authors declared no conflict of interest.

Acknowledgments

The authors thank the laboratory of the Neuroscience Research Center of [Baqiyatallah University of Medical Sciences](#) and Gholamreza Akbari.

Informed consent

Informed consent was obtained from all individual participants included in the study.

Data Availability Statement

The data supporting this study's findings are available upon request from the corresponding author.

References

- [1] Yaribeygi H, Panahi Y, Sahraei H, Johnston TP, Sahebkar A. The impact of stress on body function: A review. *EXCLI Journal*. 2017; 16:1057-72. [PMID]
- [2] Hovsepian K, al'Absi M, Ertin E, Kamarck T, Nakajima M, Kumar S. cStress: Towards a gold standard for continuous stress assessment in the mobile environment. *Proceedings of the ... ACM International Conference on Ubiquitous Computing . UbiComp (Conference)*. 2015; 2015:493-504. [DOI:10.1145/2750858.2807526] [PMID]
- [3] Mohammadi A, Emamgoli A, Shirinkalam M, Meftahi GH, Yagoobi K, Hatef B. The persistent effect of acute psychosocial stress on heart rate variability. *The Egyptian Heart Journal*. 2019; 71(1):18. [DOI:10.1186/s43044-019-0009-z] [PMID]
- [4] Altamura M, Iuso S, Balzotti A, Francavilla G, Dimitri A, Cibelli G, et al. Salivary alpha-amylase and cortisol responsiveness to stress in first episode, drug-naive patients with panic disorder. *Neuroscience Research*. 2018; 137:49-56. [DOI:10.1016/j.neures.2018.03.003] [PMID]
- [5] Billman GE. The LF/HF ratio does not accurately measure cardiac sympatho-vagal balance. *Frontiers in Physiology*. 2013; 4:26. [DOI:10.3389/fphys.2013.00026]
- [6] Parkitny L, McAuley J. The depression anxiety stress scale (DASS). *Journal of Physiotherapy*. 2010; 56(3):204. [DOI:10.1016/S1836-9553(10)70030-8] [PMID]
- [7] Chandola T, Heraclides A, Kumari M. Psychophysiological biomarkers of workplace stressors. *Neuroscience & Biobehavioral Reviews*. 2010; 35(1):51-7. [DOI:10.1016/j.neubiorev.2009.11.005] [PMID]
- [8] Johnell O, Kanis JA. An estimate of the worldwide prevalence and disability associated with osteoporotic fractures. *Osteoporosis International*. 2006; 17(12):1726-33. [DOI:10.1007/s00198-006-0172-4] [PMID]
- [9] Rondó PH, Vaz AJ, Moraes F, Tomkins A. The relationship between salivary cortisol concentrations and anxiety in adolescent and non-adolescent pregnant women. *Brazilian Journal of Medical and Biological Research*. 2004; 37(9):1403-9. [DOI:10.1590/S0100-879X2004000900016] [PMID]
- [10] Sungkono MH, Nooryanto M, Arianti DE. Association of social support with Depression and Anxiety Stress Score (DASS) and salivary cortisol level in women post curettage due to first-trimester miscarriage. *The Journal of Obstetrics and Gynecology Science*. 2020; 1(1):18-22. [Link]
- [11] Kotozaki Y, Kawashima R. Effects of the Higashi-Nihon earthquake: Posttraumatic stress, psychological changes, and cortisol levels of survivors. *Plos One*. 2012; 7(4):e34612. [DOI:10.1371/journal.pone.0034612] [PMID]
- [12] Sahebi A, Asghari MJ, Salari RS. [Validation of depression anxiety and stress scale (DASS-21) (Persian)]. *Developmental Psychology*. 2006; 1(4):299. [Link]
- [13] Mirzaee O, Saneian M, Vani JR, Shahrivar K, Peyravi M, Shariat A, et al. The psychophysiological responses of the chronic ischemic stroke patients to the acute stress were changed. *Brazilian Archives of Biology and Technology*. 2019; 62. [DOI:10.1590/1678-4324-2019180494]
- [14] Ashtiani AA, Shaygannejad V, Ghobadi F, Bathaie R, Shahyad S, Hatef B. Acute response of stress system in multiple sclerosis. *Archives of Neuroscience*. 2021; 8(4):e115781. [DOI:10.5812/ans.115781]
- [15] Vierhapper H, Nowotny P, Waldhäusl W. Sex-specific differences in cortisol production rates in humans. *Metabolism*. 1998; 47(8):974-6. [DOI:10.1016/S0026-0495(98)90353-5]
- [16] Larsson CA, Gullberg B, Råstam L, Lindblad U. Salivary cortisol differs with age and sex and shows inverse associations with WHR in Swedish women: A cross-sectional study. *BMC Endocrine Disorders*. 2009; 9:16. [DOI:10.1186/1472-6823-9-16] [PMID]
- [17] McCormick CM, Lewis E, Somley B, Kahan TA. Individual differences in cortisol levels and performance on a test of executive function in men and women. *Physiology & Behavior*. 2007; 91(1):87-94. [DOI:10.1016/j.physbeh.2007.01.020] [PMID]
- [18] Druzhkova T, Pochigaeva K, Yakovlev AA, Gersamia A, Guekht A, Gulyaeva N. Effects of childhood trauma on the biological correlates of stress in men and women with borderline mental disorders. *Neuroscience and Behavioral Physiology*. 2019; 49(7):916-20. [DOI:10.1007/s11055-019-00819-8]
- [19] Takahashi T, Ikeda K, Ishikawa M, Kitamura N, Tsukasaki T, Nakama D, et al. Anxiety, reactivity, and social stress-induced cortisol elevation in humans. *Neuroendocrinology Letters*. 2005; 26(4):351-4. [Link]
- [20] Fries E, Dettenborn L, Kirschbaum C. The cortisol awakening response (CAR): Facts and future directions. *International Journal of Psychophysiology*. 2009; 72(1):67-73. [DOI:10.1016/j.ijpsycho.2008.03.014] [PMID]
- [21] Heim C, Ehlert U, Hellhammer DH. The potential role of hypocortisolism in the pathophysiology of stress-related bodily disorders. *Psychoneuroendocrinology*. 2000; 25(1):1-35. [DOI:10.1016/S0306-4530(99)00035-9] [PMID]
- [22] Azad-Marzabadi E, Gholami Fesharaki M. Job stress and related factors among iranian male staff using a path analysis model. *Iranian Red Crescent Medical Journal*. 2016; 18(6):e34314. [DOI:10.5812/ircmj.34314] [PMID]
- [23] Beauchaine T. Vagal tone, development, and Gray's motivational theory: Toward an integrated model of autonomic nervous system functioning in psychopathology. *Development and Psychopathology*. 2001; 13(2):183-214. [DOI:10.1017/S0954579401002012] [PMID]

- [24] Electrophysiology TF. Heart rate variability: Standards of measurement, physiological interpretation, and clinical use. *Circulation*. 1996; 93(5):1043-65. [DOI:10.1161/01.CIR.93.5.1043]
- [25] Friedman BH. An autonomic flexibility-neurovisceral integration model of anxiety and cardiac vagal tone. *Biological Psychology*. 2007; 74(2):185-99. [DOI:10.1016/j.biopsycho.2005.08.009] [PMID]
- [26] Michels N, Sioen I, Huybrechts I, Bammann K, Vanaelst B, De Vriendt T, et al. Negative life events, emotions and psychological difficulties as determinants of salivary cortisol in Belgian primary school children. *Psychoneuroendocrinology*. 2012; 37(9):1506-15. [DOI:10.1016/j.psyneuen.2012.02.004] [PMID]
- [27] Crick NR, Zahn-Waxler C. The development of psychopathology in females and males: Current progress and future challenges. *Development and Psychopathology*. 2003; 15(3):719-42. [DOI:10.1017/S095457940300035X] [PMID]
- [28] Szabo M, Lovibond PF. Development and psychometric properties of the DASS-Youth (DASS-Y): An extension of the depression anxiety stress scales (DASS) to adolescents and children. *Frontiers in Psychology*. 2022; 13:766890. [DOI:10.3389/fpsyg.2022.766890] [PMID]
- [29] Michels N, Sioen I, Clays E, De Buyzere M, Ahrens W, Huybrechts I, et al. Children's heart rate variability as stress indicator: Association with reported stress and cortisol. *Biological Psychology*. 2013; 94(2):433-40. [DOI:10.1016/j.biopsycho.2013.08.005] [PMID]
- [30] Chen C, Jin Y, Lo IL, Zhao H, Sun B, Zhao Q, et al. Complexity change in cardiovascular disease. *International Journal of Biological Sciences*. 2017; 13(10):1320-8. [DOI:10.7150/ijbs.19462] [PMID]
- [31] Gunnar MR, Porter FL, Wolf CM, Rigatuso J, Larson MC. Neonatal stress reactivity: Predictions to later emotional temperament. *Child development*. 1995; 66(1):1-13. [DOI:10.1111/j.1467-8624.1995.tb00851.x] [PMID]
- [32] Fyfe-Johnson AL, Muller CJ, Alonso A, Folsom AR, Gottesman RF, Rosamond WD, et al. Heart rate variability and incident stroke: The atherosclerosis risk in communities study. *Stroke*. 2016; 47(6):1452-8. [DOI:10.1161/STROKEAHA.116.012662] [PMID]
- [33] Schommer NC, Hellhammer DH, Kirschbaum C. Dissociation between reactivity of the hypothalamus-pituitary-adrenal axis and the sympathetic-adrenal-medullary system to repeated psychosocial stress. *Psychosomatic Medicine*. 2003; 65(3):450-60. [DOI:10.1097/01.PSY.0000035721.12441.17] [PMID]
- [34] Shaverdi Y, Jahromi GP, Meftahi GH, Sharif MS, Mojabi N, Hatf B. The effect of Islamic praying on the heart rate variation in the adult Muslims. *Journal of Modern Rehabilitation*. 2024; 18(2):179-88. [DOI:10.18502/jmr.v18i2.15974]
- [35] Sobhani V, Mokari EM, Aghajani J, Hatf B. Islamic praying changes stress-related hormones and genes. *Journal of Medicine and Life*. 2022; 15(4):483-8. [DOI:10.25122/jml-2021-0167] [PMID]