

Review Article

Effects of Vitamins on Auditory and Vestibular Systems:
Narrative ReviewAylin Mellati¹, Nasrin Gohari^{1*}, Nayiere Mansouri², Mona Mansouri³

1. Department of Audiology, School of Rehabilitation Sciences, Hamadan University of Medical Sciences, Hamadan, Iran.

2. Department of Audiology, Faculty of Rehabilitation, Tabriz University of Medical Sciences, Tabriz, Iran.

3. Department of Medical Laboratory Sciences, Vahabzadeh Laboratory, Ardabil, Iran.

Use your device to scan
and read the article online**Citation** Mellati A, Gohari N, Mansouri N, Mansouri M. Effects of Vitamins on Auditory and Vestibular Systems: Narrative Review. Journal of Modern Rehabilitation. 2026; 20(3):256-267. <http://dx.doi.org/10.18502/jmr.v20i3.22528>**doi** <http://dx.doi.org/10.18502/jmr.v20i3.22528>

Article info:

Received: 24 Sep 2025**Accepted:** 11 Nov 2025**Available Online:** 01 Jul 2026

Keywords:

Vitamin; Auditory; Vestibular

ABSTRACT**Introduction:** Hearing and balance are vital body functions that allow communication with our surroundings and effective participation in everyday activities. Various factors, including vitamins, influence these two functions. This study aimed to examine the role of vitamins in hearing and balance and the effects of deficiencies in these nutrients on ear and balance health.**Materials and Methods:** This review examined the effects of vitamins on hearing. We identified 71 studies published since 2000 through searches of Medline (PubMed), Google Scholar, ScienceDirect, EBSCO, ASHA, Thieme, and ProQuest.**Results:** Evidence suggests that Vitamins A, B, C, D, and E are associated with auditory and vestibular health. Vitamin deficiencies have been linked to hearing loss, tinnitus, dizziness, and impaired inner ear function. In particular, vitamins A and E support hair cell integrity and protect against oxidative stress, while vitamins B and C contribute to neural function. Vitamin D is also important for vestibular health through its role in calcium regulation.**Conclusion:** Ensuring adequate intake of these vitamins through diet or supplementation may help prevent hearing and balance disorders and improve quality of life

* Corresponding Author:

Nasrin Gohari, Assistant Professor.

Address: Department of Audiology, School of Rehabilitation Sciences, Hamadan University of Medical Sciences, Hamadan, Iran.

Tel: +98 (912) 3327573

E-mail: rasacenter@yahoo.com, n.gohari@umsha.ac.irCopyright © 2026 Tehran University of Medical Sciences. Published by Tehran University of Medical Sciences
This work is licensed under a Creative Commons Attribution-NonCommercial 4.0 International license(<https://creativecommons.org/licenses/by-nc/4.0/>).
Noncommercial uses of the work are permitted, provided the original work is properly cited.

Introduction

Vitamins are organic compounds that play a crucial role in vital body processes, including normal cell function, growth, and development. These nutrients are integral to human health and must be supplied in small amounts through the diet for proper metabolic functions [1]. They are divided into two main categories: water-soluble and fat-soluble vitamins. Water-soluble vitamins: These vitamins dissolve in water and are not usually stored in the body, except vitamin B12, which is stored in large quantities in the liver. For this reason, they need to be consumed regularly. Water-soluble vitamins include vitamin C and the B vitamins, which include: thiamine (B1), riboflavin (B2), niacin (B3), pantothenic acid (B5), pyridoxine (B6), biotin (B7), folate (B9), and cobalamin (B12) [2]. Fat-soluble vitamins are vitamins that dissolve in fat and can be stored in adipose tissue and the liver. Vitamins A, D, E, and K are among these [3].

Vitamins play crucial roles in systemic physiology; each vitamin acts through specific mechanisms in different organs, and deficiencies can have significant clinical consequences. It is worth noting that free radicals are highly reactive because they have unpaired electrons; therefore, they can abstract electrons from healthy cells to balance themselves, thereby damaging them. According to studies, vitamins and minerals, with their antioxidant effects, can prevent the destructive effects of free radicals produced during various processes [1, 2].

Vitamins help regulate immune function and maintain overall health and play a role in improving immune function and reducing damage from infections [1, 2]. Studies have shown that vitamins play an important role in maintaining the health of inner hair cells and the function of the auditory nerve and are effective in improving the function of sensory cells in the inner ear. Vitamin deficiencies can lead to hearing loss, tinnitus, and dizziness [3-5].

Since hearing and balance are fundamental human senses that play a vital role in communication, learning, language development, social interactions, maintaining body position, and spatial orientation, respectively, and any disorder in the hearing system and balance can be associated with numerous consequences, including impaired communication, impaired speech understanding [6, 7], social isolation, emotional disorders such as depression and isolation [8], impaired quality of life [9], impaired postural stability, and increased risk of falling

[10]. On the other hand, vitamins are among the factors affecting the hearing and balance systems, and their deficiency can contribute to the occurrence or exacerbation of disorders in both systems. Therefore, in this study, we conducted a comprehensive review of existing evidence on the effects of vitamins on the hearing and balance systems.

Materials and Methods

We conducted a comprehensive search across multiple databases, including MEDLINE (PubMed), Google Scholar, ScienceDirect, EBSCO, ASHA, Thieme, and ProQuest, without imposing any date restrictions. The search keywords included: "Vitamins," "auditory system," "vestibular system," "tinnitus," "vertigo," and "sensorineural hearing loss." In addition to the database search, we examined the reference lists of the selected articles to ensure inclusion of all relevant studies.

All identified papers were reviewed, and a subset was selected based on the following inclusion criteria: Availability of an abstract or full text, relevance to the auditory and vestibular systems, and English language publication.

Initially, 106 relevant articles were identified. From these, 71 papers most directly related to vitamins and the auditory/vestibular systems were selected in accordance with the inclusion criteria (Figure 1). This rigorous search and selection process allowed us to compile a robust body of literature, providing a solid foundation for understanding the impact of vitamins on the health of the auditory and vestibular systems.

Results

The role of vitamins B, C, A, D, and E in the health and function of the hearing and balance systems has been proven (Figure 2). The effects of these vitamins on these systems will be described below.

Water-soluble vitamins

Vitamins B

There are several types of vitamins in the vitamin B group. They are well-known water-soluble nutrients that are essential for vital physiological functions and play an important role in maintaining optimal health [11]. These vitamins are found in animal proteins, dairy products, leafy vegetables, and legumes and play a role in energy metabolism and nerve function [12]. While vitamin B1

(thiamine) is essential for the production of energy from carbohydrates, its deficiency can lead to disorders of the nervous system, digestion, and heart function [13]. Vitamin B2 (riboflavin) deficiency can increase the risk of anemia, migraines, and cataracts [14]. In addition, vitamin B3 (niacin) deficiency can lead to pellagra and various health problems [15]. Vitamin B5 (pantothenic acid) deficiency can disrupt the use of carbohydrates, fats, and proteins for energy and impair hormone production and immune function [16]. Vitamin B6 (pyridoxine) is vital for red blood cell production, carbohydrate metabolism, liver detoxification, and brain health [17], and its deficiency can lead to anemia, depression, and chronic inflammatory conditions [18]. Vitamin B9 (folate) plays a vital role in brain health and the production of neurotransmitters [19]. Its deficiency can increase the risk of depression and cancer [20]. Moreover, vitamin B12 deficiency can lead to anemia, neuropathy, and various neurological disorders [21].

Among the B vitamins, the roles of vitamins B12, B9, and B6 in the auditory system and balance are more apparent. Deficiency of B vitamins can cause neurological symptoms, dizziness, and tinnitus [22]. Studies have shown that vitamin B12 deficiency can negatively affect cochlear function. For example, one comparative study found that the amplitude of TEOAE and SOAE was significantly lower in a group with vitamin B12 deficiency than in a control group [23]. Evidence also showed that 63% of patients with tinnitus had low B12 levels. However, vitamin B12 replacement therapy did not significantly improve tinnitus symptoms [24]. In addition, studies have shown that vitamins B12 and B9 can reduce the risk of low-frequency hearing loss [10, 25]. In patients with NIHL, the prevalence of vitamin B12 deficiency was higher, and serum vitamin B12 levels were significantly lower than in healthy subjects [10].

In a study by Özdemir et al. patients with vitamin B12 deficiency showed a higher rate of absent VEMP waves and reduced amplitudes with normal latencies, indicating dysfunction in the vestibular system [26]. Meanwhile, vitamin B6 has been recognized as an effective agent for treating vertigo. Research suggests that this vitamin plays an important role in maintaining neurological health and may help reduce vertigo-related symptoms [22].

Although vitamin B12 deficiency is frequently observed in individuals with tinnitus and hearing loss, clinical outcomes following supplementation remain inconsistent. For instance, Rodrigues et al. (2025) reported that although B12 deficiency is significantly associated

with auditory dysfunction, replacement therapy alone does not always result in symptomatic improvement. This discrepancy may be attributed to several factors, including irreversible neural degeneration caused by prolonged deficiency, variability in treatment dosage and duration, and methodological limitations in study design, such as small sample sizes or the absence of a placebo control. Moreover, tinnitus is a multifactorial condition influenced by oxidative stress, vascular compromise, and neurotransmitter imbalance, suggesting that B12 deficiency may be a contributing factor rather than a direct cause. Therefore, early detection and correction of deficiency, combined with multimodal management strategies, may be more effective than isolated supplementation [27-29].

A recent systematic review by Akyay et al. (2021) found that low serum vitamin B12 levels were associated with a higher prevalence of hearing impairment, particularly at frequencies 0.25-4 KHz [29]. Additionally, a 2023 case-control study observed that deficiencies in B12 and folate were correlated with worse outcomes among patients with sudden sensorineural hearing loss (SSNHL) [30]. These results imply that adequate B vitamins may support cochlear and vestibular function via neural transmission and antioxidative pathways, though larger randomized trials are needed to confirm causality.

Recent clinical evidence has strengthened the relationship between vitamin B deficiency and auditory dysfunction. In a 2023 case-control study, Kose Celebi et al. found that patients with SSNHL had significantly lower serum vitamin B12 and folate levels than healthy controls, suggesting that deficiencies in these vitamins may contribute to cochlear damage through impaired methylation and oxidative stress mechanisms [31]. Additionally, Abu-Zaid et al. (2024) reported that combination therapy including vitamin B2, magnesium, and coenzyme Q10 improved clinical outcomes in patients with vestibular migraine, indicating a potential neurometabolic role of B-vitamins in vestibular disorders [32]. These findings align with prior observations of B-vitamin-related neuroprotection and reinforce the need for further controlled human studies.

Vitamin C

Vitamin C, also known as ascorbic acid, is a water-soluble micronutrient and a potent antioxidant [33]. It is found in many food sources, such as fruits and vegetables, and is effective in protecting cells from oxidative stress and maintaining tissue health [34]. It also plays an important role in boosting immunity and supporting

Table 1. Summary of studies on the effect of vitamins on the auditory and vestibular system

No.	Source	Year	Vitamin	Study Type	Sample Size	Human/Animal	Age Range	Key Outcome	Limitations
1	Curhan et al. AJCN [46]	2015	A, C, E, Folate	Cohort	65,000	Human	NR	Large cohort; higher carotenoid and folate intake linked to reduced hearing loss risk.	Self-reported hearing; dietary recall bias; observational design limits causal inference.
2	Schmitz et al. BMJ [50]	2012	Vitamin A	RCT Follow-up	2,300	Human	Children → Teens	Childhood vitamin A supplementation linked to lower adolescent hearing loss.	Hearing not primary endpoint; follow-up loss; no noise exposure control.
3	Aydogan et al. [44]	2013	Vitamin A & E	Case-control	80	Human	Children	Lower serum A/E levels associated with otitis media with effusion.	Small sample; cross-sectional; vitamin levels fluctuate; no diet adjustment.
4	Singh et al. Noise Health [28]	2016	Vitamin B12	Pilot Clinical Trial	40	Human	Adults	B12 therapy improved tinnitus severity.	No placebo group; very small sample; limited generalizability.
5	Karli et al. [23]	2013	Vitamin B12	Case-control	40	Human	Adults	B12 deficiency associated with abnormal OAE.	Limited sample; no control for noise exposure or comorbidities.
6	Rodrigues et al. ENT J [29]	2025	Vitamin B12	Systematic Review	40+ studies	Mixed	Varied	Strong evidence linking B12 deficiency to hearing loss.	Heterogeneous studies; limited RCT evidence; potential confounding.
7	Gopinath et al. J Nutr Health Aging [47]	2011	Vitamin C & E	Cohort	2000+	Human	Older adults	Antioxidant intake associated with lower prevalence of ARHL.	No effect on incidence; dietary recall issues; lifestyle confounders.
8	McFadden et al. Hear Res [39]	2005	Vitamin C	Animal Experiment	Guinea pigs	Animal	NR	Vitamin C reduced NIHL.	Animal model; limited human translation; high-dose exposure.
9	Cao et al. ACS Chem Neurosci [41]	2024	Vitamin C	Animal tinnitus model	Rats	Animal	NR	Vitamin C increased GLT1 expression and improved tinnitus markers.	Mechanistic; animal-only; not clinically validated.
10	Ghazavi et al. Am J Otolaryngol [30]	2020	Vitamin D	Case-control	70	Human	Adults	Vitamin D deficiency was more common in SSNHL patients.	Observational; cannot show causality; vitamin D influenced by lifestyle.
11	Shen et al. Head Face Med [61]	2025	Vitamin D	RCT	102	Human	Adults	Vitamin D supplementation improved hearing recovery in deficient SSNHL patients.	Only deficient subjects included; limited generalizability.
12	Zhao et al. PLOS ONE [62]	2025	Vitamin D	Mendelian Randomization	2 cohorts	Human	Adults	Genetic evidence supports a causal role of low vitamin D in SSNHL.	MR assumptions; limited ethnic diversity.

No.	Source	Year	Vitamin	Study Type	Sample Size	Human/Animal	Age Range	Key Outcome	Limitations
13	Tokgöz et al. [40]	2012	E, B, C	Animal Ototoxicity Study	Rats	Animal	NR	Vitamin combinations reduced cisplatin ototoxicity.	Animal doses not comparable to humans; limited translation.
14	Libonati et al. Audiology Res [60]	2022	Vitamin D + antioxidants	Clinical Cohort	150	Human	Adults	Supplementation reduced recurrence of BPPV.	Non-randomized; cannot rule out placebo effect; diet uncontrolled.
15	Ahsan et al. [64]	2014	Vitamin E (Tocotrienols)	Review	-	Mixed	Varied	Tocotrienols have antioxidant and neuroprotective potential.	Narrative review; lacks direct hearing studies.
16	Soutif-Veillon et al., Maturitas [68]	2016	Vitamin K	Observational	600+	Human	Older adults	Higher vitamin K intake associated with better cognitive performance.	Indirect relation to hearing; confounders; cognitive—not auditory—end-point.

JMR

Abbreviations: SSNHL: Sudden sensorineural hearing loss; RCTs: Randomized controlled trials; NIHL: Noise-induced hearing loss; GLT-1: Glutamate transporter 1; ARHL: Age-related hearing loss.

various cellular functions in the immune system [34]. Epidemiological studies have shown an association between high vitamin C intake and improved hearing in the mid-frequency range [36]. This vitamin acts as an antioxidant in the treatment of SSNHL and may improve treatment outcomes [6].

Evidence from animal models suggests that vitamin C may have a protective effect on hearing [37]. This vitamin improves hearing function by inhibiting NLRP3 (NOD-like receptor family, pyrin domain-containing 3) inflammasome activity and reducing cytosolic mitochondrial DNA levels in the auditory pathway [38]. This vitamin reduces oxidative stress in hair cells and may prevent noise-induced damage. Combining vitamin C with other micronutrients, such as B vitamins and magnesium, may help protect hearing and reduce damage caused by noise exposure [35, 39].

McFadden et al. investigated the protective effect of vitamin C supplementation in preventing NIHL noise-induced hearing loss in guinea pigs. In this study, animals were fed three diets: a normal diet, a vitamin C-deficient diet, and a vitamin C-supplemented diet. The results showed that after noise exposure, the hearing threshold in the group receiving the supplement was approximately 15 dB lower than in the control group. In addition, vitamin C supplementation significantly reduced the change in the permanent hearing threshold, while its deficiency did not affect HL hearing loss [40]. In another study in

rats, the preventive effects of vitamins B1, B2, B6, E, C, and L-carnitine on cisplatin-induced auditory toxicity were evaluated. ABR thresholds and DPOAE distortion-induced ear acoustic emissions were measured 72 hours after drug administration, and the results showed that these compounds, including vitamin C, reduced or prevented cisplatin-induced auditory toxicity [41].

Emerging preclinical research continues to highlight the antioxidative and neuroprotective potential of vitamin Cao et al. (2024) demonstrated that vitamin C mitigates excitotoxicity in the auditory cortex of noise-induced tinnitus models by upregulating glutamate transporter-1 (GLT-1), thereby enhancing synaptic regulation and reducing neural hyperactivity [3]. This mechanistic insight complements prior reports on vitamin C's role in preserving cochlear hair-cell function under oxidative stress and suggests a promising therapeutic target for tinnitus and noise-related hearing loss [42].

Fat-soluble vitamins

Vitamin A

Vitamin A is a fat-soluble vitamin that is considered an essential nutrient for the human body. It exists in two main forms. Retinol is the active form of vitamin A and is found mainly in animal sources; carotenoids, the precursors of vitamin A, are found mainly in plant sources. Retinol can be used directly by the body and is the main source of vi-




Figure 1. The flow diagram

JMR

tamin A in the human diet. However, carotenoids, such as beta-carotene, are converted to retinol in the body [42, 43]. Beta-carotene prevents damage caused by oxidative stress and plays a role in improving immune function [1, 2].

Vitamin A plays a role in the production of inner ear fluid and the proper functioning of hair cells [42, 43]. In humans, vitamin A and its precursors, as potent antioxidants, can help reduce lipid oxidation and neutralize free radicals, and play a crucial role in protecting hair cells and repairing damaged cells in the inner ear [44, 45]. Some studies have shown that beta-cryptoxanthin

and beta-carotene are associated with a reduced risk of hearing loss, and that vitamin A deficiency can increase the prevalence of hearing loss [46, 47]. This vitamin is also used as an antioxidant in the treatment of SSNHL and may help improve treatment outcomes [45].

In addition, retinoic acid (the metabolically active form of vitamin A) is required for normal development of the inner ear. Its deficiency can lead to dose-dependent abnormalities [48]. During pregnancy, an adequate supply of this vitamin also plays a key role in preventing SNHL [49].




Figure 2. Schematic picture of relationship between vitamins and auditory and vestibular system

In contrast, a synergy between vitamin A deficiency and infections is known, and the association of vitamin A deficiency with an increased risk of otitis media has been reported; in other words, vitamin A supplementation can reduce the risk of HL caused by otitis media [50].

Vitamin D

Vitamin D is a fat-soluble vitamin that exists in two main forms, D2 and D3 [51]. It helps absorb calcium and phosphorus and is effective in maintaining bone health, strengthening the immune system, and reducing the risk of chronic diseases [52]. Vitamin D deficiency can lead to serious problems such as muscle weakness, bone pain, and an increased risk of bone diseases and infections. Adequate intake of this vitamin from sunlight and a varied diet is very important for maintaining health [53, 54].

This vitamin plays an important role in hearing function and the balance system [55]. Vitamin D deficiency can impair bone metabolism and calcium absorption, which can in turn affect hearing function. Several studies have suggested that plasma vitamin D levels may play a critical role in the development of acute otitis media and other hearing problems [26, 56, 57]. Studies have shown an association between low plasma vitamin D levels and chronic otitis media compared to controls [44, 57]. A systematic review provided evidence that deficiencies in certain nutrients, including vitamin D, are associated with middle ear diseases and infections, particularly chronic suppurative otitis media, although direct human evidence of this association remains limited [22].

Low vitamin D levels are associated with an increased risk of age-related hearing loss (ARHL) in men over 65 years [58]. According to Feng Chen et al., different types of vitamin D have different effects on hearing. High blood levels of vitamin D2 are associated with hearing problems at low and high frequencies. Certain levels of vitamin D3 are associated with a reduced risk of high-frequency hearing loss [52].

A high prevalence of vitamin D deficiency has been reported in patients with diseases of the inner ear, including benign paroxysmal positional vertigo, Ménière's disease, vestibular neuritis, idiopathic facial palsy, and acute idiopathic hearing loss and tinnitus [34, 38]. Vitamin D plays a role in maintaining the structure of the otoconia by regulating calcium levels within the semicircular canals. Therefore, low vitamin D levels can impair calcium metabolism and transport, which can, in turn, impair otoconia absorption and synthesis [59]. Finally, providing adequate vitamin D levels through diet or

supplementation may be a therapeutic approach to reduce the risk of BPPV and hearing problems [60]. In a randomized controlled trial by Shen et al. (2025), combining vitamin D supplementation with standard therapy in patients with SSNHL and vitamin D deficiency led to significantly greater improvements in hearing recovery and tinnitus relief than therapy alone [61]. Complementing this, a Mendelian randomization study by Zhao et al. (2025) found no evidence of a causal link between genetically predicted serum 25-hydroxyvitamin D levels and idiopathic sudden sensorineural hearing loss [62].

Vitamin E

Vitamin E is a fat-soluble vitamin that acts as a potent antioxidant and plays important roles in maintaining body health [63]. This vitamin is divided into two main groups: tocopherols and tocotrienols, both of which have antioxidant and anti-inflammatory properties [64]. This vitamin can neutralize free radicals and prevent glutamate-induced neurotoxicity. Vitamin E helps maintain healthy skin, strengthen the immune system, and reduce the risk of chronic diseases. Adequate intake of this vitamin through a varied diet is important for maintaining general health [65]. It is found in food sources such as vegetable oils, nuts, seeds, and leafy green vegetables [64]. Vitamin E is used as an antioxidant in the treatment of sudden-onset hearing loss and may help improve treatment outcomes [45]. In combination with other vitamins and magnesium, vitamin E may help protect hearing and reduce damage from noise exposure [38]. In patients with BPPV, vitamin E supplementation may improve function [65].

Vitamin K

There is no direct, strong, and well-established evidence demonstrating that vitamin K has a specific and reliable effect on hearing, whether in preventing or treating hearing loss. Studies examining vitamin K as an independent and specific factor related to hearing are very limited or incomplete. Regarding vitamin K itself, there are basic animal studies and some research on the effects of vitamin K₂ on the brain and aging; however, these do not provide direct evidence of any impact on the cochlea or of improved auditory thresholds in humans. Therefore, clinical conclusions cannot be reliably drawn from animal data [66, 67]. Some smaller, non-review papers—such as surveys or nutritional reports—have mentioned vitamin K intake in certain groups, but these do not provide causal links or robust clinical evidence [68, 69]. Table 1 summarizes studies on the effects of vitamins on the auditory and vestibular systems.

Discussion

The findings indicate that adequate intake of water- and fat-soluble vitamins may contribute to the prevention and adjunctive management of auditory and vestibular disorders. Key applications include: Screening elderly or at-risk populations for B12 and folate deficiencies to prevent SNHL and vestibular dysfunction, considering vitamin C supplementation in noise-exposed individuals or those receiving ototoxic drugs like cisplatin, utilizing vitamin D supplementation as an adjunct in SSNHL therapy and for patients with BPPV or other vestibular disorders and combined supplementation of vitamins B, C, and E may enhance neuroprotection and auditory recovery in clinical settings.

Despite strong preclinical evidence, several gaps remain: A, randomized controlled trials (RCTs) with standardized dosing and long-term follow-up are needed to confirm the efficacy of vitamin supplementation in hearing and balance disorders. b, investigations should assess the synergistic effects of combined vitamins on cochlear and vestibular protection, particularly in elderly populations or patients with chronic conditions. c, mechanistic studies using advanced imaging and electrophysiological measures (e.g. TEOAE, DPOAE, VEMP) are required to delineate the pathways by which vitamins influence auditory and vestibular function. d, Studies should explore personalized supplementation strategies based on genetic, dietary, and environmental factors to maximize clinical benefit. e, Longitudinal cohort studies are necessary to determine whether early correction of vitamin deficiencies can prevent ARHLs and vestibular decline.

Limitation

This review is narrative and based on heterogeneous studies varying in design, sample size, and methodology. Many of the included studies were observational or animal-based, limiting causal inference. Publication bias and the exclusion of non-English literature may also affect the completeness of our review.

Conclusion

Current evidence suggests that various vitamins and nutrients may influence auditory and vestibular health. Deficiencies in vitamins A, B, C, D, and E have been associated with hearing problems and balance disorders. In particular, vitamins A and E are suggested to support hair cell integrity and protect against oxidative stress, while vitamins B and C are linked to improved neural function and a potentially reduced risk of hearing loss. Ensuring adequate vitamin intake through diet or supplementation may be a supportive measure for maintaining hearing and balance.

Furthermore, deficiencies in vitamins B2, B3, and B6 can contribute to neurological symptoms, dizziness, and tinnitus, and vitamin D may influence vestibular function through its role in bone metabolism and calcium homeostasis. Vitamin E, as a potent antioxidant, may also help mitigate oxidative damage in inner ear tissues. Considering the relatively high prevalence of vitamin deficiencies among individuals with vestibular disorders, maintaining sufficient vitamin levels could represent a valuable adjunctive strategy for preserving auditory and vestibular function and improving quality of life.

Ethical Considerations

Compliance with ethical guidelines

There were no ethical considerations to be considered in this research.

Funding

This study did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Authors' contributions

Visualization: Aylin Mellati; Conceptualization, formal analysis, supervision, review and editing: Nasrin Gohari and Aylin Mellati; Data curation: Nasrin Gohari; Project administration: Nasrin Gohari, Nayiere Mansouri; Investigation, methodology, validation, writing the original draft, final approval: All authors.

Conflict of interest

The authors declared no conflict of interest.

References

- [1] Shastak Y, Gordillo A, Pelletier W. The relationship between vitamin A status and oxidative stress in animal production. *Journal of Applied Animal Research*. 2023; 51(1):546-53. [DOI:10.1080/09712119.2023.2239319]
- [2] Tufail T, Bader UI Ain H, Noreen S, Ikram A, Arshad MT, Abdullahi MA. Nutritional benefits of lycopene and beta-carotene: A comprehensive overview. *Food Science & Nutrition*. 2024; 12(11):8715-41. [DOI:10.1002/fsn3.4502] [PMID] [PMCID]
- [3] Jung SY, Kim SH, Yeo SG. Association of nutritional factors with hearing loss. *Nutrients*. 2019; 11(2):307. [DOI:10.3390/nu11020307] [PMID] [PMCID]

- [4] Martínez-Vega R, Garrido F, Partearroyo T, Cediel R, Zeisel SH, Martínez-Alvarez C, et al. Folic acid deficiency induces premature hearing loss through mechanisms involving cochlear oxidative stress and impairment of homocysteine metabolism. *The FASEB Journal*. 2014; 29(2):418. [DOI:10.1096/fj.14-259283] [PMID] [PMCID]
- [5] Mohan A, Kumar R, Kumar V, Yadav M. Homocysteine, vitamin B12 and folate level: Possible risk factors in the progression of chronic heart and kidney disorders. *Current Cardiology Reviews*. 2023; 19(4):66-83. [DOI:10.2174/1573403X1966623020911854] [PMID] [PMCID]
- [6] Mehrkian S, Moossavi A, Gohari N, Nazari MA, Bakhshi E, Alain C. Long latency auditory evoked potentials and object-related negativity based on harmonicity in hearing-impaired children. *Neuroscience Research*. 2022; 178:52-9. [DOI:10.1016/j.neures.2022.01.001] [PMID]
- [7] Fakhri J, Rouhbakhsh N, Hoseinabadi R, Fatahi F, Sephernejad M, Roghani Z. Investigating quick speech-in-noise comprehension in adult bimodal users. *Journal of Modern Rehabilitation*. 2021; 15(3):161-6. [DOI:10.18502/jmr.v15i3.7736]
- [8] Sayadi N, Saghar A, Nasrin G. [Determining the quality of life in the elderly with high frequency hearing loss before and after hearing aid fitting (Persian)]. *Auditory and Vestibular Research*. 2018; 27(2):111-5. [Link]
- [9] Nasrin G, Ebtesam S, Zoleikha A, Bahare K. A comparative study on the general health of the mothers of children with cochlear implant, hearing aid, and normal hearing. *Auditory and Vestibular Research*. 2020; 29(2):117-21. [Link]
- [10] Abbasi M, Pourrajab B, Tokhi MO. Protective effects of vitamins/antioxidants on occupational noise-induced hearing loss: A systematic review. *Journal of Occupational Health*. 2021; 63(1):e12217. [DOI:10.1002/1348-9585.12217] [PMID] [PMCID]
- [11] Schellack G, Harirari P, Schellack N. Vitamin B-complex deficiency, supplementation and management. *SA Pharmaceutical Journal*. 2019; 86(3):23-9. [Link]
- [12] Hanna M, Jaqua E, Nguyen V, Clay J. B Vitamins: Functions and uses in medicine. *The Permanente Journal*. 2022; 26(2):89-97. [DOI:10.7812/TPP/21.204] [PMID] [PMCID]
- [13] Mrowicka M, Mrowicki J, Dragan G, Majsterek I. The importance of thiamine (vitamin B1) in humans. *Bioscience Reports*. 2023; 43(10):BSR20230374. [DOI:10.1042/BSR20230374] [PMID] [PMCID]
- [14] Pinto J, Rivlin R. Riboflavin (vitamin B2). In: Zemleni J, Suttie JW, Gregory JF, Stover PJ, editors. *Handbook of vitamins*. Boca Raton: CRC Press; 2013. [Link]
- [15] Sarwar MF, Sarwar MH. Deficiency of vitamin B-complex and its relation with body disorders. In: Guy LeBlanc J, editor. *B-complex vitamins: Sources, intakes and novel applications*. London: IntechOpen; 2022. [Link]
- [16] Rucker RB, Bauerly K. Pantothenic acid. In: Zemleni J, Suttie JW, Gregory III JF, Stover PJ, editors. *Handbook of vitamins*. Boca Raton: CRC Press; 2013. [Link]
- [17] Parra M, Stahl S, Hellmann H. Vitamin B6 and its role in cell metabolism and physiology. *Cells*. 2018; 7(7):84. [DOI:10.3390/cells7070084] [PMID] [PMCID]
- [18] Tardy A-L, Pouteau E, Marquez D, Yilmaz C, Scholey A. Vitamins and minerals for energy, fatigue and cognition: A narrative review of the biochemical and clinical evidence. *Nutrients*. 2020; 12(1):228. [DOI:10.3390/nu12010228] [PMID] [PMCID]
- [19] Von Fraunhofer J. Vitamin B9 and Health. *Ecronicon Dental Science*. 2019; 18(6):1331-5. [Link]
- [20] Fava M, Mischoulon D. Folate in depression: efficacy, safety, differences in formulations, and clinical issues. *Journal of Clinical Psychiatry*. 2009; 70(5):12-7. [DOI:10.4088/JCP.8157su1c.03] [PMID]
- [21] Maqbool MA, Aslam M, Akbar W, Iqbal Z. Biological importance of vitamins for human health: A review. *The Journal of Agricultural Science*. 2017; 2(3):50-8. [Link]
- [22] Chiarella G, Marcianò G, Viola P, Palleria C, Pisani D, Rania V, et al. Nutraceuticals for peripheral vestibular pathology: properties, usefulness, future perspectives and medico-legal aspects. *Nutrients*. 2021; 13(10):3646. [DOI:10.3390/nu13103646] [PMID] [PMCID]
- [23] Karli R, Gül A, Uğur B. Effect of vitamin B12 deficiency on otoacoustic emissions. *Acta Otorhinolaryngologica Italica*. 2013;33(4):243. [PMID]
- [24] Berkiten G, Yildirim G, Topaloglu I, Ugras H. Vitamin B12 levels in patients with tinnitus and effectiveness of vitamin B12 treatment on hearing threshold and tinnitus. *B-ENT*. 2013; 9(2):111-6. [Link]
- [25] Akyay A, Soyulu E, Ünsal S, Demiroğlu H, Bahçeci S. Hearing status in vitamin B12-deficient children. *Journal of Paediatrics and Child Health*. 2021; 57(7):1060-6. [DOI:10.1111/jpc.15392] [PMID]
- [26] Özdemir D, Mehel DM, Küçüköner Ö, Ağrı İ, Yemiş T, Akgöl G, et al. Vestibular evoked myogenic potentials in patients with low vitamin B12 levels. *Ear, Nose & Throat Journal*. 2021; 100(5):NP231-5. [DOI:10.1177/0145561319878952] [PMID]
- [27] Mathew AR, Di Matteo G, La Rosa P, Barbati SA, Mannina L, Moreno S, et al. Vitamin B12 deficiency and the nervous system: beyond metabolic decompensation-comparing biological models and gaining new insights into molecular and cellular mechanisms. *International Journal of Molecular Sciences*. 2024; 25(1):590. [DOI:10.3390/ijms25010590] [PMID] [PMCID]
- [28] Singh C, Kawatra R, Gupta J, Awasthi V, Dungana H. Therapeutic role of Vitamin B12 in patients of chronic tinnitus: A pilot study. *Noise Health*. 2016; 18(81):93-7. [DOI:10.4103/1463-1741.178485] [PMID] [PMCID]
- [29] Rodrigues J, Anand S, Gunjawate DR, Kumar K, Ravi R. Exploring the intricate connection between vitamin B12 deficiency and hearing loss: A systematic literature review. *Ear, Nose & Throat Journal*. 2025; 01455613241298070. [DOI:10.1177/01455613241298070] [PMID]
- [30] Ghazavi H, Kargoshaie A-A, Jamshidi-Koohsari M. Investigation of vitamin D levels in patients with sudden sensory-neural hearing loss and its effect on treatment. *American Journal of Otolaryngology*. 2020; 41(2):102327. [DOI:10.1016/j.amjoto.2019.102327] [PMID]

- [31] Kose Celebi N, Deveci HS, Kulekci Ozturk S, Aslan Dundar T. Clinical role of vitamin D, vitamin B12, folate levels and hematological parameters in patients with sudden sensorineural hearing loss. *Acta Oto-Laryngologica*. 2023; 143(7):596-601. [DOI:10.1080/00016489.2023.2235398] [PMID]
- [32] Abu-Zaid A, Abu-Zaid S, Barakat M, Al-Huniti R, Khair H. Effectiveness of combination therapy of magnesium, vitamin B2 and Co-enzyme 10 supplementation on vestibular migraine: a retrospective cohort study. *Pharmacia*. 2024; 71(1):1-7. [DOI:10.3897/pharmacia.71.e112909]
- [33] Carr AC, Maggini S. Vitamin C and immune function. *Nutrients*. 2017; 9(11):1211. [DOI:10.3390/nu9111211] [PMID] [PMCID]
- [34] Traber MG, Stevens JF. Vitamins C and E: Beneficial effects from a mechanistic perspective. *Free Radical Biology and Medicine*. 2011; 51(5):1000-13. [DOI:10.1016/j.freeradbiomed.2011.05.017] [PMID] [PMCID]
- [35] Kang JW, Choi HS, Kim K, Choi JY. Dietary vitamin intake correlates with hearing thresholds in the older population: the Korean national health and nutrition examination survey. *The American Journal of Clinical Nutrition*. 2014; 99(6):1407-13. [DOI:10.3945/ajcn.113.072793] [PMID]
- [36] Choi YH, Miller JM, Tucker KL, Hu H, Park SK. Antioxidant vitamins and magnesium and the risk of hearing loss in the US general population. *The American Journal of Clinical Nutrition*. 2014; 99(1):148-55. [DOI:10.3945/ajcn.113.068437] [PMID] [PMCID]
- [37] Li H, Yang X, Song Y, Zhu Q, Liao Z, Liang Y, et al. PRRSV infection activates NLRP3 inflammasome through inducing cytosolic mitochondrial DNA stress. *Veterinary Microbiology*. 2023; 279:109673. [DOI:10.1016/j.vetmic.2023.109673] [PMID]
- [38] Le Prell CG, Gagnon PM, Bennett DC, Ohlemiller KK. Nutrient-enhanced diet reduces noise-induced damage to the inner ear and hearing loss. *Translational Research*. 2011; 158(1):38-53. [DOI:10.1016/j.trsl.2011.02.006] [PMID] [PMCID]
- [39] McFadden SL, Woo JM, Michalak N, Ding D. Dietary vitamin C supplementation reduces noise-induced hearing loss in guinea pigs. *Hearing Research*. 2005; 202(1-2):200-8. [DOI:10.1016/j.heares.2004.10.011] [PMID]
- [40] Tokgöz SA, Vuralkan E, Sonbay ND, Çalışkan M, Saka C, Beşaltı Ö, et al. Protective effects of vitamins E, B and C and L-carnitine in the prevention of cisplatin-induced ototoxicity in rats. *The Journal of Laryngology & Otology*. 2012; 126(5):464-9. [DOI:10.1017/S0022215112000382] [PMID]
- [41] Cao W, Xiong S, Ji W, Wei H, Ma F, Mao L. Neuroprotection role of vitamin C by upregulating glutamate transporter-1 in auditory cortex of noise-induced tinnitus animal model. *ACS Chemical Neuroscience*. 2024; 15(6):1197-205. [DOI:10.1021/acscchemneuro.3c00765] [PMID]
- [42] Carazo A, Macáková K, Matoušová K, Krčmová LK, Protti M, Mladěnka P. Vitamin A update: forms, sources, kinetics, detection, function, deficiency, therapeutic use and toxicity. *Nutrients*. 2021; 13(5):1703. [DOI:10.3390/nu13051703] [PMID] [PMCID]
- [43] Aydoğan F, Aydın E, Taştan E, Arslan N, Senes M, Unlu I, et al. Is there a relationship between serum levels of vitamin A, vitamin E, copper and zinc and otitis media with effusion in children? *Indian Journal of Otolaryngology and Head & Neck Surgery*. 2013; 65(Suppl 3):594-7. [DOI:10.1007/s12070-012-0564-z] [PMID] [PMCID]
- [44] Rodrigo L, Campos-Asensio C, Rodríguez MÁ, Crespo I, Olmedillas H. Role of nutrition in the development and prevention of age-related hearing loss: A scoping review. *Journal of the Formosan Medical Association*. 2021; 120(1):107-20. [DOI:10.1016/j.jfma.2020.05.011] [PMID]
- [45] Ibrahim I, Zeitouni A, da Silva SD. Effect of antioxidant vitamins as adjuvant therapy for sudden sensorineural hearing loss: Systematic review study. *Audiology and Neurotology*. 2018; 23(1):1-7. [DOI:10.1159/000486274] [PMID]
- [46] Curhan SG, Stankovic KM, Eavey RD, Wang M, Stampfer MJ, Curhan GC. Carotenoids, vitamin A, vitamin C, vitamin E, and folate and risk of self-reported hearing loss in women. *The American Journal of Clinical Nutrition*. 2015; 102(5):1167-75. [DOI:10.3945/ajcn.115.109314] [PMID] [PMCID]
- [47] Gopinath B, Flood V, McMahon C, Burlutsky G, Spankovich C, Hood LJ, et al. Dietary antioxidant intake is associated with the prevalence but not incidence of age-related hearing loss. *The Journal of Nutrition, Health & Aging*. 2011; 15(10):896-900. [DOI:10.1007/s12603-011-0119-0] [PMID] [PMCID]
- [48] Gao L, Tona Y, Omori K. Spatiotemporal expression of CWC27 spliceosome-associated protein (CWC27) in the developing mouse inner ear. *Gene Expr Patterns*. 2025; 56:119400. [DOI:10.1016/j.gep.2025.119400] [PMID]
- [49] Emmett SD, West KP, Jr. Gestational vitamin A deficiency: a novel cause of sensorineural hearing loss in the developing world? *Medical Hypotheses*. 2014; 82(1):6-10. [DOI:10.1016/j.mehy.2013.09.028] [PMID] [PMCID]
- [50] Schmitz J, West KP, Jr., Khatry SK, Wu L, Leclercq SC, Karna SL, et al. Vitamin A supplementation in preschool children and risk of hearing loss as adolescents and young adults in rural Nepal: randomised trial cohort follow-up study. *BMJ*. 2012; 344:d7962. [DOI:10.1136/bmj.d7962] [PMID] [PMCID]
- [51] Ravisankar P, Reddy AA, Nagalakshmi B, Koushik OS, Kumar BV, Anvith PS. The comprehensive review on fat soluble vitamins. *IOSR Journal of Pharmacy*. 2015; 5(11):12-28. [Link]
- [52] Wang H, Chen W, Li D, Yin X, Zhang X, Olsen N, et al. Vitamin D and chronic diseases. *Aging and Disease*. 2017; 8(3):346. [DOI:10.14336/AD.2016.1021] [PMID] [PMCID]
- [53] Kupisz-Urbańska M, Pludowski P, Marciniowska-Suchowierska E. Vitamin D deficiency in older patients-problems of sarcopenia, drug interactions, management in deficiency. *Nutrients*. 2021; 13(4):1247. [DOI:10.3390/nu13041247] [PMID] [PMCID]
- [54] Galesanu C, Mocanu V. Vitamin D deficiency and the clinical consequences. *The Medical-Surgical Journal*. 2015; 119(2):310-8. [Link]
- [55] Aiello G, Lombardo M, Baldelli S. Exploring vitamin D synthesis and function in cardiovascular health: A narrative review. *Applied Sciences*. 2024; 14(11):4339. [DOI:10.3390/app14114339]

- [56] Taneja MK, Taneja V. Role of vitamin D in prevention of deafness. *Indian Journal of Otolaryngology*. 18(2):55-57. [DOI:10.4103/0971-7749.100692]
- [57] Li HB, Tai XH, Sang YH, Jia JP, Xu ZM, Cui XF, et al. Association between vitamin D and development of otitis media: A PRISMA-compliant meta-analysis and systematic review. *Medicine*. 2016; 95(40):e4739. [DOI:10.1097/MD.0000000000004739] [PMID] [PMCID]
- [58] Lee YJ, Jung JH, Chung JW. The relationship between lower vitamin D levels and hearing loss in older Adults. *Journal of Audiology & Otolaryngology*. 2024; 28(4):252. [DOI:10.7874/jao.2023.00458] [PMID] [PMCID]
- [59] Hamayal M, Khurshid S, Zahid MA, Khurshid N, Shahid W, Ali M, et al. Exploring the significance of vitamin D levels as a biomarker in ear diseases: A narrative review. *Cureus*. 2024;16(2):e54812. [DOI:10.7759/cureus.54812] [PMID] [PMCID]
- [60] Libonati GA, Leone A, Martellucci S, Gallo A, Albera R, Lucisano S, et al. Prevention of recurrent benign paroxysmal positional vertigo: The role of combined supplementation with vitamin D and antioxidants. *Audiology Research*. 2022; 12(4):445-56. [DOI:10.3390/audiolres12040045] [PMID] [PMCID]
- [61] Shen X, Yang M, Tian J, Xie L, Feng N, Ma R. Clinical efficacy of vitamin D combined with conventional therapy for sudden sensorineural hearing loss in patients with vitamin D deficiency: A randomized controlled trial. *Head & Face Medicine*. 2025; 21(1):68. [DOI:10.1186/s13005-025-00545-2] [PMID] [PMCID]
- [62] Zhao Y, Yu C, Sun H, Xie F, Shen J, Li X, et al. Exploring the causal link between serum 25-hydroxyvitamin D concentrations and idiopathic sudden sensorineural hearing loss: Insights gained from a Mendelian randomization study involving two independent samples. *PLoS One*. 2025; 20(5):e0322898. [DOI:10.1371/journal.pone.0322898] [PMID] [PMCID]
- [63] Rafeeq H, Ahmad S, Tareen MBK, Shahzad KA, Bashir A, Jabeen R, et al. Biochemistry of fat soluble vitamins, sources, biochemical functions and toxicity. *Haya: The Saudi Journal of Life Sciences*. 2020; 5(6):188-96. [DOI:10.36348/sjls.2020.v05i09.007]
- [64] Ahsan H, Ahad A, Iqbal J, Siddiqui WA. Pharmacological potential of tocotrienols: A review. *Nutrition & Metabolism*. 2014; 11:1-22. [DOI:10.1186/1743-7075-11-52] [PMID] [PMCID]
- [65] Dattola A, Silvestri M, Bennardo L, Passante M, Scali E, Patruno C, et al. Role of vitamins in skin health: A systematic review. *Current Nutrition Reports*. 2020; 9(3):226-35. [DOI:10.1007/s13668-020-00322-4] [PMID]
- [66] Elkattawy HA, Ghoneim FM, Eladl MA, Said E, Ebrahim HA, El-Shafey M, et al. Vitamin K2 (Menaquinone-7) reverses age-related structural and cognitive deterioration in naturally aging rats. *Antioxidants*. 2022; 11(3):514. [DOI:10.3390/antiox11030514] [PMID] [PMCID]
- [67] Lin X, Wen X, Wei Z, Guo K, Shi F, Huang T, et al. Vitamin K2 protects against A β 42-induced neurotoxicity by activating autophagy and improving mitochondrial function in *Drosophila*. *Neuroreport*. 2021; 32(6):431-7. [DOI:10.1097/WNR.0000000000001599] [PMID] [PMCID]
- [68] Soutif-Veillon A, Ferland G, Rolland Y, Presse N, Boucher K, Féart C, et al. Increased dietary vitamin K intake is associated with less severe subjective memory complaint among older adults. *Maturitas*. 2016; 93:131-6. [DOI:10.1016/j.maturitas.2016.02.004] [PMID]
- [69] Camacho-Barcia L, García-Gavilán J, Martínez-González M, Fernández-Aranda F, Galié S, Corella D, et al. Vitamin K dietary intake is associated with cognitive function in an older adult Mediterranean population. *Age and Ageing*. 2022; 51(2):afab246. [DOI:10.1093/ageing/afab246] [PMID]