

# Research Paper: The Effect of Eight Weeks of Iyengar Yoga With an Emphasis on Spine and Shoulder Exercises on the Upper Cross Syndrome in Middle-aged Women

Shilan Sohrabi<sup>1</sup>, Mohammad Rahimi<sup>2</sup>, Mojtaba Babaei-Mobarakeh<sup>3</sup>, Hashem Piri<sup>4\*</sup>

1. Department of Physical Education & Sport Sciences, Faculty of Literature, Humanities and Social Sciences, Tehran Science and Research Branch, Islamic Azad University, Tehran, Iran.

2. Corrective Exercises and Sport Injuries, Faculty of Sport Sciences, Shahid Rajaei Teacher Training University, Tehran, Iran.

3. Department of Biomechanics and Sports Injuries, Faculty of Physical Education and Sport Sciences, Kharazmi University, Tehran, Iran.

4. Department of Corrective Exercise & Sports Injuries, Faculty of Physical Education and Sport Sciences, Allameh Tabataba'i University, Tehran, Iran.



**Citation:** Sohrabi Sh, Rahimi M, Babaei-Mobarakeh M, Piri H. The Effect of Eight Weeks of Iyengar Yoga With an Emphasis on Spine and Shoulder Exercises on the Upper Cross Syndrome in Middle-aged Women. *Journal of Modern Rehabilitation*. 2020; 14(3):159-168. <http://dx.doi.org/10.18502/jmr.v14i3.7714>

**doi** <http://dx.doi.org/10.18502/jmr.v14i3.7714>



## Article info:

**Received:** 21 Feb 2020

**Accepted:** 28 Apr 2020

**Available Online:** 01 Jul 2020

## Keywords:

Iyengar, Syndrome, Exercise

## ABSTRACT

**Introduction:** Upper Crossed Syndrome (UCS) is a combination of forward head, rounded shoulder, and hyperkyphosis deformities. Yoga is a non-competitive physical exercise with the potential to correct postural imbalances in the human body. Iyengar yoga is a form of Hatha yoga.

**Materials and Methods:** The purpose of present study was to evaluate the effect of Iyengar yoga with an emphasis on spine and shoulder exercises on the UCS in middle-aged women. In this quasi-experimental applied research, 15 subjects were purposefully recruited out of middle-aged women affected with UCS. The participants performed Iyengar yoga exercises with an emphasis on the spine and shoulder. The photogrammetry method was used to measure UCS.

**Results:** At the end of 8 weeks exercises, there was a significant increase in the cervical ( $P < 0.001$ ) and shoulder angles ( $P < 0.005$ ), and a significant decrease in thoracic flexion angle ( $P < 0.001$ ).

**Conclusion:** All relevant coaches and therapists are recommended considering Iyengar yoga as an alternative for training programs in middle-aged women affected by UCS.

## 1. Introduction

To evaluate the effect of Iyengar yoga with an emphasis on spine and shoulder exercises on the UCS in middle-aged women. Posture is defined as the appropriate arrangement of the body parts [1]. In a balanced structure, local and global muscles activate accordingly to protect the body

and spine from injuries during activities such as lifting, sitting, and standing [2]. However, a faulty relationship between various parts of the body with abnormal spinal curves may impair this balance [3]. In this regard, Vladimir Janda recognized three muscle imbalance patterns that accompany with distinctive pain syndromes: the upper crossed, pelvic crossed, and stratification syndromes [4]. Upper Crossed Syndrome (UCS) is also referred to

\* Corresponding Author:

**Hashem Piri, PhD.**

**Address:** Department of Corrective Exercise & Sports Injuries, Faculty of Physical Education and Sport Sciences, Allameh Tabataba'i University, Tehran, Iran.

**Tel:** +98 (919) 9033390

**E-mail:** [hpiri1984@gmail.com](mailto:hpiri1984@gmail.com)

as forward head and rounded shoulder posture, forward head posture and protracted shoulders, and proximal or shoulder girdle crossed syndrome [4-6]. UCS has been defined as tightness or facilitation of postural muscles, namely, the pectoralis major, pectoralis minor, subscapularis, latissimus dorsi, levator scapulae, upper trapezius, teres major, sternocleidomastoid, upper trapezius, and scalenes, and weakness or inhibition of phasic muscles, namely, the rhomboids, serratus anterior, middle and lower trapezius, teres minor, infraspinatus, and the deep neck flexors [7, 8]. Janda called this syndrome “upper crossed” because when the weakened and tightened muscles are joined in the upper body, they form a cross [9].

UCS results from muscle imbalances between two types of muscles, namely, postural or tonic and phasic muscles [10]. Predominantly postural or tonic muscles such as pectoralis major tend to become tight, whereas predominantly phasic muscles such as deep-neck flexors tend to become weak [11]. People who afflicted with UCS will show particular postural changes, including Forward Head Posture (FHP), rounded back, elevation and protraction of the shoulders, winging of the scapula, and diminished mobility of the thoracic spine [4, 8].

FHP and Rounded Shoulder Posture (RSP) as the components of UCS are prevalent in people of all ages and with different jobs [6, 12-15]. In an Iranian academic dental staff, the prevalence rates of the FHP and RSP have been reported as 85.5%, 68.8%, respectively [14]. The prevalence of UCS in laundry workers is reported as 28% [11].

Many disorders are linked with UCS, including migraine headaches; chronic headaches; Subacromial impingement; biceps tendinitis; thoracic outlet syndrome; degeneration of the cervical spine; and joint dysfunction at the C1-C2 segment, C4-C5 segment, cervicothoracic joint, and T4-T5 segment [1, 4, 8, 9, 16]. Also, dysfunction in the sternoclavicular joint, acromioclavicular joint and thoracic and cervical facet joints have been proposed as potential joint dysfunctions in individuals with UCS [8]. Altered scapular kinematics and muscle activity, modification in biomechanics and neuromuscular control of the shoulder complex and upper extremity, and reduction in glenohumeral stability are among the other consequences of UCS [1, 4, 5, 17]. UCS-related disorders greatly affect the safety and health status worldwide and it may increase in the near future [18].

Yoga is a way of living. It was first introduced by Swami Vivekananda in the United States in 1893, and now yoga practice has evolved to target health and beauty [19]. Yoga has been traditionally used to treat many pains or

malpostures like hyperkyphosis in various populations [20]. Iyengar yoga is a form of Hatha yoga with key aspects of correct alignment of the body and sequences of postures using props such as blocks, belts (rope), and chairs [21]. To the best of our knowledge, the effect of Iyengar yoga with an emphasis on spine and shoulder exercises on UCS has not been studied. Previous studies have examined the effects of general yoga exercises on hamstring flexibility, and discrete abnormalities such as hyperkyphosis [22-27]. Also, the effects of different exercises such as scapular stabilization, strengthening and stretching, and comprehensive exercises on the discrete component of UCS such as FHP, RSP, and kyphosis have been investigated [28-30].

Considering the high prevalence of UCS and its frequent complications, there is no evidence about the effectiveness of Iyengar yoga with an emphasis on spine and shoulder exercises for the management of UCS. So, regarding the safety of yoga exercises and the possibility of doing these exercises by a person alone at home, this study aimed to evaluate the effect of Iyengar yoga with an emphasis on spine and shoulder exercises on correcting UCS in middle-aged women.

## 2. Materials and Methods

In this quasi-experimental applied study, 15 participants were selected, using a convenience sampling method from middle-aged women afflicted with UCS, aged between 45 and 61 years. Considering  $\alpha=0.05$ , the power of 80%, and an effect size of 0.80, the sample size was determined as 15 subjects by using G\*Power software [31]. The inclusion criteria for the participants were having a shoulder angle less than 52°, a cervical angle less than 50°, and a thoracic flexion angle of more than 50° [32, 33]. The participants would be excluded if they had a vertebral fracture, scoliosis, severe dysplasia, rheumatic disease, torticollis, persistent respiratory problems, uncontrolled hypertension, pulmonary disease, cervical spine, and shoulder instability. They would also be excluded if they missed practice for two consecutive sessions or more. All participants were informed about the procedures and signed a consent form before the initiation of the study.

Yoga exercises are performed three times a week. The participants performed general Iyengar yoga 1 day per week (1 hour each day) (Table 1) and Iyengar yoga home-based training with an emphasis on spine and shoulder exercises 2 days (half an hour each day) per week (Table 2). They were monitored through phone calls, and an exercise checklist was completed by each participant. All exercise sessions began with the moun-

Table 1. General Iyengar yoga exercises

Number	Movement Name	Description
1	Tadasana	Mountain pose
2	Utkatasana	Chair Pose
3	Ardha Uttanasana	Standing Half Forward Bend
4	Virabhadrasana I	Warrior Pose I
5	Virabhadrasana II	Warrior Pose II
6	Vrikshasana	Tree Pose
7	Utthita trikonasana	Triangle Pose
8	Utthita parśvakonasana	Extended Side Angle Pose
9	Parsvottanasana	Pyramid Pose
10	Darnikasana	Child pose
11	Adho-Mukha Śvanasana	Downward-facing dog pose
12	Urdhva-mukha Śvanasana	Upward Facing Dog Pose
13	Parivrtta-trikonasana	Revolved Triangle Pose
14	Parivrtta Parsvakonasana	Revolved Side Angle Pose
15	Salabhasana	Locust Pose
16	Salamba bhujangasana	Sphinx
17	Dhanurasana	Bow Posture
18	Ardha urdhva Dhanurasana	Half Upper Bow Pose
19	Dandasana	Staff Pose
20	Gomukhasana	Cow Face Pose
21	Paschimottanasana	Seated Forward Bend
22	Parivrtta Paschimottanasana	Revolved Seated Forward Bend
23	Baddha Konasana	Bound Angle Pose
24	Upavistha Konasana	Wide-Angle Seated Forward Bend
25	Marichyasana	Sage's Pose
26	Urdhva Prasarita Padasana	Upward Extended Feet Pose
27	Anantasana	Sleeping Vishnu Pose
28	Setu Bandha Sarvangasana	Bridge Pose
29	Sarvangasana	Shoulder Stand
30	Viparita Karani	Legs Up the Wall Pose

JMR

tain posture (Tadasana), followed by positions focused primarily on elongating the spine and improving flexibility, balance, muscle strength, and endurance [22]. Yoga exercises were done for 8 weeks.

The intended angles of the body were obtained by commercial cameras (16.2-MP Samsung Digital Camera, WB30F, made in South Korea) using optimal resolution without zooming [34]. The camera lens was perpendicular to the floor and was placed on a tripod parallel to the participants. The camera was set at 50% of the participant's height from the ground. To normalize the angle of the camera lens and increase the measurement accuracy,

each participant's height was used to calibrate the images. The body of the participant was aligned with the plumb line [34]. The privacy and comfort of the participants were guaranteed [35].

To capture the participant's natural alignment, each person was asked to bent forward 3 times and place hands on head 3 times, then look naturally and calmly to the imaginary point on the opposite wall [28]. Then the experimenter takes a photo after 5 seconds pause [28]. Finally, the photos were transferred to the computer and processed using the Kinovea program [36]. This program has been used in previous studies, and it is both

**Table 2.** The Iyengar yoga home-based exercises with emphasis on spine and shoulder

Number	Movement Name	Description
1	Tadasana	Mountain pose
2	Bhujangasana	Cobra pose
3	Adho-Mukha Śvanasana	Downward-facing dog pose
4	Ustrasana	Camel pose
5	Ardha Uttanasana	Standing Half forward bend
6	Gomukhasana	Cow Face Pose
7	Vajrasana	Diamond Pose
8	Viparita Dandasana	Inverted Staff Pose
9	Salabhasana	Locust pose

**JMR**

valid and reliable [36]. The following measurements were used to assess UCS.

### Cervical angle

The cervical angle is a reliable angle to measure the FHP [32]. The angle between the horizontal line passing through the spinal process of C7 and the line that attach the central point of the tragus of the ear to the spinal process of C7 considered as the cervical angle [32] (Figure 1). A skin marker was placed on the spinal process of C7. If the cervical angle was less than  $50^\circ$ , the subject would be considered to have FHP [32].


### Shoulder angle

To assess shoulder posture, the angle between the horizontal line passing through the spinal process of the C7, and a line drawn from the midpoint of the shoulder joint to the spinal process of C7 was measured [37] (Figure 1). A skin marker was placed on the spinal process of C7. If the shoulder angle less than  $52^\circ$ , the subject would be considered to have forward shoulder posture [32].

### Thoracic flexion angle

The thoracic kyphosis was determined by measuring the thoracic flexion angle [38]. Skin markers were placed on the spinous process of the seventh cervical vertebra (C7) and the 12<sup>th</sup> thoracic vertebra (T12). The point where perpendicular lines to the skin surface through T12 and C7 markers cross each other shapes the thoracic flexion angle [38] (Figure 2). If the thoracic flexion angle was more than  $50^\circ$ , the subject would be considered to have a hyperkyphosis posture [33].

**Figure 1.** Cervical angle (CA) and Shoulder Angle (SA)**JMR**



**Figure 2.** Thoracic Flexion Angle (TFA)

**JMR**

After uploading the images in the software, using a high precision pen with the ability to sense 1024 pressure levels, anatomical points, including C7, T12, ear tragus, and the midpoint of the shoulder joint were identified. Then, with the Kinovea software and photogrammetric technique, the images were processed. The cervical, shoulder, and thoracic flexion angles were measured individually. Kinovea is a reliable tool that generates valid data and calculates the acceptable level of angular and linear measurements derived from the digitization of the x and y coordinates [36]. Measurements were carried out before the first yoga class (pre-test) and after the last yoga class (post-test). Data were analyzed using IBM SPSS version 20. Descriptive statistics were calculated for all variables. The Shapiro-Wilk test was used to test normality given the small sample size. The paired-sample t test was used to compare the within-group differences. The level of statistical significance was set at  $P < 0.05$ .

### 3. Results

**Table 3.** Demographic characteristics of the participants

Variables	Intervention Group (n=15)
	Mean±SD
Age (y)	53.1±6.47
Height (cm)	159.2±6.84
Weight (kg)	66.3±8.46
BMI ( $\text{kg}/\text{m}^2$ )	26.2±3.39

**JMR**

In this study, 20 middle-aged women with UCS were screened and five participants dropped out of the study considering the inclusion and exclusion criteria. Thus, 15 subjects performed Iyengar yoga home-based training with an emphasis on spine and shoulder exercises (Table 3).

The study data were subjected to the Shapiro-Wilk test and all the variables were found to be normally distributed with a P-value of more than 0.05. Hence we used the paired-sample t test to assess the postural changes in participants before and after 8 weeks.

In participants' posture, a significant change in the cervical ( $P < 0.001$ ), shoulder ( $P < 0.005$ ) and thoracic flexion ( $P < 0.001$ ) angles after 8 weeks of the Iyengar yoga with an emphasis on spine and shoulder exercises were noted when compared with those at baseline (Table 4).

### 4. Discussion

**Table 4.** Pre-test comparison of variables

Variables	Mean±SD		t	P
	Pre-test	Post-test		
Cervical angle	38.4±10.1	49.53±7.45	-7.48	0.001*
Shoulder angle	29.4±4.37	44.54±3.51	-17.57	0.001*
Thoracic flexion angle	66±6.54	46.51±5.33	32.85	0.001*

\* Significant (The significance level is considered to be  $P < 0.05$ ).

This study aimed to investigate the effect of the Iyengar yoga with an emphasis on spine and shoulder exercises on the cervical, shoulder, and thoracic flexion angles changes in middle-aged women with UCS. To the best of our knowledge, the present study is the first study that investigated the effect of Iyengar yoga with an emphasis on spine and shoulder exercises on UCS. The results of the current study showed that the implementation of eight weeks of the Iyengar yoga spine and shoulder training program had a significant effect on the cervical, shoulder, and thoracic flexion angles in middle-aged women with UCS. Few studies investigated the effects of training programs on the UCS. Rajalaxmi et al. (2018) reported that yoga exercises significantly reduced forward head shift and shoulder protraction in people with UCS. In this study, the subjects received yoga Asanas, including Bhujangasana, Dhanurasana, Virabhadrasana, Eka Pada Bhekasana, Pranayama, Salabhasana, Adho Mukha Shvanasana, and Matsyendrasana. These asanas involve the lengthening of the spinal cord, and extension at the cervicothoracic joint and mobilization of the upper limb, and thoracic cage, and through these changes, they can improve posture [39].

In the present study, which was performed on middle-aged women suffering from hyperkyphosis with a mean angle of 66 degrees, 8 weeks of the Iyengar yoga with an emphasis on spine and shoulder exercises, decreased the kyphosis angle by 12 degrees. The current study result is consistent with previous studies results [23-27]. Greendale et al. (2009) proved a substantial decrease in the kyphosis angle in people participating in yoga classes for 24 weeks, compared to people not participating in these classes [23]. The improvement in spinal posture as the result of yoga is also confirmed by the investigation of children's and adults' posture [24, 25, 27]. Increment in strength and flexibility and enhanced awareness of posture may be the mechanisms by which postural improvements occurred [26]. Another explanation for the decrease in kyphosis angle is that most yoga asanas are performed on the elongated spine. Elongation of the

spine could reduce pelvic anteversion and lumbar lordosis, and because of the chain reaction in spinal curvatures, the kyphosis angle of the thoracic area will reduce [24].

In the current study, the forward head angle was measured as an angle drawn between a line from the tragus of the ear to the last cervical vertebra and the horizontal line [40]. In this method of measurement, the participants with forward head posture have a smaller angle relative to people with normal head posture. Based on the results of the present study, the cervical angle increased by 11 degrees after eight weeks of the Iyengar yoga with an emphasis on the spine and shoulder. This finding agrees with previous investigations [41, 42]. Ghiasinezhad et al. (2016) found that eight weeks of yoga exercises significantly improved neck proprioception, motor control, and cervical posture of 15-17 years old girls [41].

In another study, Harman et al. (2005) found that 10 weeks of targeted and progressive home exercise program improved postural alignment related to FHP [42]. A probable reason for the effectiveness of the Iyengar yoga program is that the Iyengar yoga with an emphasis on spine and shoulder exercises program focused on balancing FHP-related major muscle groups, including stretching of the cervical extensors and pectoral muscles and strengthening of the deep cervical flexors, and shoulder retractors. Another explanation for the effect of the Iyengar yoga program on forward head posture is that yoga could improve proprioception and motor control, hence cervical spine posture [41].

In the current study, the shoulder angle was determined by measuring the angle between the horizontal line passing through the spinal process of the C7, and the line attached the midpoint of the shoulder to the spinal process of C7. With this way of angle calculation, the participants with rounded shoulder posture have a smaller angle. The results showed that after 8 weeks of the Iyengar yoga with an emphasis on spine and shoulder exercises, the angle of the shoulder in women increased by

15 degrees, suggesting that the movements trained the tissues that contribute to improved posture. This finding is consistent with previous studies [43, 44]. Complex training has been shown to ameliorate postural deviations, namely, FHP and RSP [44].

Lynch et al. (2010) reported improvement in postural alignment in swimmers with round shoulder posture following an 8-week intervention [43]. Considering that rounded shoulder is a multifactorial deformity that may be affected by hypomobility of any segment of the spine, rib articulation, or joint of the shoulder girdle, including the acromioclavicular, sternoclavicular, and shoulder joints, as well as shortness of the muscles related to any segment of the spine, ribcage, and shoulder girdle [45], Iyengar yoga with an emphasis on spine and shoulder exercises could improve posture by addressing hypomobility and shortness.

This study has several limitations. A control group was not included in the current study. The sample size was small and this work should be done with a larger sample size. The yoga exercises are designed to be performed both at home and under supervision of coach at the gym, and greater changes could likely occur if they were performed under the full supervision of a coach and or corrective exercise specialist. The present study did not investigate the durability of postural changes and it remains to be determined how long the effects would continue.

## 5. Conclusion

The results of the present study showed that the implementation of eight weeks of the Iyengar yoga with an emphasis on spine and shoulder exercises had a significant effect on the cervical, shoulder, and thoracic flexion angles in middle-aged women with UCS. Generally, maintaining an active and healthy lifestyle through yoga exercise can be a key element in correcting the UCS in these women. The Iyengar yoga is a style of Hatha yoga with key aspects of correct alignment of the body and sequencing of the postures. It seems that Iyengar yoga with an emphasis on spine and shoulder exercises can correct UCS. It is a safe and cheap intervention to use in the prevention and rehabilitation area, too.

## Ethical Considerations

### Compliance with ethical guidelines

All ethical principles were considered in this article. The participants were informed about the purpose of the research and its implementation stages; they were also

assured about the confidentiality of their information. Moreover, they were allowed to leave the study whenever they wish, and if desired, the results of the research would be available to them.

## Funding

The present article was extracted from the MSc. thesis of the first author, Department of Physical Education and Sport Sciences, Tehran Science and Research Branch, Islamic Azad University.

## Authors contributions

All authors contributed in preparing this article.

## Conflict of interest

The authors declared no conflict of interest.

## References

- [1] Oberoi H, Kaur H. Effect of cervicothoracic taping along with stretching strengthening exercise program for upper cross syndrome [PhD. dissertatio]. Punjab: Lovely Professional University; 2015. <http://localhost:8080/xmlui/handle/123456789/3792>
- [2] Peterson J. Teaching Pilates for postural faults, illness and injury: A practical guide, 1<sup>st</sup> edition. Philadelphia: Butterworth Hememann Elsevier; 2009.
- [3] Peterson-Kendall F, Kendall-McCreary E, Geise-Provance P, McIntyre-Rodgers M, Romani W. Muscles testing and function with posture and pain. Philadelphia: Lippincott Williams & Wilkins, Ltd; 2005.
- [4] Page P, Frank C, Lardner R. Assessment and treatment of muscle imbalance: The Janda approach, 1<sup>st</sup> edition. Human Kinetics, Inc.; 2009.
- [5] Thigpen CA, Padua DA, Michener LA, Guskiewicz K, Giuliani C, Keener JD, et al. Head and shoulder posture affect scapular mechanics and muscle activity in overhead tasks. *Journal of Electromyography and Kinesiology*. 2010; 20(4):701-9. [DOI:10.1016/j.jelekin.2009.12.003] [PMID]
- [6] Ruivo RM, Pezarat-Correia P, Carita AI. Effects of a resistance and stretching training program on forward head and protracted shoulder posture in adolescents. *Journal of Manipulative and Physiological Therapeutics*. 2017; 40(1):1-10. [DOI:10.1016/j.jmpt.2016.10.005] [PMID]
- [7] Thacker D, Jameson J, Baker J, Divine J, Unfried A. Management of upper cross syndrome through the use of active release technique and prescribed exercises [PhD. dissertation]. United States: Logan College of Chiropractic. 2011. <https://>

- [pdfs.semanticscholar.org/ed40/d7b401417169d724af4770d-5cc8812bf9be0.pdf](https://pdfs.semanticscholar.org/ed40/d7b401417169d724af4770d-5cc8812bf9be0.pdf)
- [8] Clark M, Lucett S. *NASM essentials of corrective exercise training*. Philadelphia: Lippincott Williams & Wilkins; 2010.
- [9] Moore MK. Upper crossed syndrome and its relationship to cervicogenic headache. *Journal of Manipulative and Physiological Therapeutics*. 2004; 27(6):414-20. [DOI:10.1016/j.jmpt.2004.05.007] [PMID]
- [10] Yoo WG, Yi CH, Kim MH. Effects of a ball-backrest chair on the muscles associated with upper crossed syndrome when working at a VDT. *Work*. 2007; 29(3):239-44. <https://yonsei.pure.elsevier.com/en/publications/effects-of-a-ball-backrest-chair-on-the-muscles-associated-with-u>
- [11] Mujawar JC, Sagar JH. Prevalence of upper cross syndrome in laundry workers. *Indian Journal of Occupational and Environmental Medicine*. 2019; 23(1):54-6. [DOI:10.4103/ijoom.IJOEM\_169\_18] [PMID] [PMCID]
- [12] Verma SL, Shaikh J, Mahato RK, Sheth MS. Prevalence of forward head posture among 12-16 year old school going students: A cross sectional study. *Applied Medical Research*. 2018; 4(2):18-21. [DOI:10.5455/amr.20180805064752]
- [13] Gh ME, Alilou A, Ghafurinia S, Fereydounnia S. Prevalence of faulty posture in children and youth from a rural region in Iran. *Biomedical Human Kinetics*. 2012; 4(1):121-6. [DOI:10.2478/v10101-012-0023-z]
- [14] Vakili L, Halabchi F, Mansournia MA, Khami MR, Irandoost S, Alizadeh Z. Prevalence of common postural disorders among academic dental staff. *Asian Journal of Sports Medicine*. 2016; 7(2):e29631. [DOI:10.5812/asjms.29631] [PMID] [PMCID]
- [15] Batistão MV, Moreira RdFC, Coury HJCG, Salasar LEB, Sato Td O. Prevalence of postural deviations and associated factors in children and adolescents: A cross-sectional study. *Fisioterapia em Movimento*. 2016; 29(4):777-86. [DOI:10.1590/1980-5918.029.004.ao14]
- [16] Page P. Shoulder muscle imbalance and subacromial impingement syndrome in overhead athletes. *International Journal of Sports Physical Therapy*. 2011; 6(1):51-8. [PMID] [PMCID]
- [17] Sahrman S, Azevedo DC, Van Dillen L. Diagnosis and treatment of movement system impairment syndromes. *Brazilian Journal of Physical Therapy*. 2017; 21(6):391-9. [DOI:10.1016/j.bjpt.2017.08.001]
- [18] Gillani SN, Qurat-Ul- Ain, Rehman SU, Masood T. Effects of eccentric muscle energy technique versus static stretching exercises in the management of cervical dysfunction in upper cross syndrome: A randomized control trial. *The Journal of the Pakistan Medical Association*. 2020; 70(3):394-8. [DOI:10.5455/JPMA.300417] [PMID]
- [19] Sengupta P. Health impacts of yoga and pranayama: A state-of-the-art review. *International Journal of Preventive Medicine*. 2012; 3(7):444-58. [PMID] [PMCID]
- [20] Sharan D, Manjula M, Urmi D, Ajeesh P. Effect of yoga on the Myofascial Pain Syndrome of neck. *International Journal of Yoga*. 2014; 7(1):54-9. [DOI:10.4103/0973-6131.123486] [PMID] [PMCID]
- [21] Amin DJ, Goodman M. The effects of selected asanas in Iyengar yoga on flexibility: Pilot study. *Journal of Bodywork and Movement Therapies*. 2014; 18(3):399-404. [DOI:10.1016/j.jbmt.2013.11.008] [PMID]
- [22] Grabara M, Szopa J. Effects of hatha yoga exercises on spine flexibility in women over 50 years old. *Journal of Physical Therapy Science*. 2015; 27(2):361-5. [DOI:10.1589/jpts.27.361] [PMID] [PMCID]
- [23] Greendale GA, Huang MH, Karlamangla AS, Seeger L, Crawford S. Yoga decreases kyphosis in senior women and men with adult-onset hyperkyphosis: Results of a randomized controlled trial. *Journal of the American Geriatrics Society*. 2009; 57(9):1569-79. [DOI:10.1111/j.1532-5415.2009.02391.x] [PMID] [PMCID]
- [24] Grabara M. Effects of 8-months yoga training on shaping the spine in people over 55. *Biomedical Human Kinetics*. 2013; 5(1):59-64. [DOI:10.2478/bhk-2013-0009]
- [25] Grabara M, Szopa J. Effects of hatha yoga on the shaping of the antero-posterior curvature of the spine. *Human Movement*. 2011; 12(3):259-63. [DOI:10.2478/v10038-011-0028-4]
- [26] Greendale GA, McDivit A, Carpenter A, Seeger L, Huang M-H. Yoga for women with hyperkyphosis: Results of a pilot study. *American Journal of Public Health*. 2002; 92(10):1611-4. [DOI:10.2105/AJPH.92.10.1611] [PMID] [PMCID]
- [27] Savić K, Pfau D, Skorić S, Pfau J, Spasojević N. The effect of Hatha yoga on poor posture in children and the psychophysiological condition in adults. *Medicinski Pregled*. 1990; 43(5-6):268-72. [PMID]
- [28] Hajhosseini E, Norasteh A, Shamsi A, Daneshmandi H. The effects of strengthening, stretching and comprehensive exercises on forward shoulder posture correction. *Physical Treatments-Specific Physical Therapy Journal*. 2014; 4(3):123-32.
- [29] Kang J-I, Choi H-H, Jeong D-K, Choi H, Moon Y-J, Park J-S. Effect of scapular stabilization exercise on neck alignment and muscle activity in patients with forward head posture. *Journal of Physical Therapy Science*. 2018; 30(6):804-8. [DOI:10.1589/jpts.30.804] [PMID] [PMCID]
- [30] Seidi F, Rajabi R, Ebrahimi I, Alizadeh MH, Minoonejad H. The efficiency of corrective exercise interventions on thoracic hyper-kyphosis angle. *Journal of Back and Musculoskeletal Rehabilitation*. 2014; 27(1):7-16. [DOI:10.3233/BMR-130411] [PMID]
- [31] Faul F, Erdfelder E, Lang A-G, Buchner A. G\*Power 3: A flexible statistical power analysis program for the social, behavioral, and biomedical sciences. *Behavior Research Methods*. 2007; 39(2):175-91. [DOI:10.3758/BF03193146] [PMID]
- [32] Ruivo RM, Pezarat-Correia P, Carita AI. Cervical and shoulder postural assessment of adolescents between 15 and 17 years old and association with upper quadrant pain. *Brazilian Journal of Physical Therapy*. 2014; 18(4):364-71. [DOI:10.1590/bjpt-rbf.2014.0027] [PMID] [PMCID]
- [33] Rodrigues A, Romeiro C, Patrizzi L. Evaluation of thoracic kyphosis in older adult women with osteoporosis by means of computerized biophotogrammetry. *Revista Brasileira de Fisioterapia*. 2009; 13(3):205-9. [DOI:10.1590/S1413-35552009005000036]
- [34] Porto AB, Okazaki VHA. Procedures of assessment on the quantification of thoracic kyphosis and lumbar lordosis by radiography and photogrammetry: A literature review. *Journal*



- of Bodywork and Movement Therapies. 2017; 21(4):986-94. [DOI:10.1016/j.jbmt.2017.01.008] [PMID]
- [35] Watson AWS. Procedure for the production of high quality photographs suitable for the recording and evaluation of posture. *Rev Fisioter Univ São Paulo*. 1998; 5(1).
- [36] Puig-Diví A, Padullés-Riu JM, Busquets-Faciaben A, Padullés-Chando X, Escalona-Marfil C, Marcos-Ruiz D. Validity and reliability of the kinovea program in obtaining angular and distance dimensions. *Preprints 2017*; 2017100042. [DOI:10.20944/preprints201710.0042.v1]
- [37] Lewis JS, Wright C, Green A. Subacromial impingement syndrome: the effect of changing posture on shoulder range of movement. *Journal of Orthopaedic & Sports Physical Therapy*. 2005; 35(2):72-87. [DOI:10.2519/jospt.2005.35.2.72] [PMID]
- [38] Singla D, Veqar Z, Hussain ME. Photogrammetric assessment of upper body posture using postural angles: A literature review. *Journal of Chiropractic Medicine*. 2017; 16(2):131-8. [DOI:10.1016/j.jcm.2017.01.005] [PMID] [PMCID]
- [39] Rajalaxmi V, Paul J, Nithya M, Lekha SC, Likitha B. Effectiveness of three dimensional approach of schroth method and yoga on pulmonary function test and posture in upper crossed syndrome with neck Pain-A double blinded study. *Research Journal of Pharmacy and Technology*. 2018; 11(5):1835-9. [DOI:10.5958/0974-360X.2018.00341.4]
- [40] Selvaganapathy K, Rajappan R, Dee TH. The effect of smartphone addiction on craniocervical angle and depression status among university students. *International Journal of Integrative Medical Sciences*. 2017; 4(5):537-42. [DOI:10.16965/ijims.2017.118]
- [41] Ghiasinezhad S, Hadadnezhad M, Letafatkar A. The effects of eight weeks of yoga training on motor control, proprioception and forward head angle among girls diagnosed with forward head posture. *International Journal of Medical Research & Health Sciences*. 2016; 5(11):40-6.
- [42] Harman K, Hubley-Kozey CL, Butler H. Effectiveness of an exercise program to improve forward head posture in normal adults: A randomized, controlled 10-week trial. *Journal of Manual & Manipulative Therapy*. 2005; 13(3):163-76. [DOI:10.1179/106698105790824888]
- [43] Lynch SS, Thigpen CA, Mihalik JP, Prentice WE, Padua D. The effects of an exercise intervention on forward head and rounded shoulder postures in elite swimmers. *British Journal of Sports Medicine*. 2010; 44(5):376-81. [DOI:10.1136/bjism.2009.066837] [PMID]
- [44] Park H-C, Kim Y-S, Seok S-H, Lee S-K. The effect of complex training on the children with all of the deformities including forward head, rounded shoulder posture, and lumbar lordosis. *Journal of Exercise Rehabilitation*. 2014; 10(3):172-5. [DOI:10.12965/jer.140113] [PMID] [PMCID]
- [45] Wong CK, Coleman D, Song J, Wright D. The effects of manual treatment on rounded-shoulder posture, and associated muscle strength. *Journal of Bodywork and Movement Therapies*. 2010; 14(4):326-33. [DOI:10.1016/j.jbmt.2009.05.001] [PMID]

---

This Page Intentionally Left Blank

---