

Bioinformatics screening of therapeutic targets in chemoresistance human colon cancer HCT8 cell line

Zohreh Jahanafrooz ^{1*}

1

1. Department of Biology,
Faculty of Sciences, University of
Maragheh, Maragheh, Iran

*Corresponding author:
Zohreh Jahanafrooz, Ph.D.
Department of Biology, Faculty of
Sciences, University of Maragheh,
Maragheh, Iran.

Tel: +98-41-37278001523
Fax: +98-41-3727800

Email: jahanafrooz@maragheh.ac.ir
jahanafrooz2018@gmail.com

ABSTRACT

Background: Chemoresistance is still one of the main challenges in treatment of cancers, including colon adenocarcinoma (COAD). COAD is a common cancer with a high mortality. The goal of this study was to identify and evaluate the differentially expressed mRNAs (DEmRNAs) associated with both 5-fluorouracil (5-Fu) and cisplatin (DDP) resistance in human COAD cell line.

Methods: Common DEmRNAs, DEMiRNAs, and DELncRNAs were obtained from the gene expression profile GSE173606 between acquired two chemoresistance (5-Fu and DDP) and sensitive HCT8 cells. PPI network of overlapped DEmRNAs was obtained based on STRING database and Cytoscape and Gephi software. Degree algorithm in CytoHubba was used to determine hub genes. The hub genes were evaluated in TIMER2.0 and GEPIA databases. miRTarBase database was used to find DEMiRNAs which target the common DEmRNAs. Then DELncRNAs which have interaction with the selected DEMiRNAs were obtained from RNAInter database. LncRNA-miRNA-mRNA competing endogenous RNA (ceRNA) network visualized using Cytoscape software.

Results: A high number of common DEmRNAs (about 1780) in chemoresistance HCT8 cells were identified. TIMER2.0 database showed that the expression of high-expressed hub genes, including EGFR, TGFB1, ESR1, ICAM1, PECAM1, CAV1, and CCL5 has significant positive correlations with tumor infiltration of cancer-associated fibroblasts in COAD. CeRNA networks included the low-expressed mRNAs (as targets of upregulated miR-675-3p, miR-6084, and miR-1182) whose expression is also down-regulated in COAD tissues based on GEPIA database. MDM2 (as a target of downregulated miR-4635 and miR-4306 in the ceRNA network) was an upregulated gene in both chemoresistance cells and COAD tumor tissues. RNAactDrug database confirmed the association of the high expression of four mRNAs of the ceRNA network (i.e., EFNB2, F2RL2, FLT1, and ADGRF1) with decreased drug sensitivity of DDP.

Conclusion: The results of this study can offer therapeutic targets. For example, inhibition of CCL5, oncogenic miR-675-3p, and MDM2 might be a good choice for gene targeted therapy to overcome the multi-drug resistance in COAD. Moreover, it can provide multiple mRNAs and miRNAs for predicting chemoresistance COAD.

Keywords: 5-Fluorouracil, Cisplatin, Colon adenocarcinoma, Drug resistance, HCT8 cell line

INTRODUCTION:

Colon adenocarcinoma (COAD) is one of the leading cancer-dependent causes of death in both men and women worldwide. Surgery, chemotherapy, radiation therapy, targeted therapy, and immunotherapy are the common types of treatment for people with COAD [1]. 5-fluorouracil (5-Fu) and cisplatin (DDP) are two examples of chemotherapy drugs. 5-Fu by damaging the RNA and DNA leads to cell death [2]. DDP by modulation of DNA synthesis disturbs the S phase of the cell cycle and repair mechanisms leading to cell death [3]. However, a heterogeneous population of cancer cells and complex interconnection between cancer cells and noncancer cells inside the tumor microenvironment (TME) usually trigger drug resistance. Drug resistance decreases the therapeutic efficacies of chemotherapy drugs as well as increases metastasis and recurrence [4]. According to studies, resistance mechanisms can be different among cancer patients; therefore, finding common approaches for reversing drug resistance is challenging. On the other hand, cancer cells usually develop different resistance mechanisms against different chemotherapy drugs. Various drug resistance cellular mechanisms have been reported, including increased drug efflux, increased drug metabolism, and apoptosis escape [4, 5]. The mentioned modified mechanisms are because of genetic factors either mutations or transcriptome changes or both of them [6]. Moreover, multi-drug resistance happens frequently in cancers [7]; therefore, determining common mutations and differentially expressed genes (DEGs) associated with chemoresistance can be useful in overcoming multi-drug resistance. This study using gene expression profile GSE173606 data looked for genes with differential expression between 5-Fu resistance and sensitive human COAD HCT8 cell line as well as between DDP resistance and sensitive cells. This research aimed to explore the potential common hub genes related to chemoresistance because targeting the key upregulated genes could be effective in reversing the resistance to chemotherapy drugs. In addition, the correlation of hub genes with cancer-associated fibroblasts (CAFs) infiltration to TME was investigated. Overlapped mRNA-miRNA-lncRNA ceRNA networks associated with chemoresistance were constructed to find key miRNAs. The identified common genes can be used in diagnostic and therapeutic approaches.

2. Methods:

2-1 Data processing and determining DEGs

The GSE173606 data package containing the microarray data of acquired 5-Fu and DDP-resistant and sensitive HCT8 cell line from the GEO database were obtained. DEGs with $|\log_2FC| > 0$ and adjusted P value ≤ 0.05 between drug resistance and sensitive groups were identified via GEO2R online tools and R program. DEGs with $|\log_2FC| \geq 1$ were selected for further analysis.

2-2 Determining hub genes

We used STRING database [8] for construction of a protein-protein interaction (PPI) network for DEmRNAs associated with both 5-Fu and DDP resistance in the HCT8 cell line. The results were visualized with CytoScape software. To select the hub genes, CytoHubba, as a plug-in of CytoScape, was employed to identify the hub genes. Degree method as a network centrality measure was used to identify hub genes participating in PPI network. Genes with the higher interactions have the higher values (Score > 100) in Degree method and regarded as hub genes.

2-3 Evaluations of hub genes

We evaluated the correlation between upregulated hub genes and CAF infiltration. According to TIMER2.0 database [9], different methods (EPIC, XCELL, TIDE, and MCP-COUNTER) were performed for the association analysis. The Gene Expression Profiling Interactive Analysis (GEPIA) database was used for the evolution of the expression of the identified hub genes.

2-4 Construction of common ceRNA networks

Targets of common DEmiRNAs (mRNAs and lncRNAs) were determined from the miRTarBase [10] and RNAInter [11]. Overlapping of the targets with common DEmRNAs or DElncRNAs was found by Venn diagram. Finally, the mRNA-miRNA-lncRNA ceRNA network was visualized using the Cytoscape software platform. The drug sensitivity analysis of DEmRNAs was analyzed using the RNAactDrug database [12].

3. Results:

3-1 Different gene expression between resistant and sensitive cells

About 1544 genes, including 1203 mRNAs and 341 ncRNAs were high-expressed and 1900 genes, including 1272 mRNAs and 628 ncRNAs were low-expressed in 5-Fu resistance cells. In DDP resistance cells, 1328 high-

expressed and 1029 low-expressed mRNAs and 434 high-expressed and 608 low-expressed ncRNAs were detected (Fig 1). Venn diagram showed the 1780 overlapped DEMRNAs in the 5-Fu and DDP resistance cells. These common DEMRNAs and their related DEMiRNAs and DELncRNAs (with $|\log_2FC| \geq 1$ and adjusted P value ≤ 0.05) were selected for future analyses.

3-2 PPI network construction of common 5-Fu and DDP related genes and selection of hub genes

Degree algorithm in CytoHubba was used to determine the interactions. The PPI network for the top 100 genes was reconstructed by Gephi software and reanalyzed to identify the hub genes (Fig. 2). Seven overexpressed common hub genes, including EGFR, TGFB1, ESR1, ICAM1, PECAM1, CAV1, and CCL5 and five downregulated, including CD44, CXCL8, CCL2, HIF1A, and PTGS2 were determined (Table 1). CCL5 and ICAM1 were also high-expressed in COAD compared to normal tissues and CCL2 and PTGS2 were also low-expressed in COAD compared to normal tissues based on GEPIA database. However, EGFR, TGFB1, ESR1, PECAM1, and CAV1 were downregulated and CD44, CXCL8, and HIF1A were upregulated in COAD compared to normal tissues based on GEPIA database (Supplementary Figure 1).

TIMER2.0 database showed positive correlations ($0.75 \geq \text{methods}$) for all the seven high-expressed hub genes with CAFs infiltration (Fig. 3).

3-3 mRNA-miRNA-lncRNA ceRNA network associated with chemoresistance in human colon cancer HCT8 cell line

As depicted in Fig. 4, there are 8 downregulated and 9 upregulated miRNAs in common ceRNA networks of chemoresistance cells. Based on miRTarBase database, 50 high-expressed mRNAs have interaction with the 8 miRNAs and 18 low-expressed mRNAs have interaction with the 9 miRNAs. The high-expressed lncRNAs namely HOXA-AS3 can act as a competing endogenous RNA for miR-4306 and cause the upregulation of its targets.

According to RNAactDrug database, the expression of EFNB2, F2RL2, FLT1, ADGRF1, TCF4, and CCDC80 are negatively related to the drug sensitivity of DDP ($P < 0.01$). mRNAs of EFNB2, F2RL2, and ADGRF1 genes were increased in both chemoresistance cells and in COAD tissues. However, mRNAs of TCF4 and CCDC80 genes were increased in chemoresistance cells and decreased in COAD tissues based on GEPIA database (Fig.5).

4. Discussion

Although there have been reported several routes (such as small molecules or natural anticancer agents) for reversing drug resistance in cancer, chemoresistance is still an unsolved problem in cancer treatment [13, 14]. Therefore, finding the key genes and molecular mechanisms involved in drug resistance seems to be necessary. This study aimed to identify the key genes




Figure 1. Tolcano plots of GSE173606 for acquired 5-Fu and DDP resistance COAD cells obtained from GEO2R analysis. X axis indicates the $\log_2FC$ (or $\log_2$ fold change) of genes and Y axis is related to the adjusted P value which is the negative $\log_{10}$ of the adjusted P value. Red dots represent upregulated genes ($\log_2FC > 0$) and blue dots represent downregulated genes ($\log_2FC < 0$). In this study, genes with $\log_2FC \geq 1$ were considered high-expressed and genes with $\log_2FC \leq -1$ were considered low-expressed genes in chemoresistance cells.




Figure 2. PPI network construction. Venn diagram showed the 1780 overlapped DEmRNAs in the 5-Fu and DPP resistance cells. PPI network of top 100 common chemoresistance associated DEmRNAs in Gephi (version 0.1) based on Degree algorithm, genes with more interactions are depicted with the high intensity of blue color and larger circles.

Table 1. Common identified hub genes in acquired 5-Fu and DDP- resistance COAD cells based on Degree algorithm in CytoHubba.

Rank	Name	Score	Log2FC	
			5-Fu-R compared to 5-Fu-S	DDP-R compared to DDP-S
1	EGFR (epidermal growth factor receptor)	243	1.29	1.72
2	CD44 (CD44 molecule)	170	-2.69	-2.84
3	TGFB1 (transforming growth factor beta 1)	163	1.50	1.78
4	ESR1 (estrogen receptor 1)	141	3.08	2.33
5	CXCL8 (C-X-C motif chemokine ligand 8)	140	-2.45	-4.00
6	CCL2 (C-C motif chemokine ligand 2)	134	-4.16	-4.06
7	ICAM1 (intercellular adhesion molecule 1)	124	1.38	1.60
7	HIF1A (hypoxia inducible factor 1 subunit alpha)	124	-1.10	-1.07
9	PECAM1 (platelet and endothelial cell adhesion molecule 1)	119	1.10	2.04
10	PTGS2 (prostaglandin-endoperoxide synthase 2)	115	-3.5	-6.5
11	CAV1 (caveolin 1)	112	2.81	2.53
12	CCL5 (C-C motif chemokine ligand 5)	101	1.71	1.90





Figure 3. Correlation of high-expressed hub genes ($\log_2FC > 1$, P value < 0.05) with CAFs infiltration in COAD. A) According to TIMER2.0 database, different methods (EPIC, XCELL, TIDE, and MCP-COUNTER) were performed for the association analysis. B) Bar chart represents the percentages of methods showed positive correlation between upregulated hub genes and CAFs tumor infiltration for each gene.

Ye et al. [15] showed that overexpression of ESR1 or ESR- α was related to decreased chemosensitivity to 5-Fu and the present study reinforced this. The blockade of the three upregulated hub genes, namely EGFR, TGFB1, and CCL5 has been suggested as some treatment routes in COAD in previous studies [16-18]. CCL5 is a significantly high-expressed gene in COAD chemoresistance cells and COAD tissues. Overexpression of CCL5 has been associated with immune cell infiltration, drug resistance, and metastasis in several studies [19, 20]. According to the TIMER2.0 database, the upregulated hub genes (EGFR, TGFB1, ESR1, ICAM1, PECAM1, CAV1, and CCL5) can recruit CAFs to the TME milieu. The TME can have an essential role in drug resistance properties [21]. CAFs

In ceRNA networks, three upregulated miR-675-3p ($\log_2\text{FC} > 2$), miR-6084 ($\log_2\text{FC} > 1.6$), and miR-1182 ($\log_2\text{FC} > 1.2$) target mRNAs which their expression is downregulated both in chemoresistance cells and in COAD compared to normal tissues. These mRNAs include IGF1R and EHD3 (targets of miR-675-3p and miR-6084), ZNF704 (target of miR-675-3p), TNFAIP2 (a target of miR-6084), and CD1D and KLHL3 (targets of miR-1182). The oncogenic role of miR-675-3p in COAD



23-32: Vol 16 ,No 1 ,2024 ,Basic & Clinical Cancer Research




Figure 5. The molecular level of EFNB2, F2RL2, FLT1, ADGRF1, TCF4, and CCDC80 genes (which are high-expressed in chemoresistance cells) are negatively related to drug sensitivity of DDP based on RNAactDrug database ($P < 0.01$). mRNAs of EFNB2, F2RL2, and ADGRF1 genes are upregulated in COAD tissues. However, mRNAs of TCF4 and CCDC80 genes are decreased in COAD tissues based on GEPIA database.

has been confirmed previously [24]. Therefore, the knockdown of miR-675-3p might be a good choice for gene targeted therapy in chemoresistance COAD. EHD3, as a target of miR-675-3p, is frequently downregulated by several tumor types and confers lower chemotherapy response rates in COAD [25].

In the ceRNA network, MDM2 is under the control of two downregulated microRNAs (miR-4635 and miR-4306). It was reported negative association between MDM2 overexpression and chemosensitivity [26]. MDM2 is

a crucial inhibitor of the TP53 tumor suppressor [27] which is an upregulated gene in both COAD tumor tissues (according to GEPIA) and chemoresistance cells in this study. Normally, TP53 causes cell cycle arrest or apoptosis in damaged cells but most cancer cells lose this crucial brake because of many reasons such as mutations or MDM2 upregulation [28]. Blocking of TP53 through MDM2 prevents cell cycle arrest or apoptosis induction in chemotherapy induced DNA damages. MDM2 inhibitors led to TP53 reactivation and increased

chemosensitivity [26]. TLE3 as another upregulated mRNA in the ceRNA network is under the control of two downregulated microRNAs (miR-6739 and miR-4306); however, high TLE3 expression is associated with increased chemosensitivity in taxane-treated breast and ovarian cancers [29].

The expression of EFNB2, FLT1, F2RL2, and ADGRF1 was negatively correlated with the drug sensitivity of DDP according to the RNAactDrug database. Considering that the mentioned four genes are upregulated in COAD based on GEPIA, they can be also suggested as the genes in targeted therapy to overcome the chemoresistance in COAD. EFNB2 and FLT1 are overexpressed genes in several cancers and correlated with the malignant progression of tumors, including COAD. So, blocking EFNB2 and FLT1 has been suggested as alternative therapies for COAD [30,31]. Silencing of EFNB2 inhibited cell growth and migration of COAD cells [31].

5. Conclusion

In general, the common DEGs in chemoresistance cells were screened out. It was found some mRNAs and miRNAs that their manipulation could increase the drug sensitivity of COAD cells; for example, the knockdown of EGFR, TGFB1, CCL5, and miR-675-3p. EGFR, TGFB1, and CCL5, as upregulated hub genes, by recruiting the CAFs can induce immune abnormalities in TME, which is important in chemoresistance. In addition to therapeutic potential, some hub genes (e.g., EGFR, TGFB1, ESR1, PECAM1, CAV1, CD44, CXCL8, and HIF1A) can be used as chemoresistance diagnostic markers because their changes in chemoresistance cells are in the contrary of the COAD tissues (based on GEPIA). Further research is needed to confirm these results obtained from bioinformatic methods.

Conflict of Interests: The authors report there are no competing interests to declare.

Acknowledgments: This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Reference:

1. Jahanafrooz Z, Mosafer J, Akbari M, Hashemzaei M, Mokhtarzadeh A, Baradaran B. Colon cancer therapy by focusing on colon cancer stem cells and their tumor microenvironment. *J Cell Physiol.* 2020;235:4153-66. DOI: 10.1002/jcp.29337
2. Azwar S, Seow HF, Abdullah M, Faisal Jabar M, Mohtarrudin N. Recent updates on mechanisms of resistance to 5-fluorouracil and reversal strategies in colon cancer treatment. *Biology.* 2021;10:854. DOI: 10.3390/biology10090854
3. Romani AM. Cisplatin in cancer treatment. *Biochem Pharmacol.* 2022;206:115323. DOI: 10.1016/j.bcp.2022.115323
4. Chen L, Yang F, Chen S, Tai J. Mechanisms on chemotherapy resistance of colorectal cancer stem cells and research progress of reverse transformation: A mini-review. *Front Med.* 2022;9:995882. DOI: 10.3389/fmed.2022.995882
5. Haider T, Pandey V, Banjare N, Gupta PN, Soni V. Drug resistance in cancer: mechanisms and tackling strategies. *Pharmacol Rep.* 2020;72:1125-51. DOI: 10.1007/s43440-020-00138-7
6. Bukowski K, Kciuk M, Kontek R. Mechanisms of multidrug resistance in cancer chemotherapy. *Int J Mol Sci.* 2020;21:3233. DOI: 10.3390/ijms21093233
7. Ashique S, Bhowmick M, Pal R, Khatoon H, Kumar P, Sharma H, et al. Multi drug resistance in colorectal cancer-approaches to overcome, advancements and future success. *Adv Cancer Biol: Metastasis.* 2024;100114. DOI: 10.1016/j.adcanc.2024.100114
8. Szklarczyk D, Kirsch R, Koutrouli M, Nastou K, Mehryary F, Hachilif R, et al. The STRING database in 2023: protein-protein association networks and functional enrichment analyses for any sequenced genome of interest. *Nucleic Acids Res.* 2023;51:D638-D46. DOI: 10.1093/nar/gkac1000
9. Li T, Fu J, Zeng Z, Cohen D, Li J, Chen Q, et al. TIMER2. 0 for analysis of tumor-infiltrating immune cells. *Nucleic Acids Res.* 2020;48:W509-W14. DOI: 10.1093/nar/gkaa407
10. Huang HY, Lin YC, Cui S, Huang Y, Tang Y, Xu J, et al. miRTarBase update 2022: an informative resource for experimentally validated miRNA-target interactions. *Nucleic Acids Res.* 2022;50:D222-D30. DOI: 10.1093/nar/gkab1079
11. Kang J, Tang Q, He J, Li L, Yang N, Yu S, et al.

- RNAInter v4. 0: RNA interactome repository with redefined confidence scoring system and improved accessibility. *Nucleic Acids Res.* 2022;50:D326-D32. DOI: 10.1093/nar/gkab997
12. Dong Q, Li F, Xu Y, Xiao J, Xu Y, Shang D, et al. RNAactDrug: a comprehensive database of RNAs associated with drug sensitivity from multi-omics data. *Brief Bioinform.* 2020;21:2167-74. DOI: 10.1093/bib/bbz142
 13. Ye J, Wu J, Liu B. Therapeutic strategies of dual-target small molecules to overcome drug resistance in cancer therapy. *Biochim Biophys Acta Rev Cancer.* 2023;1878:188866. DOI:10.1016/j.bbcan.2023.188866
 14. Chen T, Xiao Z, Liu X, Wang T, Wang Y, Ye F, et al. Natural products for combating multidrug resistance in cancer. *Pharmacol Res.* 2024;202:107099. DOI: 10.1016/j.phrs.2024.107099
 15. Ye SB, Cheng YK, Deng R, Deng Y, Li P, Zhang L, et al. The Predictive Value of Estrogen Receptor 1 on Adjuvant Chemotherapy in Locally Advanced Colorectal Cancer: A Retrospective Analysis With Independent Validation and Its Potential Mechanism. *Front Oncol.* 2020;10:214. DOI: 10.3389/fonc.2020.00214
 16. Li X, Wu Y, Tian T. TGF- β signaling in metastatic colorectal cancer (mCRC): from underlying mechanism to potential applications in clinical development *Int J Mol Sci.* 2022;23:14436. DOI: 10.3390/ijms232214436
 17. Pabla B, Bissonnette M, Konda VJ. Colon cancer and the epidermal growth factor receptor: Current treatment paradigms, the importance of diet, and the role of chemoprevention. *World J Clin Oncol.* 2015;6:133. DOI: 10.5306/wjco.v6.i5.133
 18. Aldinucci D, Borghese C, Casagrande N. The CCL5/CCR5 axis in cancer progression. *Cancers.* 2020;12:1765. DOI: 10.3390/cancers12071765
 19. Schlechter BL, Stebbing J. CCR5 and CCL5 in metastatic colorectal cancer. *J immunotherap cancer.* 2024;12. DOI: 10.1136/jitc-2023-008722
 20. Battaglin F, Baca Y, Millstein J, Yang Y, Xiu J, Arai H, et al. Ccr5 and Ccl5 gene expression in colorectal cancer: comprehensive profiling and clinical value. *J Immunother Cancer.* 2024;12. DOI: 10.1136/jitc-2023-007939
 21. Erin N, Grahovac J, Brozovic A, Efferth T. Tumor microenvironment and epithelial mesenchymal transition as targets to overcome tumor multidrug resistance. *Drug Resist Updat.* 2020;53:100715. DOI: 10.1016/j.drug.2020.100715
 22. Feng B, Wu J, Shen B, Jiang F, Feng J. Cancer-associated fibroblasts and resistance to anticancer therapies: status, mechanisms, and countermeasures. *Cancer Cell Int.* 2022;22:166. DOI: 10.1186/s12935-022-02599-7
 23. Bu L, Baba H, Yasuda T, Uchihara T, Ishimoto T. Functional diversity of cancer-associated fibroblasts in modulating drug resistance. *Cancer Sci.* 2020;111:3468-77. DOI: 10.1111/cas.14578
 24. Yang X, Lou Y, Wang M, Liu C, Liu Y, Huang W. miR 675 promotes colorectal cancer cell growth dependent on tumor suppressor DMTF1. *Mol Med Rep.* 2019;19:1481-90. DOI: 10.3892/mmr.2018.9780
 25. Wang YH, Chang SC, Ansar M, Hung CS, Lin RK. Eps15 homology domain-containing protein 3 hypermethylation as a prognostic and predictive marker for colorectal cancer. *Biomedicines.* 2021;9:453. DOI: 10.3390/biomedicines9050453
 26. Grigoreva T, Sagaidak A, Novikova D, Tribulovich V. New Insights into Chemoresistance Mediated by Mdm2 Inhibitors: The Benefits of Targeted Therapy over Common Cytostatics. *Biomedicines.* 2024;12:547. DOI:10.3390/biomedicines12030547
 27. Elshazli RM, Toraih EA, Elgaml A, Kandil E, Fawzy MS. Genetic polymorphisms of TP53 (rs1042522) and MDM2 (rs2279744) and colorectal cancer risk: An updated meta-analysis based on 59 case-control studies. *Gene.* 2020;734:144391. DOI: 10.1016/j.gene.2020.144391
 28. Kim KM, Ahn AR, Park HS, Jang KY, Moon WS, Kang MJ, et al. Clinical significance of p53 protein expression and TP53 variation status in colorectal cancer. *BMC cancer.* 2022;22:940. DOI: 10.1186/s12885-022-10039-y
 29. Ring BZ, Murali R, Soslow RA, Bowtell DDL, Fereday S, deFazio A, et al. Transducin-Like Enhancer of

- Split 3 (TLE3) Expression Is Associated with Taxane Sensitivity in Nonserous Ovarian Carcinoma in a Three-Cohort Study. *Cancer Epidemiol Biomarkers Prev.* 2018;27:680–8. DOI: 10.1158/1055-9965.EPI-17-1101
30. Wei SC, Tsao PN, Weng MT, Cao Z, Wong JM. Flt-1 in colorectal cancer cells is required for the tumor invasive effect of placental growth factor through a p38-MMP9 pathway. *J Biomed Sci.* 2013;20:1-12. DOI: 10.1186/1423-0127-20-39
31. Li P, Chen W, Wang Y, Fu X, Wen K, Qian J, et al. Effects of ephrinB2 gene siRNA on the biological behavior of human colorectal cancer cells. *Oncol Rep.* 2015;33:758-66. DOI: 10.3892/or.2014.3633