



Use of Pesticides and Health Risks to Chinese Water Chestnut Farmers in Suphanburi Province, Thailand

Aittiphol Duangchinda¹, Chuanpit Siripaiboon¹, Srisurang Kehanak¹, Kunthida Kingsawad¹,
Sutthayot Yimpoonsap², Umarat Sirijaroonwong³, *Anurak Khruetakham⁴

1. Sirindhorn College of Public Health, Suphanburi Praboromarajchanok Institute, Ministry of Public Health, Suphanburi, Thailand
2. Faculty of Science and Technology, Phranakorn Si Ayutthaya Rajabhat University, Phranakorn Si Ayutthaya, Thailand
3. Division of Occupational Health and Safety, Faculty of Public and Environmental Health, Huachiew Chalermprakiet University, Samut Prakan, Thailand
4. Division of Research Administration and Academic Services, Kasetsart University, Chalermprakiat Sakon Nakhon Province Campus, Sakon Nakhon, Thailand

*Corresponding Author: Email: csnark@ku.ac.th

(Received 10 Nov 2022; accepted 15 Jan 2023)

Abstract

Background: We aimed to study the relationship between the use of pesticides and the health risks faced by Chinese water chestnut farmers in this country.

Methods: This cross-sectional survey was conducted among 425 farmers in Sriprachan district, Suphanburi Province, Thailand in 2021. Samples were recruited using the cluster sampling method, and data collection took place through questionnaires. The questionnaire consisted of 3 parts, 76 items, namely personal information of farmers (12 items checklist), the use of pesticides by chemical risk assessment was a 5-point rating scale (40 items), and health risk assessment exposure to pesticides was a 5-point rating scale (24 items). The content validity index for scale (S-CVI) was 0.963 and the reliability with Cronbach's alpha coefficient was 0.904. Data were analyzed using descriptive statistics, chi-square, relative risk, and 95% CI.

Results: The overall pesticide use among farmers was at a moderate level (Mean=3.26, SD=0.60). Farmers' signs and symptoms of exposure were compared to their use of pesticides. The pesticide use increased health risks by 15.57 (95% CI: 12.33 to 18.14). Hazard identification was 10.79 higher (95% CI: 8.19 to 13.40). Dose-response assessment was -16.23 higher (95% CI: -17.63 to -14.82). Exposure assessment was 11.49 higher (95% CI: 9.87 to 13.10) and the risk characteristic was -7.46 (95% CI: -8.49 to -6.44). It was statistically significant at <.001.

Conclusion: Careless and incorrect use of pesticides by Chinese water chestnut farmers can lead to health risks from exposure to toxic substances.

Keywords: Use of pesticides; Health risks; Farmers; Water chestnut farming

Introduction

Thailand has the largest cultivation area of Chinese water chestnuts, this practice is especially

well-known in Sriprachan District, Suphanburi Province (1, 2). Pesticides are widely used in the



Copyright © 2023 Duangchinda et al. Published by Tehran University of Medical Sciences.
This work is licensed under a Creative Commons Attribution-NonCommercial 4.0 International license.
(<https://creativecommons.org/licenses/by-nc/4.0/>). Non-commercial uses of the work are permitted, provided the original work is properly cited

agricultural sector to protect yield and prevent pests such as weeds, insects, shellfish, rodents, fungi, and many plant diseases. Pesticides have impacts on health and the environment (3, 4) and even put farmers and consumers at risk of poisoning (5), but farmers using them fail to recognize these problems. Thailand has one of the top three pesticide use cases in ASEAN (6), with chemical insecticides and herbicides used for monocropping in large areas like rice fields, cornfields, and Chinese water chestnut fields (7,8). Thailand's import data shows a year-on-year trend of increasing use of pesticides. In 2015, the country imported 149,458 tons of these substances; this increased to 160,667 and 197,646 tons in 2016 and 2017, respectively (9). This puts 64.1 million Thai people at risk of exposure to pesticides at a level of more than 2.6 kilograms per person per year (10). The country's report data on occupational disease shows that pesticide use according to food group T600 (ICD-10) results in a 12.37 sickness rate per 100,000 population. In this respect, farmers are found most frequently (37.3%), followed by hired workers (28.8%) (11,12). The careless use of agricultural chemicals by farmers is both acutely toxic and

chronically poisonous, affecting the nervous system and causing cancer (13,14). It is a serious health problem for farmers caused by the use of pesticides and incorrect preventive behavior of farmers (15). There are many factors related to the use of pesticides and impacts on health, such as gender, age, educational attainment, and congenital diseases (16). Factors related to the use of pesticides include chemical type, level of chemical consumption, method, period, frequency, and use of personal protective equipment (PPE) (17,18). There are additional factors regarding knowledge, attitude, and behavior regarding pesticide use (19). In addition, there is the toxicological risk assessment of chemicals such as hazard identification, assessment response to quantity, chemical exposure assessment, and risk characterization (20,21) (Fig. 1).

Given the importance of this topic, the researchers wish to conduct a study related to the use of pesticides and the health risks faced by Chinese water chestnut farmers in Suphanburi Province. In this study, principles of risk assessment for mixed chemicals applied to the use of health risks of farmers from signs and symptoms were outlined in the study (22).




Fig. 1: The toxicological risk assessment of chemicals and health risks (20,21)

Methods

Population and Sample Group

Population - using purposive selection from 5 areas where Chinese water chestnuts are planted most from a total of 9 sub-districts in Siphachan District in 2021. Overall, 812 households was one representative each (1). The sam-

ple group size was obtained by using the Cochran formula (23).

$$n = p(1 - p)z_{\alpha}^2 / e^2.$$

n = sample size

p = the population proportion

e = acceptable sampling error

The sample group consisted of 425 farmers in Siphachan District, Suphanburi Province (14°37'11"N 100°8'40"E) (Fig. 2).




Fig. 2: Location of study area (Siphachan district, Suphanburi Province)

Research Ethics

The participants filled out the questionnaires with their consent and cooperated with this research, and their personal information was given anonymously in the research.

Data Analysis

Data were analyzed by using SPSS software version 24.0 (IBM Corp., Armonk, NY, USA), with the statistical significance level set at .05. The following calculations were made:

1. Descriptive statistics - percentage, frequency, and standard deviation.
2. Chi-Square or Fisher's Exact Test, Relative Risk (RR), and 95% CI of RR using Multiple Linear Regression to analyze the relationship between the use of pesticides and health risks of the Chinese water chestnut farmers.

Results

Personal Characteristics

More than one-half of the respondents were males who were 28-74 yr old. The majority were married elementary school graduates. Their average monthly income range was 5,001-7,500 baht. Most of the respondents suffered from chronic disease, with high blood pressure being the most prevalent. Most had universal health insurance cards and did not smoke or drink. The farmers had been cultivating Chinese water chestnut for an average of 9.74 yr on an average farm size of 3.70 acres. Most of them did not hire workers to conduct pesticide spraying (Table 1).

Table 1: Personal characteristics of the respondents (n=425)

| <i>Personal characteristics</i> | <i>N</i> | <i>%</i> |
|---|----------|----------|
| Gender | | |
| Male | 245 | 57.65 |
| Female | 180 | 42.35 |
| Age (yr) | | |
| Below 40 | 158 | 37.18 |
| 40 - 49 | 60 | 14.12 |
| 50 - 59 | 118 | 27.76 |
| 60 and above | 89 | 20.94 |
| Education attainment | | |
| Elementary school (6 yr) | 222 | 52.24 |
| Secondary school or vocational certificate | 91 | 21.41 |
| Diploma or higher vocational certificate | 88 | 20.70 |
| Bachelor's degree or equivalent | 24 | 5.65 |
| Average monthly income (baht) | | |
| Less than 5,000 | 112 | 26.35 |
| 5,001 - 7,500 | 122 | 28.70 |
| 7,501 - 10,000 | 81 | 19.06 |
| 10,001 - 12,500 | 71 | 16.71 |
| More than 12,500 | 39 | 9.18 |
| (\bar{X} = 8,717.18, SD = 2,829.18, Min = 2,400, Max = 25,000) | | |
| Chronic disease | | |
| No | 162 | 38.12 |
| Yes | 263 | 61.88 |
| - Diabetes Mellitus | 84 | 31.94 |
| - Hypertension | 97 | 36.88 |
| - Hyperlipidemia | 40 | 15.21 |
| - Chronic Obstructive Pulmonary Disease | 18 | 6.84 |
| - Osteoarthritis of Knee | 20 | 7.61 |
| - Heart Disease | 4 | 1.52 |
| Right to treatment | | |
| Government official/State enterprise employee | 81 | 19.6 |
| Health insurance card | 344 | 80.94 |
| Smoking | | |
| No | 352 | 82.82 |
| Yes | 73 | 17.18 |
| Drinking | | |
| No | 370 | 87.06 |
| Yes | 55 | 12.94 |
| Duration of farming (years) | | |
| Less than 5 | 168 | 39.53 |
| 5-9 | 48 | 11.30 |
| 10- 14 | 110 | 25.88 |
| 15 and above | 99 | 23.29 |
| (\bar{X} = 9.74 yr, SD = 6.054, Min = 2, Max = 27) | | |
| Farm size (acres) | | |
| Less than 1.98 | 158 | 37.18 |
| 1.99- 3.56 | 60 | 14.12 |
| 3.57-5.53 | 118 | 27.76 |
| 5.54 and above | 89 | 20.94 |
| (\bar{X} = 3.696 acres, SD = 5.078, Min = 0.79, Max = 8.70) | | |
| Contracted pesticides spraying | | |
| No | 398 | 93.65 |
| Yes | 36 | 6.35 |

Use of Pesticides and Health Risks

According to Table 2, the respondents use pesticides at a moderate level (\bar{X} =3.26, SD=0.60). The following were also found at a moderate level: assessment of chemicals exposure/use of PPE (\bar{X} = 3.56, SD=0.66); risk characterization/use of

pesticides method (\bar{X} =3.31, SD=0.77); hazard identification/type of pesticides (\bar{X} =3.15, SD=0.20); and quantitative response assessment/amount, duration, and frequency of use of pesticides (\bar{X} =3.01, SD=0.93), respectively.

Table 2: Use of pesticides among the respondents (N = 425)

| Use of pesticides | Level of use of pesticides | | |
|--|----------------------------|------|-------------|
| | M | SD | Description |
| 1. Hazardous identification - Type of pesticides | 3.15 | 0.20 | Moderate |
| 1.1 Use of herbicides | 4.82 | 0.47 | High |
| 1.2 Use of insecticides | 4.89 | 0.33 | High |
| 1.3 Use of pesticides for shellfish and rodents | 4.09 | 0.88 | High |
| 1.4 Use of fungicide and plant disease killer | 4.85 | 0.47 | High |
| 1.5 Use of more than two types of pesticides each time | 2.01 | 1.13 | Low |
| 1.6 Switched the types of pesticides used to prevent pesticide resistance | 3.95 | 0.96 | High |
| 1.7 Mixing different pesticides are more effective than just one | 2.14 | 0.89 | Low |
| 1.8 Use of natural or biological extracts instead of chemicals such as neem juice and biological fermentation | 2.64 | 1.11 | Moderate |
| 1.9 Use of at least one of the following pesticides: paraquat (Grammoxone), glyphosate (Round Up), chlorpyrifos (Triazophos) | 1.64 | 0.85 | Low |
| 1.10 Choose pesticides that match the type of pests | 3.57 | 0.62 | Moderate |
| 2. Quantitative response assessment, e.g. quantity, period, and frequency of use of chemicals | 3.01 | 0.93 | Moderate |
| 2.1 Strictly use the number of pesticides as specified on the package label | 3.28 | 0.69 | Moderate |
| 2.2 Use of pesticides in the amount enough to spray out each time | 3.01 | 0.90 | Moderate |
| 2.3 Use of pesticides based on one experience or habit | 2.73 | 1.21 | Moderate |
| 2.4 Use of pesticides more than as specified by the package label for more effectiveness | 2.87 | 1.12 | Moderate |
| 2.5 Immediately spray pesticides when plant disease and insects are found | 3.67 | 1.43 | High |
| 2.6 Spray pesticides for not more than an hour | 2.88 | 1.75 | Moderate |
| 2.7 Spraying pesticides for more than two hours per day on average | 2.80 | 1.33 | Moderate |
| 2.8 Leave the time for spraying pesticides before harvesting as specified by the packaging label | 2.90 | 1.53 | Moderate |
| 2.9 Spray pesticides often or twice a month to prevent insects and pests | 2.48 | 1.15 | Low |
| 2.10 Spray pesticides immediately when learning that neighbor's Chinese water chestnut fields are spreading insects or disease | 3.24 | 1.34 | Moderate |
| 3. Assessment of chemical exposure, e.g. use of personal protective equipment (PPE) | 3.56 | 0.66 | Moderate |
| 3.1 Put on a complete set of PPE (hat, rubber gloves, face mask, apron skirt, chemical goggles, and boots) | 4.27 | 0.73 | High |

| | | | |
|--|------|------|----------|
| 3.2 Use a cloth to cover the nose and mouth instead of wearing a chemical mask when spraying pesticides | 3.56 | 1.09 | Moderate |
| 3.3 Use chemical goggles when preparing or mixing chemicals or pesticides | 3.31 | 0.74 | Moderate |
| 3.4 Wear rubber gloves to prevent chemical exposure when talking or pouring chemicals | 3.31 | 0.74 | Moderate |
| 3.5 Check that PPE is in good condition before use | 3.93 | 0.64 | High |
| 3.6 Wearing personal protective equipment is inconvenient for spraying pesticides | 3.30 | 0.87 | Moderate |
| 3.7 Remove PPE immediately at the workplace after spraying pesticides | 2.65 | 2.18 | Moderate |
| 3.8 Wash clothing or PPE separately from other clothing | 3.40 | 0.56 | Moderate |
| 3.9 Wear a general mask instead of a chemical mask when spraying pesticide because it is more convenient | 3.57 | 0.77 | Moderate |
| 3.10 Wash body with water for at least 15 min and take a soapy shower if exposed to spilled chemicals to prevent harm | 3.24 | 0.72 | Moderate |
| 4. Risk characterization - use of pesticides method | 3.31 | 0.77 | Moderate |
| 4.1 Read the package label carefully for instructions on how to use it, protect against danger, and cure poisoning before using | 2.91 | 0.49 | Moderate |
| 4.2 Have been trained to use pesticides properly | 2.89 | 1.53 | Moderate |
| 4.3 Strictly spray pesticides by the method as specified by the package label | 3.20 | 1.43 | Moderate |
| 4.4 Blow or suck the spray nozzle when the sprayer is clogged or damaged | 3.20 | 1.40 | Moderate |
| 4.5 Mix many kinds of pesticides in the same tank | 2.31 | 1.35 | Low |
| 4.6 Spray pesticides in the morning or afternoon when there is no bright sunlight and always stand against the wind's direction | 4.40 | 0.56 | High |
| 4.7 Spray pesticides in advance to prevent insects or pests | 3.17 | 1.00 | Moderate |
| 4.8 Make an appointment to spray pesticides together with neighboring water chestnut fields to prevent or eliminate pests and plant diseases | 3.29 | 0.65 | Moderate |
| 4.9 Do not smoke or drink alcohol while spraying pesticides | 3.15 | 1.67 | Moderate |
| 4.10 Be careful not to let chemicals get into your mouth, nose, eyes, skin, and face while using it | 4.01 | 0.93 | High |
| Total | 3.26 | 0.60 | Moderate |

Health risks

The respondents had a low level of health risks (\bar{X} =1.69, SD=0.40). Based on its details, the following was found: group I (mild), at moderate level (\bar{X} =2.60, SD =0.79), group II (moderate) at a moderate level (\bar{X} = 1.44, SD = 0.40), and group III (severe) at a low level (\bar{X} =1.05, SD=0.11).

Relationship between personal characteristics and health risks

The following variables had a statically significant relationship with health risks among the respondents: gender, age, right to treatment, smoking, drinking, duration of farming, and contracted pesticide spraying. However, marital status, educational attainment, average monthly income, chronic diseases, and farm size had no relationship with health risks (Table 3).

Table 3: Relationship between personal characteristics and health risks of the respondents (n=425)

| Personal Characteristics | Low level of health risks | Moderate level of health risks - High level of health risks | χ^2 , (P-value) |
|---|---------------------------|---|----------------------------|
| | N (%) | N (%) | |
| Gender | | | $\chi^2(b) = 45.009^{**}$ |
| Male | 175 (41.18) | 70 (16.47) | $P < 0.001$ |
| Female | 174 (40.94) | 6 (1.41) | df = 1 |
| Age ($P_{50} = 56.00$ yr) | | | $\chi^2(a) = 4.684^*$ |
| 28 – 55 | 167 (63.77) | 26 (36.17) | $P = 0.030$ |
| 56 – 74 | 182 (36.23) | 50 (63.83) | df = 1 |
| Marital status ($P_{50} =$ Single) | | | $\chi^2(b) = 0.091$ |
| Single | 41 (9.65) | 8 (1.88) | $P = 0.846$ |
| Married, widowed, divorced, Separated | 308 (72.47) | 68 (16.00) | df = 1 |
| Educational attainment ($P_{50} =$ Elementary school) | | | $\chi^2(a) = 2.550$ |
| Elementary school | 176 (41.41) | 46 (10.82) | $P = 0.110$ |
| Higher than elementary school | 173 (40.71) | 30 (7.06) | df = 1 |
| Average monthly income ($P_{50} = 8,500$ baht) | | | $\chi^2(a) = 2.086$ |
| 1,500 – 8,500 | 161 (37.88) | 42 (9.88) | $P = 0.149$ |
| 8,501 – 25,000 | 188 (44.24) | 34 (8.00) | df = 1 |
| Chronic disease | | | $\chi^2(a) = 1.644$ |
| No | 166 (39.06) | 30 (7.06) | $P = 0.200$ |
| Yes | 183 (43.06) | 46 (10.82) | df = 1 |
| Right to treatment ($P_{50} =$ Health insurance card) | | | $\chi^2(a) = 7.532^*$ |
| Health insurance card | 291 (68.47) | 53 (12.47) | $P = 0.006$ |
| Other rights | 58 (13.65) | 23 (5.41) | df = 1 |
| Smoking ($P_{50} =$ No smoking) | | | $\chi^2(b) = 292.335^{**}$ |
| No | 340 (80.00) | 12 (2.82) | $P < 0.001$ |
| Yes | 9 (2.19) | 64 (5.41) | df = 1 |
| Drinking ($P_{50} =$ No drinking) | | | $\chi^2(b) = 240.998^{**}$ |
| No | 345 (81.18) | 25 (5.88) | $P < 0.001$ |
| Yes | 4 (0.94) | 51 (12.00) | df = 1 |
| Duration of water chestnut farming ($P_{50} = 9.00$ yr) | | | $\chi^2(a) = 3.941^*$ |
| 2 - 9 | 177 (41.65) | 29 (6.82) | $P = 0.047$ |
| 10 - 27 | 172 (40.47) | 47 (11.06) | df = 1 |
| Farm size ($P_{50} = 2.77$ acres) | | | $\chi^2(a) = 0.601$ |
| 0.79 - 2.77 | 162 (38.12) | 39 (9.18) | $P = 0.438$ |
| 2.78 - 6.32 | 187 (44.00) | 37 (8.70) | df = 1 |
| Contracted pesticides spraying | | | $\chi^2(b) = 50.053^{**}$ |
| No | 335 (78.82) | 54 (12.71) | $P < 0.001$ |
| Yes | 14 (3.29) | 22 (5.18) | df = 1 |

**P-value < 0.01, *P-value <) 0.05 by Chi-Square test)
 χ^2 a) = Pearson Chi-Square, χ^2 b) = Fisher's Exact Test

Relationship between use of pesticides and health risks of the respondents

The respondents were at risk who did not have health risks. Based on its details, the use of pesticides has positive health risks including haz-

ard identification and exposure assessment. However, use of pesticides having health risks in the opposite direction included dose-response assessment and risk characterization with a statistical significance level (P -value < 0.001) (Table 4).

Table 4: Relationships between the use of pesticides and health risks of the respondents

| Use of pesticides | Unstandardized Coefficients | | Standardized Coefficients | t | Sig. | 95% Interval for B | Confidence |
|----------------------------|-----------------------------|------------|---------------------------|---------|------|--------------------|------------|
| | B | Std. Error | | | | | |
| (Constant) Reference* | 15.57 | 1.477 | | 10.311 | .000 | 12.33 | 18.14 |
| • Hazard identification | 10.79 | 1.325 | .17 | 8.148 | .000 | 8.19 | 13.40 |
| • Dose-response assessment | -16.23 | .716 | -1.18 | -22.677 | .000 | -17.63 | -14.82 |
| • Exposure assessment | 11.49 | .820 | .59 | 14.006 | .000 | 9.87 | 13.10 |
| • Risk characterization | -7.46 | .520 | -.45 | -14.36 | .000 | -8.49 | -6.44 |

*Health risks of the respondents based on signs and symptoms due to exposure to pesticides

Discussion

The use of pesticides by Chinese water chestnut farmers in Sriprachan District, Suphanburi Province had a relationship with their health risks at a moderate level ($\bar{X} = 3.26$, $SD = 0.60$) (10). Most of the farmers were at risk from the use of pesticides at a moderate level. As a whole, the health risks of the farmers in the Sriprachan District based on signs and symptoms due to exposure to pesticides were at risk 15.57 times those having no health risks with a statistical significance level (<0.001). This was consistent with the hypothesis of the set.

Use of pesticides on hazard identification: the type of pesticides had a positive relationship with health risks (10.79 times). This could be explained by the fact that most Chinese water chestnut farmers in Sriprachan District preferred the use of insecticide and herbicide. They used insecticide at a high level ($\bar{X} = 4.89$, $SD = 0.33$) and followed by herbicide ($\bar{X} = 4.82$, $SD = 0.47$) (15), which left most farmers at risk from exposure to insecticides and herbicides. Chemical toxicity depends on the chemical properties or comments of a particular chemical. Hence, it may affect health risks from signs and symptoms.

Quantitative response assessment included amount, times span, and frequency of use of pesticides having relationships with health risks in the opposite direction (-16.23 times). Health risks

due to signs and symptoms caused by the use of pesticides of the farmers' group have fewer risk factors than those without risk factors. These are considered protective factors. In other words, the farmers using pesticides in customized quantities, not as specified by the package table and those spraying immediately when insects or plant diseases are found can be observed at a high level ($\bar{X} = 3.67$, $SD = 1.43$). This is followed by those spraying pesticides immediately when finding that water chestnut fields of neighbors have an outbreak of diseases or insects ($\bar{X} = 3.24$, $SD = 1.34$). This will be more health risks from using pesticides than farmers who use chemicals at the specified rate by the package level (10) which found that farmers are at risk of using pesticides in large quantities and having long exposure are at the greatest risk. This is particularly on the overuse of chemicals.

Assessment of chemical exposure including the use of PPE has a positive relationship with health risks (11.49 times). This can be explained that most farmers completely were PPE (i.e., hat, rubber gloves, mask face, chemical apron shirt, chemical goggles, and boots) when using pesticides and it is found at a high level ($\bar{X} = 4.27$, $SD = 0.73$). This is followed by checking PPE before using it ($\bar{X} = 3.93$, $SD = 0.64$). This results in a low level of risk of chemical exposure (10), found that inappropriate wearing of PPE is at risk most. Moreover, academic data indicate that

the nature of toxicity depends on the do received a period of exposure in areas exposed to chemicals, and little self-defense of farmers. Moreover, the pesticide prevention behaviors of the farmers are at the least average mean score ($\bar{X} = 1.86$, SD = 0.41) (16).

Risk characterization - use of pesticide method has a relationship with health risk in the opposite direction (-7.46 times). The farmers have their use of pesticides method. They mostly spray pesticides in the morning or late afternoon when there is no bright sunlight (standing against the wind direction) and it is found at a high level ($\bar{X} = 4.40$, SD = 0.56). This is followed by being careful not to let the chemical enter the mouth, nose, eyes, or sleep into the skin or face when using it ($\bar{X} = 4.01$, SD = 0.93) (19) which found that farmers perceive the risks of pesticides while mixing them with bare hand and using more chemicals than the label says. More than 64.90% find that having a lot of pests is a high chance of getting the chemical into the body. They perceive this risk through shop suggestion, package label, and their self-practice method (12), found that the use of pesticides has a negative relationship with farming size and the period of farmers ($r = -0.548$). However, it does not conform to a study (10) which showed that the use of pesticide traits in the cultivation of farmers is very risky but in the positive direction 2.36 times of those not engaged in cultivation.

Conclusion

The overall use of pesticides among Chinese water chestnut farmers was at a moderate level. The use of pesticides compared to the health risk of farmers' exposure from signs and symptoms, and the overall pesticides use increased by 15.57 higher risk than that of the farmers who had no health risks. Therefore, health officials and related parties should be monitored for the impact on farmers' health with appropriate risk mitigation management to ensure the safety of using pesticides, especially for high-risk farmers.

Journalism Ethics considerations

Ethical issues (Including plagiarism, informed consent, misconduct, data fabrication and/or falsification, double publication and/or submission, redundancy, etc.) have been completely observed by the authors.

Acknowledgements

The authors would like to acknowledge Sirindhorn College of Public Health Suphanburi for providing academic support. Siprachan District Public Health Office and Suphanburi Provincial Agriculture Extension Office for to carry out this study.

Conflict of interest

The authors declare that they have no competing interests.

References

1. Suphan Buri Provincial Agriculture and Cooperatives Office)2019(. Chinese water chestnut farming. [Cited 14 September 2019]. Available from: <http://www.supban.biz/waternut.htm>.
2. Khruksing P, Thamma-apipon S) 2017(. The environment and farming system of water chestnut farmers in Suphanburi province. *Veridian E-Journal, Silpakorn University*, 10(2): 1828-1842. [In Thai]
3. Peter N, Mildred N, Alex M, Hilda O) 2018(. Pesticides use in pest management: A case study of Ewaso Narok Wetland small-scale vegetable farmers, Laikipia County, Kenya. *J Agric Ecol*, 14(2): 1-8.
4. European Parliament Committee on Development (2021).The use of pesticides in developing countries and their impact on health and the right to food. [cited 31 January 2021]. Available from: <https://www.europarl.europa.eu/cmsdata/219887/Pesticides%20health%20and%20food.pdf>.
5. Jallow F, Awadh G, Mohammed S, et al (2017). Pesticide knowledge and safety practices among farm workers in Kuwait: Results of a

- survey. *Int J Environ Res Public Health*, 14(4): 340.
6. Phuong T (2016). Pesticide use in Thailand. [Cited 9 June 2020]. Available from: <https://ap.jficc.org.tw/article/1154>
 7. Petchuay C (2017). Application of agrochemicals in the Lower Mekong Basin. *Sci Tech J, Ubon Ratchathani University*, 19(1): 111-122.
 8. Malaja E, Liber K, Morrissey C (2020). Spatial distribution of agricultural pesticide use and predicted wetland exposure in the Canadian Prairie Pothole Region. *Sci Total Environ*, 718: 134765.
 9. Ministry of Agriculture and Cooperatives. 2019. Summary of Imported Pesticides into Thailand [in Thai]. Office of Agricultural Economics, Bangkok. [Accessed 22 Dec 2019].
 10. Wichai D, Kasetsoomboon P, Sunwijit N (2018). Risk behaviors and relationship with risk of pesticide using of farmers in Lumpathao Dam Watershed Area, Chaiyaphum province. *Journal of the Seventh Diseases Prevention and Control*, Khon Kaen Province, 25(2): 22-34. [In Thai]
 11. Bureau of Occupational and Environmental Diseases (2020). Agricultural risk assessment form for farmers from exposure to pesticides. [Cited 22 December 2019]. [In Thai].
 12. Ketbumroong V, Photipim M, Prathumyo J (2020). Health situation of farmers from the use of pesticides in group T60.0, Mueang district, Nakhon Ratchasima province. *Health Research and Development Journal*. Nakhon Ratchasima Public Health Province Office, 6(2): 76-91. [In Thai]
 13. Sapbarnr R, Nata S (2014). Health symptoms related to pesticide exposure and agricultural tasks among rice farmers from Northern Thailand. *Environ Health Prev Med*, 19(1): 12-20.
 14. Asghar U, Mallk F, Javed A (2016). Pesticide exposure and human health: A review. *J Ecosys Ecograph*, S5(5): 1-4.
 15. Raksanam B, Taneepanichskul S, Siriwong W, Robson G (2012). Factors associated with pesticide risk behaviors among rice farmers in rural community, Thailand. *Environ Earth Sci*, 2(2): 32-39.
 16. Somboon S, Rawiworakul T, Wittayapun Y, Phummarak S (2018). The relationships between personal factors, Health Belief Model, and pesticide poisoning prevent behaviors from pesticide toxicity. *J Med Health Sci*, 25(3): 1-13.
 17. Thaiou T, Thiphom S, Yasaka S, Chanthorn W (2017). Factors related to behaviors on pesticide protection of pesticide spraying workers in Dong Prakhom sub-district, Phrom Phiram district, Phitsanulok province. *Diseases Control Journal*, 43(3): 293-305.
 18. The Ohio State University (2019). Pesticide protective equipment. [Cited 27 December 2020]. Available from: <https://ohioline.osu.edu/factsheet/aex-59131>.
 19. Toeihom N, Soonthornchaikul N (2017). Determinants of risk perception regarding pesticide use among farmers in Nongchang district, Uthai Thani province. *Journal of Safety and Health*, 10(37): 21-34. [In Thai]
 20. Nielson E, Ostergaard G, Larsen C (2008). *Toxicological risk assessment of chemicals: A practical guide*. 1st ed. Informa, New York: Health Care Inc.
 21. U.S. EPA - United States of Environmental Protection Agency (2014). Framework for human health risk assessment to inform decision making, risk assessment forum. Office of the Science Advisor. U.S. Environmental Protection Agency. [Cited 9 August 2020]. Available from: <http://www2.epa.gov>.
 22. Ruangwises N (2016). *Chemical health risk assessment*. Bangkok: Chulalongkorn University Printing.
 23. Cochran G (1977). *Sampling techniques*. 3rd ed. New York: John Wiley & Sons.