



Effect of Low Protein Diet on Patients with Cardiovascular Disease: A Systematic Review and Meta-Analysis

Mingtao Ding¹, *Lei Liu²

1. Department of Health Management Centre, Jinan Central Hospital, No. 105, Jiefang Road, Jinan, 250013, China
2. Department of Internal Medicine, Jinan Central Hospital, No. 105, Jiefang Road, Jinan, 250013, China

*Corresponding Author: Email: liulei19810526@163.com

(Received 14 Oct 2023; accepted 17 Jan 2024)

Abstract

Background: Low protein diet plays an important role in cardiovascular diseases. However, exact role is unclear so far. We aimed to find out the effect of low protein diet on patients with cardiovascular disease.

Methods: The PRISMA guidelines were followed throughout the research project until 10th Apr 2023. MeSH phrases and Boolean operators were used to search PubMed for suitable studies. The entire estimate was expressed as a 95% confidence interval around the mean difference. The model was picked because of the discrepancies found in the research. Choi's Q test and I² statistics were used to determine the degree of variation between experiments. The funnel plot was used to qualitatively examine the publishing bias.

Results: Low-protein diets have a greater impact on waist circumference [-8.82 (-9.51, -8.13), $P < 0.0000.1$] and high-density lipoprotein (HDL) [-0.05 (-0.07, -0.03), $P < 0.0000.1$] alteration than non-LPD diets, as measured by the standard mean difference (SMD). Further, significant changes were observed in weight loss [1.51 (1.25, 1.77), $P < 0.00001$], BMI, [0.46 (0.25, 0.67), $P < 0.0001$], change systolic [2.48 (1.20, 3.77), $P = 0.0002$] and diastolic blood pressure [1.49 (0.72, 2.26), $P = 0.0002$], low density lipoprotein [0.09 (0.06, 0.12), $P < 0.00001$], triglyceride [0.52 (0.49, 0.55), $P < 0.00001$], in non-LPD group as compared to LPD group.

Conclusion: The results indicated the role of low protein diet on patients with cardiovascular disease.

Keywords: Low protein diet; Cardiovascular diseases; Systematic review; Meta-analysis; Blood pressure; High-density lipoprotein (HDL)

Introduction

Diet has recently been recognised as a useful measure in avoiding the onset or progression of cardiovascular illnesses (1). In animals, dietary proteins are required for the formation of all amino acids, hence appropriate protein consumption is critical (2). The minimum recommended dietary allowance (RDA) for protein in the general population is 0.8 g/kg of body weight, re-

gardless of age or gender (3). Patients with cardiovascular illnesses, on the other hand, may benefit from a higher protein intake since they have a higher protein need due to anabolic resistance and lower muscle perfusion. Nonetheless, due to physical impairments, socioeconomic constraints, and comorbidities, they frequently consume less protein. This disparity in demand and supply may



worsen the clinical outcomes of individuals suffering from cardiovascular illnesses (4-9).

The effects of a very low carbohydrate diet on body composition and cardiovascular risk factors were also studied in a randomised controlled trial. Results from a randomised controlled trials suggest that very low carbohydrate diets are more effective than low-fat diets for rapid weight loss (10). The mechanism of low protein diets on cardiovascular diseases is unclear so far. A low-protein diet raises energy intake and expenditure at the same time. Hyperphagia brought on by a low-protein diet may be mediated by elevated levels of ghrelin and fibroblast growth factor-21 (FGF21) in the bloodstream as well as hypothalamic neuropeptide Y (NPY) expression.

Abete et al. evaluated the impact of four low-calorie diets, each with a distinct dietary distribution or high protein content, on the metabolic alterations and mitochondrial oxidation that accompany weight reduction. The use of legumes or foods with a high protein content within a diet low in calories could activate mitochondrial oxidation, which could involve additional benefits in addition to those related with weight loss (11). According to the findings of the research, an increase in the consumption of whey protein did not result in a statistically significant change in the amount of weight lost or total fat reduction (12). People who suffer from cardiovascular disease do not yet have a clear picture of what the long-term repercussions of a diet low in protein will be.

Therefore, to investigate the effect of a low protein diet on cardiovascular illnesses, we did a comprehensive review and meta-analysis of randomised controlled trials (RCTs).

Methods

Search strategy

This study followed the PRISMA [Preferred Reporting Items for Systematic Review and Meta-analysis] guidelines. A thorough systematic search in PubMed was carried out utilising Medical Sub-

ject Headings (MeSH) and Boolean operators since from inception to 30th Apr 2023.

Study selection

The PICOS was considered as: Patients: Cardiovascular diseases; Intervention: low protein diet; Comparator: non-low protein diet, Outcomes: weight, body mass index (BMI), total cholesterol (TC), low-density lipoprotein cholesterol (LDL-C), high-density lipoprotein cholesterol (HDL-C), and triglyceride (TG) levels, systolic blood pressure (SBP), and diastolic blood pressure (DBP). The food and anti-oxidants supplements and physical activity were considered as confounders. Following the exclusion of duplicate publications, two writers independently assessed study titles, abstracts, or full text to identify relevant articles. Finally, original papers were considered for inclusion in the current meta-analysis if they met the following criteria: 1) were randomized clinical trials studies; 2) administered diet as an intervention 3) enrolled adult participants (Age 18 yr and older); and 4) reported a variety of risk factors for cardiovascular disease, including weight, body mass index (BMI), total cholesterol (TC), low-density lipoprotein cholesterol (LDL-C), high-density lipoprotein cholesterol (HDL-C), and triglyceride (TG) levels, systolic blood pressure (SBP), and diastolic blood pressure (DBP). Studies that did not include a suitable comparator group were not included since they did not meet the requirements. Case reports, animal studies, review papers, and meta-analyses that had already been published were not considered.

Data extraction

MD assessed the data, and any differences were handled by LL. Researchers extracted data on the study's author, publication year, country, number of case and placebo groups, participants' gender, mean age (year), mean body mass index, diabetes type, diabetes duration, protein intake in the intervention and control groups, clinical trial design, and means and standard deviations (SD) of various cardiovascular disease factors.

Quality assessment

MD checked for the possibility of bias using the Cochrane Risk of Bias Tool for Randomised Controlled Trials. Random sequence generation adequacy, allocation concealment, blinding, identification of missing outcome data, selective outcome reporting, and other potential sources of bias were all taken into account by the quality assessment method. Each domain's risk of bias was rated as "Low," "High," or "Unclear" according to criteria laid out in the Cochrane Handbook. Any discrepancies in the data extraction or assessment of the possibility of bias were settled by a LL.

Statistical analysis

RevMan V.5.3 software was used for the statistical analysis. The overall estimate was computed as a mean difference with a 95% confidence range. The model was chosen based on differences across research. The Chochrane Q test and I² statistics were used to calculate study heterogeneity. The funnel plot was used to examine the publishing bias subjectively.

Results

A flowchart of the research selection procedure including exclusion criteria (Fig. 1).




Fig. 1: Flowchart of the process of the study selection.

This figure contains 594 publications obtained from the aforementioned electronic databases.

After removing duplicate studies, 456 papers remained. These publications were then reviewed

by study title/abstract, with 408 articles being rejected because they did not fit the inclusion criteria. During the secondary screening, full-text retrieval yielded 48 publications. Eleven of those experiments were abandoned for various reasons. Overall, 36 research (12-47) investigating the effect of a low protein diet on patients with cardiovascular disease were judged to be relevant for the meta-analysis. Fig. 1 depicts the study selection as a PRISMA flow chart.


Study characteristics

The characteristics of included studies were compiled in Table 1 (Supplementary material).

Meta-analysis

Effect of low protein diet on Weight loss

The overall estimate measure in terms of standard mean difference was found to be 1.51 [1.25, 1.77] which indicate significant effect of non-low protein diet on the weight loss as compared to low protein diet as shown in Fig. 2a. The heterogeneity among studies was found to be 81% as indicated by I^2 statistics. The symmetrical shape of funnel plot indicated a less involvement of publication bias (Fig. S1).



a)




Fig. 2: Forest plot showing effects of low protein diet (LPD) as compared to non-low protein diet on a) weight loss b) BMI c) Waist circumference

Effect of low protein diet on Body Mass Index (BMI)

The overall standard mean difference (SMD) was found to be 0.46 [0.25, 0.67] which indicates significant effect of non-LPD on BMI as compared to the LPD group (Fig. 2b). The heterogeneity among studies was found to be 19%. There is an involvement of publication bias as indicated by funnel plot (Fig. S2).

Waist circumference


The overall estimate measure in terms of standard mean difference was found to be -8.82 [-9.51, -8.13] which indicates significant effect of low protein diet on the waist circumference as compared to non-LPD (Fig. 2c). However, the heterogeneity among studies was found to be very high as indicated by I² statistics. Fig. S3 indicated an involvement of publication bias.

Change in systolic blood pressure


The effect of low protein diet on change in blood pressure was also checked. The overall estimate was found to be 2.48 [1.20, 3.77] which indicates significant change in systolic blood pressure in non-LPD group as compared to LPD group as shown in Fig. 3a. There is an involvement of publication bias (Fig. S4).

Change in diastolic blood pressure

The overall estimate measure in terms of standard mean difference was found to be 1.49 [0.72, 2.26] which indicates significant effect of non-LPD as compared to LPD on change in diastolic blood pressure. Further, no heterogeneity among studies was found (Fig. 3b) and less involvement of publication bias (Fig. S5).



a)



b)


Fig. 3: Forest plot showing effects of low protein diet (LPD) as compared to non-low protein diet on a) change in systolic blood pressure b) Change in diastolic blood pressure

Change in total cholesterol


The overall estimate measure in terms of standard mean difference was found to be 0.04 [-0.03, 0.11] which indicate non-significant effect of low protein diet on change in total cholesterol. The heterogeneity among studies was found to be 58% as indicated by I² statistics (Fig. 4a). Fig. S6 represents somewhat symmetrical shape of funnel plot which indicated less involvement of publication bias.

Change in high density lipoprotein


The overall estimate measure in terms of standard mean difference was found to be -0.05[-0.07, -0.03] which indicate significant effect of low protein diet on change in high density lipoprotein. The heterogeneity among studies was found to be 73% as indicated by I² statistics (Fig. 4b). The funnel plot has indicated high involvement of publication bias (Fig. S7).



a)



b)



c)

Fig. 4: Forest plot showing effects of low protein diet (LPD) as compared to non-low protein diet on a) Change in total cholesterol. b) high density lipoprotein c) change in low density lipoprotein

Change in low density lipoprotein

The overall estimate measure in terms of standard mean difference was found to be 0.09 [0.06, 0.12] which indicate significant effect of non-low protein diet on change in low density lipoprotein. The heterogeneity among studies was found to be 45% as indicated by I² statistics (Fig. 4c). The funnel plot has indicated high involvement of publication bias (Fig. S8).


Change in C-reactive protein

The overall estimate measure in terms of standard mean difference was found to be 0.20 [-0.25, 0.64] non-which indicate significant effect of low


protein diet on change in CRP. The heterogeneity among studies was found to be 1% as indicated by I² statistics (Fig. 5a). Publication bias was not assessed due to availability of limited number of studies.

Changes in Triglyceride

The significant change in triglyceride was found in non-LPD as compared to LPD group as indicated by overall mean difference i.e. 0.52 [0.49, 0.55] (Fig. 5a). However, very high heterogeneity among studies was observed (Fig. 5b). Fig. S9 represents asymmetrical shape of funnel plot which indicated involvement of publication bias.



a)



b)

Fig. 5: Forest plot showing effects of low protein diet (LPD) as compared to non-low protein diet on a) Change in CRP b) change in triglyceride

Discussion

Heart disease is the leading killer worldwide. In 2019, CVDs were responsible for an estimated 32.5% of all deaths worldwide, or 17.9 million people. It is anticipated that cardiovascular diseases claimed the lives of 17.9 million people worldwide in 2019, making up 32% of all deaths (48). The majority of these deaths (85%) were caused by cardiovascular events such as heart attacks and strokes. The vast majority of cardiovascular diseases can be avoided by addressing behavioural risk factors such as smoking, eating a bad diet, being overweight, not being physically active enough, and drinking too much alcohol. In the present investigation, we conducted a thorough literature search and meta-analysis to investigate whether or not a low-protein diet is associated with an increased risk of cardiovascular disease. Sohoulou et al. have conducted a systematic review and meta-analysis of randomized-controlled trials evaluating the influence of low protein and high protein diets on cardiovascular risk factors and kidney function in diabetic nephropathy. These researchers looked at the

trials to determine which diet was more beneficial. Urine urea and HbA1c levels both dropped noticeably when participants switched to a diet that contained much less protein (49). Zhu et al. carried out an exhaustive analysis of randomised controlled trials in order to assess whether or not a low-protein diet is effective in the treatment of diabetic nephropathy. A diet reduced in protein does not appear to be associated with an improvement in renal function in individuals who have type 1 or type 2 diabetic nephropathy (50). A systematic review and meta-analysis of controlled trials using GRADE to investigate the effect of the Portfolio dietary pattern on the primary therapeutic lipid target for the prevention of cardiovascular disease, low-density lipoprotein cholesterol (LDL-C), as well as other well-established cardiometabolic risk factors. According to the findings of the study, following the portfolio dietary pattern can lead to clinically significant changes in LDL-C levels, in addition to other well-established cardiometabolic risk factors and an estimated 10-year increase in the risk of developing CHD (51). Schwingshackl et al. have carried out a network meta-analysis in addi-

tion to a systematic review in order to examine the impact of various dietary approaches on blood pressure in individuals who are hypertensive or are at risk of developing hypertension. The Dietary Approaches to Stop Hypertension (DASH) may be the most effective dietary approach for lowering blood pressure in hypertensive and pre-hypertensive patients based on data of a high quality (52).

The term "heterogeneity" describes the variations in study findings among investigations. Being heterogeneous just means that your data is variable, which is nothing to be afraid about (53-56). The findings of this research have also demonstrated the existence of heterogeneity among studies. A funnel plot is typically utilised in order to conduct a qualitative analysis of publication bias (57), which is yet another essential factor to investigate. The findings of the most recent inquiry have pointed to the possible role of publication bias.

The studies published in English language are only considered. Further, the only one database i.e. PubMed was searched for relevant studies. The heterogeneity among studies was found to be a little bit high.

Conclusion

The results of the current study have indicated the beneficial effects of low protein diet on cardiovascular disorders. It is advised that further research be conducted in order to throw light on these findings.

Journalism Ethics considerations

Ethical issues (Including plagiarism, informed consent, misconduct, data fabrication and/or falsification, double publication and/or submission, redundancy, etc.) have been completely observed by the authors.

Conflict of interest

The authors declare that there is no conflict of interests.

Data availability

All supplementary data not published here will be sent to the respected readers for reasonable application. Please contact the corresponding author.

References

1. Kromhout D (2001). Diet and cardiovascular diseases. *J Nutr Health Aging*, 5(3): 144-149.
2. Moughan PJ (2012). Dietary protein for human health. *Br J Nutr*, 108 Suppl 2:S1-2.
3. Wolfe RR, Miller SL, Miller KB (2008). Optimal protein intake in the elderly. *Clin Nutr*, 27(5): 675-684.
4. Appel LJ (2003). The effects of protein intake on blood pressure and cardiovascular disease. *Curr Opin Lipidol*, 14(1): 55-59.
5. Chahoud G, Aude YW, Mehta, J L (2004). Dietary recommendations in the prevention and treatment of coronary heart disease: do we have the ideal diet yet? *Am J Cardiol*, 94(10): 1260-1267.
6. Hu FB (2005). Protein, body weight, and cardiovascular health. *Am J Clin Nutr*, 82(1 Suppl):242S-247S.
7. Forman D, Bulwer BE (2006). Cardiovascular disease: optimal approaches to risk factor modification of diet and lifestyle. *Curr Treat Options Cardiovasc Med*, 8(1):47-57.
8. Van Horn L, McCoin M, Kris-Etherton P M, et al(2008). The evidence for dietary prevention and treatment of cardiovascular disease. *J Am Diet Assoc*, 108(2):287-331.
9. Schultz WM, Kelli HM, Lisko JC Varghese T, et al (2018). Socioeconomic Status and Cardiovascular Outcomes: Challenges and Interventions. *Circulation*, 137(20):2166–2178.
10. Hu T, Mills K T, Yao L, et al (2012). Effects of low-carbohydrate diets versus low-fat diets on metabolic risk factors: a meta-analysis of randomized controlled clinical trials. *Am J Epidemiol*, 176 Suppl 7(Suppl 7):S44-54.
11. Abete I, Parra D, Martinez JA (2009). Legume-, fish-, or high-protein-based hypocaloric diets: effects on weight loss and

- mitochondrial oxidation in obese men. *J Med Food*, 12(1):100-8.
12. Aldrich ND, Reicks MM, Sibley SD, et al (2011). Varying protein source and quantity do not significantly improve weight loss, fat loss, or satiety in reduced energy diets among midlife adults. *Nutr Res*, 31(2):104-112
 13. Aude YW, Agatston AS, Lopez-Jimenez F, et al (2004). The national cholesterol education program diet vs a diet lower in carbohydrates and higher in protein and mono-unsaturated fat: a randomized trial. *Arch Intern Med*, 164(19):2141-6.
 14. Ballesteros-Pomar MD, Calleja-Fernandez AR, Vidal-Casariago A, et al (2010). Effectiveness of energy-restricted diets with different protein:carbohydrate ratios: the relationship to insulin sensitivity. *Public Health Nutr*, 13(12):2119-26.
 15. Belobrajdic DP, Frystyk J, Jeyaratnaganthan N, et al (2010). Moderate energy restriction-induced weight loss affects circulating IGF levels independent of dietary composition. *Eur J Endocrinol*, 162(6):1075-1082.
 16. Brehm BJ, Seeley RJ, Daniels SR, D'Alessio DA (2003). A randomized trial comparing a very low carbohydrate diet and a calorie-restricted low fat diet on body weight and cardiovascular risk factors in healthy women. *J Clin Endocrinol Metab*, 88(4):1617-23.
 17. Brehm BJ, Spang SE, Lattin BL, et al (2005). The role of energy expenditure in the differential weight loss in obese women on low-fat and low-carbohydrate diets. *J Clin Endocrinol Metab*, 90(3):1475-82.
 18. Brinkworth GD, Noakes M, Buckley JD, Keogh JB, Clifton PM (2009). Long-term effects of a very-low-carbohydrate weight loss diet compared with an isocaloric low-fat diet after 12 mo. *Am J Clin Nutr*, 90:23-32.
 19. Dansinger ML, Gleason JA, Griffith JL, et al (2005). Comparison of the Atkins, Ornish, Weight Watchers, and Zone diets for weight loss and heart disease risk reduction: a randomized trial. *JAMA*, 293(1):43-53.
 20. Das SK, Gilhooly CH, Golden JK, et al (2007). Long-term effects of 2 energy-restricted diets differing in glycemic load on dietary adherence, body composition, and metabolism in CALERIE: a 1-y randomized controlled trial. *Am J Clin Nutr*, 85(4):1023-30.
 21. Delbridge EA, Prendergast LA, Pritchard JE, Proietto J (2009). One-year weight maintenance after significant weight loss in healthy overweight and obese subjects: Does diet composition matter? *Am J Clin Nutr*, 90(5):1203-1214.
 22. Due A, Toubro S, Skov AR, Astrup A (2004). Effect of normal-fat diets, either medium or high in protein, on body weight in overweight subjects: a randomised 1-year trial. *Int J Obes Relat Metab Disord*, 28(10):1283-90.
 23. Dyson PA, Beatty S, Matthews DR, et al (2007). A low-carbohydrate diet is more effective in reducing body weight than healthy eating in both diabetic and non-diabetic subjects. *Diabet Med*, 24(12):1430-5.
 24. Foster GD, Wyatt HR, Hill JO, et al (2003). A randomized trial of a low-carbohydrate diet for obesity. *N Engl J Med*, 348(21):2082-90.
 25. Jenkins DJA, Wong JMW, Kendall CWC, et al (2009). The effect of a plant-based low-carbohydrate ("eco-atkins") diet on body weight and blood lipid concentrations in hyperlipidemic subjects. *Arch Intern Med*, 169(11):1046-1054.
 26. Keogh JB, Brinkworth GD, Clifton PM, et al (2007). Effects of weight loss on a low-carbohydrate diet on flow-mediated dilatation, adhesion molecules and adiponectin. *Br J Nutr*, 98(4):852-9.
 27. Labayen I, Diez N, Gonzalez A, Parra D, Martinez JA, Labayen Idoia, et al (2003). Effects of protein vs. carbohydrate-rich diets on fuel utilisation in obese women during weight loss. *Forum Nutr*, 56:168-70.
 28. Landers P, Wolfe MM, Glore S, Guild R, Phillips L, Landers Patti, et al (2002). Effect of weight loss plans on body composition and diet duration. *J Okla State Med Assoc*, 95(5):329-31.

29. Lasker DAW, Evans EM, Layman DK (2008). Moderate carbohydrate, moderate protein weight loss diet reduces cardiovascular disease risk compared to high carbohydrate, low protein diet in obese adults: A randomized clinical trial. *Nutr Metab (Lond)*, 5:30.
30. Layman DK, Evans EM, Erickson D, et al (2009). A moderate-protein diet produces sustained weight loss and long-term changes in body composition and blood lipids in obese adults. *J Nutr*, 139(3):514-21.
31. Leidy HJ, Carnell NS, Mattes RD et al (2007). Higher protein intake preserves lean mass and satiety with weight loss in pre-obese and obese women. *Obesity (Silver Spring)*, 15(2):421-9.
32. Mahon AK, Flynn MG, Stewart LK, et al (2007). Protein intake during energy restriction: effects on body composition and markers of metabolic and cardiovascular health in postmenopausal women. *J Am Coll Nutr*, 26(2):182-9.
33. Maki KC, Rains TM, Kaden VN, Raneri KR, Davidson MH, Maki KC, et al (2007). Effects of a reduced-glycemic-load diet on body weight, body composition, and cardiovascular disease risk markers in overweight and obese adults. *Am J Clin Nutr*, 85(3):724-34.
34. Morgan LM, Griffin BA, Millward DJ, et al (2009). Comparison of the effects of four commercially available weight-loss programmes on lipid-based cardiovascular risk factors. *Public Health Nutr*, 12(6):799-807.
35. Noakes M, Keogh JB, Foster PR, Clifton PM (2005). Effect of an energy-restricted, high-protein, low-fat diet relative to a conventional high-carbohydrate, low-fat diet on weight loss, body composition, nutritional status, and markers of cardiovascular health in obese women. *Am J Clin Nutr*, 81(6):1298-306.
36. Sacks FM, Bray GA, Carey VJ, et al (2009). Comparison of weight-loss diets with different compositions of fat, protein, and carbohydrates. *N Engl J Med*, 360(9):859-73.
37. Stamets K, Taylor DS, Kunselman A, et al (2004). A randomized trial of the effects of two types of short-term hypocaloric diets on weight loss in women with polycystic ovary syndrome. *Fertil Steril*, 81(3):630-7.
38. Stoernell CK, Tangney CC, Rockway SW (2008). Short-term changes in lipoprotein subclasses and C-reactive protein levels of hypertriglyceridemic adults on low-carbohydrate and low-fat diets. *Nutr Res*, 28(7):443-9.
39. Te Morenga LA, Levers MT, Williams SM, et al (2011). Comparison of high protein and high fiber weight-loss diets in women with risk factors for the metabolic syndrome: A randomized trial. *Nutr J*, 10:40.
40. Volek JS, Sharman MJ, Gomez AL, et al (2003). An isoenergetic very low carbohydrate diet improves serum HDL cholesterol and triacylglycerol concentrations, the total cholesterol to HDL cholesterol ratio and postprandial lipemic responses compared with a low fat diet in normal weight, normolipidemic women. *J Nutr*, 133(9):2756-61.
41. Volek JS, Sharman MJ, Gomez AL, et al (2004). Comparison of a very low-carbohydrate and low-fat diet on fasting lipids, LDL subclasses, insulin resistance, and postprandial lipemic responses in overweight women. *J Am Coll Nutr*, 23(2):177-84.
42. Volek JS, Sharman MJ, Gomez AL, et al (2004). Comparison of energy-restricted very low-carbohydrate and low-fat diets on weight loss and body composition in overweight men and women. *Nutr Metab*, 1(1):13.
43. Yancy WS Jr, Olsen MK, Guyton JR, et al (2004). A low-carbohydrate, ketogenic diet versus a low-fat diet to treat obesity and hyperlipidemia: a randomized, controlled trial. *Ann Intern Med*, 140(10):769-77.
44. Gardner CD, Kiazand A, Alhassan S, et al (2007). Comparison of the Atkins, Zone, Ornish, and LEARN diets for change in weight and related risk factors among overweight premenopausal women: the A TO Z Weight Loss Study: a randomized trial. *JAMA*, 297(9):969-77.
45. Johnston CS, Tjonn SL, Swan PD, et al (2004). High-protein, low-fat diets are effective for weight loss and favorably alter

- biomarkers in healthy adults. *J Nutr*, 134(3):586-91.
46. Seshadri P, Samaha FF, Stern L, et al (2005). Adipocytokine changes caused by low-carbohydrate compared to conventional diets in obesity. *Metab Syndr Relat Disord*, 3(1):66-74.
 47. Appel LJ, Sacks FM, Carey VJ, et al (2005). Effects of protein, monounsaturated fat, and carbohydrate intake on blood pressure and serum lipids: results of the OmniHeart randomized trial. *JAMA*, 294(19):2455-64.
 48. WHO. Cardiovascular diseases (CVDs). [https://www.who.int/news-room/fact-sheets/detail/cardiovascular-diseases-\(cvds\)](https://www.who.int/news-room/fact-sheets/detail/cardiovascular-diseases-(cvds))
 49. Sohoulı MH, Mirmiran P, Seraj SS, et al (2022). Impact of low-protein diet on cardiovascular risk factors and kidney function in diabetic nephropathy: A systematic review and meta-analysis of randomized-controlled trials. *Diabetes Res Clin Pract*, 191:110068.
 50. Zhu HG, Jiang ZS, Gong PY, et al (2018). Efficacy of low-protein diet for diabetic nephropathy: a systematic review of randomized controlled trials. *Lipids Health Dis*, 17(1):141.
 51. Chiavaroli L, Nishi SK, Khan TA, et al (2018). Portfolio Dietary Pattern and Cardiovascular Disease: A Systematic Review and Meta-analysis of Controlled Trials. *Prog Cardiovasc Dis*, 61(1):43-53.
 52. Schwingshackl L, Chaimani A, Schwedhelm C, et al (2019). Comparative effects of different dietary approaches on blood pressure in hypertensive and pre-hypertensive patients: A systematic review and network meta-analysis. *Crit Rev Food Sci Nutr*, 59(16):2674-2687.
 53. Srivastava R, Kumar A (2021). Use of aspirin in reduction of mortality of COVID-19 patients: A meta-analysis. *Int J Clin Pract*, 75(11):e14515.
 54. D V, Sharma A, Kumar A, Flora SJS (2021). Neurological Manifestations in COVID-19 Patients: A Meta-Analysis. *ACS Chem Neurosci*, 12: 2776-2797.
 55. Thakur M, Babu A, Khatik GL, et al (2023). Role of baricitinib in COVID-19 patients: A systematic review and meta-analysis. *World J Meta-Anal*, 11(4):125-133.
 56. Singh RK (2023). A meta-analysis of the impact on gastrectomy versus endoscopic submucosal dissection for early stomach cancer. *International Journal of Clinical Medical Research*, 1(3):11.
 57. Kumar A. Meta-analysis in Clinical Research: Principles and Procedures. 2023. Springer Nature. first edition.