



Cytotoxic Evaluation of *Daphne pontica* L. Aerial Part Extracts on Three Cancerous Cell Lines by MTT Assay

Behshad Eskandari¹, Maliheh Safavi², Seyedeh Nargess Sadati Lamardi³,
Mahdi Vazirian^{1,4*}

¹Department of Pharmacognosy, Faculty of Pharmacy, Tehran University of Medical Sciences, Tehran, Iran

²Department of Biotechnology, Iranian Research Organization for Science and Technology (IROST), Tehran, Iran

³Department of Traditional Pharmacy, School of Persian Medicine, Tehran University of Medical Sciences, Tehran, Iran

⁴Medicinal Plants Research Center, Tehran University of Medical Sciences, Tehran, Iran

Received: 3 May 2019

Revised: 12 May 2019

Accepted: 14 May 2019

Abstract

Nowadays, cancer is the second prevalent cause of mortality after cardiovascular diseases in developed and the third one in developing countries. Adverse effects of chemotherapeutic agents bring the necessity of investigating about new medications with fewer side effects. *Daphne* L. genus is one of the natural sources with valuable reported anticancer effects. This study aimed to assess the cytotoxic effect of some extracts from the aerial parts of *Daphne pontica* collected from North of Iran on cancer cell lines. Extraction of the plant material was performed by maceration (3×72 h) of 200 g of sample with petroleum ether, chloroform, ethyl acetate, and methanol, respectively. The total extract was also obtained by maceration of the sample with 80% ethanol. Different concentrations of the dried extracts were prepared to assess their cytotoxic effect by 24 h incubation of cell lines with different extracts and then MTT (dimethyl thiazolyl diphenyl tetrazolium) assay on three cancerous cell lines (MDA-MB-231, MCF-7 and T47D), performed in triplicate. IC₅₀ was then estimated from curves constructed by plotting cell survival (%) versus sample concentration (µg/ml). Results indicated that ethyl acetate fraction of *D. pontica* had the most potent cytotoxic effect in MTT assay with IC₅₀ = 977.46 µg/ml; while other fractions were weaker in toxicity (IC₅₀ > 1000 µg/ml). By comparing to potent cytotoxic effects of other *Daphne* species, it seems that the cytotoxic properties of *D. pontica* is different from other species of this genus since according to this study, no significant antineoplastic properties against the three breast cancer cell lines were determined. Further studies on other pharmacological activities of this plant are recommended.

Keywords: Cytotoxic; MTT; *Daphne pontica*; Cancer cell line

Citation: Eskandari B, Safavi M, Sadati Lamardi SN, Vazirian M. Cytotoxic Evaluation of *Daphne pontica* L. Aerial Part Extracts on Three Cancerous Cell Lines by MTT Assay. Trad Integr Med 2019; 4(2): 58-63.

*Corresponding Author: Mahdi Vazirian

Department of Pharmacognosy, Faculty of Pharmacy, Tehran University of Medical Sciences, Tehran, Iran

Tel/Fax: +98-2164121223

Email: vazirian_m@sina.tums.ac.ir

Introduction

Cancer is currently one of the most important challenges in medical sciences and despite all the advances made, no definitive treatment has been found. One of the approaches to find anticancer agents is the screening of plants to find natural compounds that have cytotoxic effects on human cancer cell lines, which has so far led to the discovery of remarkably effective therapeutic compounds in a variety of cancers including the alkaloids of *Vinca rosea* and *Taxus brevifolia*. Phytochemical studies of plants with a history of traditional or folk use for the treatment of cancer often results in the isolation of compounds with antitumor activity [1].

The genus *Daphne* belongs to the Thymelaeaceae family and has a history of therapeutic use in different countries. Various pharmacological effects have been reported from different species of this plant, such as analgesic and anti-inflammatory effects [2-4], wound healing properties [5], antimicrobial effects [6], and antioxidant effects [6,7].

Daphne pontica L., known with the local name of *Bargebooei*, is one of the *Daphne* species in Iran. The leaf infusion of *D. pontica* is used by native people in Mazandaran province to treat low back and abdominal pain, diarrhea, heartburn, and kidney infections [8]. In an in vivo study, anti-inflammatory and antinociceptive activity of ethyl acetate extract of the roots of *D. pontica* was reported [4]. Considering the cytotoxic effects of various *Daphne* species in different types of cancer cells, for example, *D. gnidium* and *D. genk-*

wa on lung cancer, and *D. mucronata* on breast and leukemia cell lines [9-18], the aim of this study was to investigate the cytotoxic effect of some extracts from the aerial part of *D. pontica* collected from North of Iran against T-47D, MDA-MB231, MCF-7 cancer cell lines.

Methods

Plant material

The aerial part (leave and stem) of the *Daphne pontica* was collected from the altitudes of Sari in Mazandaran province on June 2015 and a voucher specimen deposited in *Herbarium Ministrii Iranici Agriculturae* (Iran) (Code No. IRAN70100/1).

Extraction

The air-dried aerial parts of *D. pontica* (200 g) were powdered and extracted by maceration (3×72 h) with petroleum ether, chloroform, ethyl acetate and methanol, respectively. The total extract was also obtained by maceration of the plant with 80% ethanol. Removal of the solvents with a rotary evaporator resulted in the production of petroleum ether (PEE), chloroform (ChE), ethyl acetate (EAE) and methanol (MEE) extracts.

Cell lines and cell culture

Human breast cancer cell lines comprising MDA-MB231(ATCC^R HTB-26), MCF-7 (ATCC^R HTB-22) and T47D (ATCC^R HTB-133) cells were purchased from the National Cell Bank of Iran, Pasteur Institute, Tehran, Iran.

Determination of cell viability by MTT assay

The *in vitro* cytotoxic activity of all samples was assessed against three human breast cancer cell lines by dimethyl thiazolyl diphenyl tetrazolium (MTT; Merck, Germany) colorimetric test (19). Cancer cell lines were grown in RPMI-1640 medium supplemented with 10% FBS (fetal bovine serum protein) (Gibco BRL), 100 µg/ml streptomycin and 100 U/ml penicillin (Roche, Germany). All cell lines were cultured at 37° C in air/carbon dioxide (95:5) atmosphere.

Samples were dissolved in DMSO (dimethyl sulfoxide) and were further diluted with cell culture medium. The final DMSO concentration used was 1% of total volume of medium in all treatments, including the control group. For the MTT assay, 5×10^4 cells /wells were plated into 96-well plates (Nunc, Denmark) and incubated for 24 h before addition of the extracts. The plates were incubated overnight at 37°C with 5% CO₂. Then 5 µl of the media having various concentrations of the samples was added per well in triplicate. The plates were

then incubated for 72 h. In each plate, there were three control wells (cells without test samples) and three blank wells (the medium with 0.1% DMSO) for cell viability. After treatment, the medium was removed and 200 µl phenol red-free medium containing MTT (1 mg/ml), was added to the wells, followed by 4 h incubation at 37°C. After incubation, the culture medium was then replaced with 100 µl of DMSO and the absorbance of each well was measured by using a microplate reader at 570 nm. The cytotoxicity value for each sample was presented as IC₅₀ (the median growth inhibitory concentration) of the reagents compared with the control. IC₅₀ values were calculated by Sigmaplot 9.0 software [20].

Results

The *in vitro* cytotoxic activity of total extract (TE) and other extracts including PEE, ChE, EAE, and MEE were assessed against three human breast cancer cell lines including MCF-7, T47D, and MDA-MB-231. The IC₅₀ for the samples are represented in Table 1-3.

Table 1. *In vitro* effect of *Daphne pontica* different extracts on viability (Mean ± SD) of MDA-MB 231 cell line.

Samples ^a	(Concentration (µg/ml))				IC ₅₀ (µg/ml)
	1000	100	50	10	
PEE	107.19±11.19	82.19±7.75	86.23±10.81	71.38±11.12	>1000
ChE	59.69±12.03	75.59±10.00	78.52±3.01	77.02±6.28	>1000
EAE	48.85±7.43	87.64±14.14	104.38±11.63	89.45±9.38	977.46
MEE	111.21±4.68	88.51±15.05	77.24±5.32	85.14±4.51	>1000
TE	104.70±12.36	102.89±4.80	90.14±10.27	92.50±4.70	>1000
CON+DMSO	94.18±5.71				
DMSO	3.83±0.16				

^a Samples; PEE Petroleum ether extract, ChE: Chloroform extract, EAE: Ethyl acetate extract, MEE: Methanol extract, TE: Total extract, CON+DMSO: Control+DMSO

Table2. *In vitro* effect of *Daphne pontica* different extracts on viability (Mean±SD) of T47D cell line.

Samples ^a	(Concentration (µg/ml))				IC ₅₀ (µg/ml)
	1000	100	50	10	
PEE	54.06±4.35	79.85±6.87	90.00±8.54	68.13±7.18	>1000
ChE	63.07±1.86	115.60±4.97	96.92±13.96	92.63±3.88	>1000
EAE	59.19±2.21	120.54±12.27	97.50±14.27	93.62±11.18	>1000
MEE	94.26±8.59	107.03±12.74	73.73±9.16	73.84±10.56	>1000
TE	84.61±3.61	93.69±5.82	81.90±12.72	68.02±10.41	>1000
CON+DMSO	91.83±7.06				
DMSO	9.15±0.41				

^a Samples; PEE Petroleum ether extract, ChE: Chloroform extract, EAE: Ethyl acetate extract, MEE: Methanol extract, TE: Total extract, CON+DMSO: Control+DMSO

Table3. *In vitro* effect of *Daphne pontica* different extracts on viability (Mean±SD) of MCF-7 cell line.

Samples ^a	(Concentration (µg/ml))				IC ₅₀ (µg/ml)
	1000	100	50	10	
PEE	70.41±6.17	113.88±4.49	107.19±6.74	94.37±6.06	>1000
ChE	81.85±1.53	129.67±23.22	112.28±3.04	95.94±12.10	>1000
EAE	77.58±0.12	171.23±18.60	153.72±22.07	85.39±15.01	>1000
MEE	120.32±18.47	135.69±7.12	101.57±4.63	96.91±3.33	>1000
TE	183.18±19.59	125.95±14.88	110.43±9.63	78.76±5.13	>1000
CON+DMSO	98.76±8.34				
DMSO	9.18±0.76				

^a Samples; PEE Petroleum ether extract, ChE: Chloroform extract, EAE: Ethyl acetate extract, MEE: Methanol extract, TE: Total extract, CON+DMSO: Control+DMSO

Discussion

According to the results, among all samples, EAE showed the highest cytotoxic effect on MDA-MB231 cell line with IC₅₀ value 977.46 µg/ml and the remaining samples did not show significant cytotoxicity (IC₅₀ > 1000 µg /ml). A review of previous studies showed that the

compounds in ethyl acetate, hydroalcoholic, and chloroform extracts of other *Daphne* species had significant cytotoxic effects. Ethyl acetate extract of *D. gnidium* inhibited the growth of lung cancer cell lines (A549), liver cancer cell line (SMMC-7721) and breast cancer cell line (MCF-7) by inducing apopto-

sis after 48 hours [18]. Also, daphnane diterpenoids from *D. genkwa* have been shown to inhibit proliferation of the A549 lung cancer cell line [9].

In other studies, diterpenoids with cytotoxic effects were isolated from *D. genkwa*, *D. mucronata*, *D. mezereum* and *D. linearifolia*, such as yuanhuahine, yuanhualine, yuanhuacine, yuanhuadine, yuanhuagine [9], gnidilatinonein [12,13], mezerein [11], and bioflavonoids including 4-methylgenkwanol A, and 2-hydroxygenkwanol A [21].

Several compounds with diterpene ester and secolignans structure which had a moderate cytotoxic effect were isolated from the ethyl acetate fraction of *D. acutiloba* and *D. feddei* [10, 22]. The results of this study, which was performed on three breast cancer cell lines including T47D cell line which was evidently susceptible to progesterone [23], an estrogen-positive (MCF-7) and an estrogen-negative (MDA-MB-231) human breast cancer cell line [24], showed that the ethyl acetate extract had a poor effect on the MDA-MB-231.

Conclusion

Based on the results, the most potent effect of *D. pontica* ethyl acetate extract on MDA-MB231 cancer cell line was found, and the rest of the extracts did not show cytotoxicity on breast cancer cell lines. Future studies on pure compounds of the ethyl acetate extract as well as other cancer cell lines can be done to elucidate further biological effects of this plant.

Conflicts of Interest

The authors confirm that this article content has no conflicts of interest.

Acknowledgments

None.

References

- [1] Evans WC. Trease and Evans Pharmacognosy. 16th ed. SAUNDERS 2009; pp 416-417.
- [2] Harizi H, Chaabane F, Ghedira K, Chekir-Ghedira L. Inhibition of proinflammatory macrophage responses and lymphocyte proliferation in vitro by ethyl acetate leaf extract from *Daphne gnidium*. Cell Immunol 2011;267:94-101.
- [3] Khodadadian Z, Hassanpour-Ezatti M, Mousavi SZ, Asgarpanah J. Analgesic and anti-inflammatory potential of aerial parts of the *Daphne mucronata* Royle extract in mice: Opioid-independent action. Asian Pac J Trop Biomed 2016;6:198-201.
- [4] Kupeli E, Tosun A, Yesilada E. Assessment of anti-inflammatory and antinociceptive activities of *Daphne pontica* L. (Thymelaeaceae). J Ethnopharmacol 2007;113:332-337.
- [5] Süntar I, Küpeli Akkol E, Keles H, Yesilada E, Sarker SD, Arroo R. Efficacy of *Daphne oleoides* subsp. kurdica used for wound healing: Identification of active compounds through bioassay guided isolation technique. J Ethnopharmacol 2012;141:1058-1070.
- [6] Balkan İA, Taşkın T, Doğan HT, Deniz İ, Akaydın G, Yesilada E. A comparative investigation on the in vitro anti-inflammatory, antioxidant and antimicrobial potentials of subextracts from the aerial parts of *Daphne oleoides* Schreb. subsp. *oleoides*. Ind Crops Prod 2017;95:695-703.
- [7] Sanda MA, Zengin G, Aktumsek A, Cakmak YS. Evaluation of antioxidant potential of two *Daphne* species (*D. gnidioides* and *D. pontica*) from Turkey. Emirates Journal of Food and Agriculture [EJFA] 2015:4878-494.
- [8] Gholipour A, Ghorbani Nohooji M, Rasuli N. An Ethnobotanical Study on the Medicinal Plants of Zarm-rood Rural District of Neka (Mazandaran Province). Journal of Medicinal Plants 2014;4:101-121.
- [9] Jo S-K, Hong J-Y, Park HJ, Lee SK. Anticancer activity of novel daphnane diterpenoids from *Daphne genkwa* through cell-cycle arrest and suppression of Akt/STAT/ Src signalings in human lung cancer cells. Biomol Ther (Seoul) 2012;20:513.
- [10] Huang SZ, Zhang XJ, Li XY, Kong LM, Jiang HZ, Ma QY. Daphnane-type diterpene esters with cytotoxic and

- anti-HIV-1 activities from *Daphne acutiloba* Rehd. *Phytochemistry* 2012;75:99-107.
- [11] Kupchan SM, Baxter RL. Mezerein: antileukemic principle isolated from *Daphne mezereum* L. *Science* 1975;187:652-653.
- [12] Yazdanparast R, Moosavi M. Gnidilatimonoein from *Daphne mucronata* induced differentiation and apoptosis in leukemia cell lines. *Iran J Pharm Res* 2010;76-76.
- [13] Mianabadi M, Yazdanparas R. The Anti-metastatic Potency of Gnidilatimonoein, from *Daphne Mucronata*, Compared to Two Glycosylation Inhibitors: Castanospermine and Tunicamyci. *DARU* 2004;12:44-48.
- [14] Amirghofran Z, Miri R, Javidnia K, Davoodi M. Study of cytotoxic activity of *Daphne mucronata* Royle grown in Iran. *Iran J Med Sci* 2015;26:146-151.
- [15] Yazdanparast R, Sadeghi H. Antineoplastic effects of *Daphne mucronata*: inhibition of DNA and RNA synthesis. *Acta Med Iran* 2003;202-206.
- [16] Panahi Kokhdan E, Mianabadi M, Sadeghi H. The Effects of Two Species of *Daphne*, Betulin and Betulinic Acid on Alkaline Phosphatase Activity in Two Human Cancer Cell lines, K562 and MCF-7. *Armaghane danesh* 2014;18:900-909.
- [17] Kizaibek M, Daniar M, Li L, Upur H. Antiproliferative activity of different extracts from *Daphne altaica* Pall. on selected cancer cells. *J Med Plant Res* 2011;5:3448-3452.
- [18] Chaouki W, Meddah B, Hmamouchi M. Antiproliferative and apoptotic potential of *Daphne gnidium* L. root extract on lung cancer and hepatoma cells. *Pharmazie* 2015;70:205-210.
- [19] Mosmann T. Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays. *J Immunol Methods* 1983;65:55-63.
- [20] Alipour E, Mousavi Z, Safaei Z, Pordeli M, Safavi M, Firoozpour L. Synthesis and cytotoxic evaluation of some new [1,3]dioxolo[4,5-g]chromen-8-one derivatives. *Daru* 2014;22:41.
- [21] Malafronte N, Vassallo A, Dal Piaz F, Bader A, Braca A, De Tommasi N. Biflavonoids from *Daphne linearifolia* Hart. *Phytochem Lett* 2012;5:621-625.
- [22] Lu Y, Li X, Mu H, Huang H, Li G-P, Hu Q. Bioactive phenylpropanoids from *Daphne feddei*. *J Braz Chem Soc* 2012;23:656-660.
- [23] Yu S, Kim T, Yoo KH, Kang K. The T47D cell line is an ideal experimental model to elucidate the progesterone-specific effects of a luminal A subtype of breast cancer. *Biochem Biophys Res Commun* 2017;486:752-758.
- [24] Panjehpour M, Castro M, Klotz KN. Human breast cancer cell line MDA-MB-231 expresses endogenous A2B adenosine receptors mediating a Ca²⁺ signal. *Br J Pharmacol* 2005;145:211-218.